

Guidelines on Chronic Pelvic Pain

M. Fall (chair), A.P. Baranowski, S. Elneil, D. Engeler, J. Hughes,
E.J. Messelink, F. Oberpenning, A.C. de C. Williams

TABLE OF CONTENTS

PAGE

1.	INTRODUCTION	5
1.1	Background	5
1.1.1	Definition of pain (WHO)	5
1.1.2	Innervation of the urogenital system	6
1.1.3	References	6
1.2	Pain evaluation and measurement	7
1.2.1	Pain evaluation	7
1.2.2	Pain measurement	7
1.2.3	References	8
2.	CHRONIC PELVIC PAIN	8
2.1	Background	8
2.1.1	Introduction to urogenital pain syndromes	8
2.2	Definitions of chronic pelvic pain and terminology (Table 2)	10
2.3	Classification of chronic pelvic pain syndromes	11
	Table 1: EAU classification of chronic urogenital pain syndromes (page 9)	
	Table 2: Definitions of chronic pain terminology (page 10)	
	Table 3: ESSIC classification of types of bladder pain syndrome according to the results of cystoscopy with hydrodistension and of biopsies (page 12)	
2.4	References	12
2.5	An algorithm for chronic pelvic pain diagnosis and treatment	12
2.5.1	How to use the algorithm	12
2.6	Chronic prostate pain/chronic prostatitis associated with chronic pelvic pain syndrome (CP/CPPS)	14
2.6.1	Introduction	15
2.6.2	Definition	15
2.6.3	Pathogenesis	15
2.6.4	Diagnosis	16
2.6.5	Treatment	16
2.6.5.1	Alpha-blockers	16
2.6.5.2	Antibiotic therapy	16
2.6.5.3	Non-steroidal anti-inflammatory drugs (NSAIDs)	16
2.6.5.4	Corticosteroids	16
2.6.5.5	Opioids	17
2.6.5.6	5-alpha-reductase inhibitors	17
2.6.5.7	Allopurinol	17
2.6.5.8	Phytotherapy	17
2.6.5.9	Muscle relaxants	17
2.6.5.10	Supportive therapies	17
2.6.5.11	Surgical management	17
2.6.6	References	18
2.7	Bladder pain syndrome/interstitial cystitis (BPS/IC)	24
2.7.1	Introduction	24
2.7.2	Definition	24
2.7.3	Pathogenesis	25
2.7.4	Epidemiology	27
2.7.5	Association with other diseases	27
2.7.6	Diagnosis	27
2.7.7	Biological markers	28
2.7.8	IC in children and males	28
2.7.9	Medical treatment	28
2.7.10	Intravesical treatment	30
2.7.11	Interventional treatments	32
2.7.12	Treatments of limited efficacy and absence of recent publications	34
2.7.13	Alternative and complementary treatments	34
2.7.14	Surgical treatment	35
2.7.15	References	39

2.8	Scrotal pain	58
2.8.1	Management of different conditions	59
2.8.2	References	60
2.9	Urethral pain syndrome	61
2.9.1	Treatment	62
2.9.2	References	62
3.	PELVIC PAIN IN GYNAECOLOGICAL PRACTICE	62
3.1	Introduction	62
3.2	Clinical history	62
3.3	Clinical examination	63
3.3.1	Investigations	63
3.4	Dysmenorrhoea	63
3.5	Infection	63
3.5.1	Treatment	63
3.6	Endometriosis	63
3.6.1	Treatment	63
3.7	Gynaecological malignancy	64
3.8	Injuries related to childbirth	64
3.9	Conclusion	64
3.10	References	64
4.	NEUROLOGICAL ASPECTS	65
4.1	Physiology of the urogenital system	65
4.2	Physiology of the bladder	65
4.2.1	Bladder filling	66
4.2.2	Afferent innervation of the bladder	66
4.2.3	Efferent innervation of the bladder	66
4.2.4	Central control of micturition	66
4.2.5	Physiology of the genital organs	67
4.3	Sexual dysfunction in men and women	69
4.4	References	70
5.	NEUROGENIC CONDITIONS	73
5.1	Introduction	73
5.2	Pudendal nerve entrapment	73
5.3	Other neurogenic conditions	74
5.4	References	74
6.	PELVIC FLOOR FUNCTION AND DYSFUNCTION	74
6.1	Introduction	74
6.2	Function	74
6.3	Dysfunction	75
6.4	Myofascial trigger points	75
6.5	Therapy	75
6.6	References	76
7.	PSYCHOLOGICAL FACTORS IN CHRONIC PELVIC PAIN	77
7.1	Introduction	77
7.2	Psychological models of pain	77
7.3	Methodology	78
7.4	Psychological factors in assessment of pelvic pain	78
7.4.1	Psychological risk factors in the development of pelvic pain and adaptation to it	78
7.4.2	Anxiety	78
7.4.3	Depression and catastrophizing	79
7.4.4	Impact of pain	79
7.5	Summary: assessment recommendations	79
7.5.1	Anxiety	80
7.5.2	Depression	80
7.5.3	Sexual and physical abuse in childhood	80

7.6	Psychological factors in treatment of pelvic pain	80
7.7	References	81
8.	GENERAL TREATMENT OF CHRONIC PELVIC PAIN	84
8.1	Simple analgesics	85
8.1.1	Paracetamol	85
8.1.2	Acidic antipyretic analgesics	85
8.1.3	Guidelines for use of NSAIDs and COX2 selective agents	85
8.2	Neuropathic analgesics	86
8.2.1	Tricyclic antidepressants	86
8.2.2	Anticonvulsants	86
8.2.3	N-methyl-D-aspartate (NMDA) antagonists	86
8.2.4	Sodium channel blockers	86
8.3	Opioids	87
8.3.1	Guidelines for the use of opioid-like agents in chronic/non-acute urogenital pain	87
8.3.2	Morphine	87
8.3.3	Transdermal fentanyl	87
8.3.4	Methadone	87
8.3.5	Oxycodone	88
8.3.6	Other opioids and opioid-like agents	88
8.4	References	90
8.5	Nerve blocks	93
8.5.1	References	94
8.6	Transcutaneous electrical nerve stimulation (TENS)	94
8.6.1	Results of suprapubic TENS in BPS/IC	95
8.7	Sacral neuromodulation in pelvic pain syndromes	95
8.7.1	References	95
9.	ABBREVIATIONS USED IN THE TEXT	98

1. INTRODUCTION

1.1 Background

1.1.1 Definition of pain (WHO)

'Pain management is a necessity in the work of each physician.' F. Sauerbruch, 1936

Pain can be defined as an unpleasant sensory and emotional experience associated with either real or potential tissue damage, or it can be described in terms of such damage (1). Pain is the most common symptom of any illness. In its management, firstly, the physician needs to discover and treat the cause of the pain; secondly, to treat the pain itself, whether or not the underlying cause is treatable; and thirdly to relieve the suffering caused by the pain.

One function of the nervous system is to provide information about the occurrence of or the threat of injury. The sensation of pain, by its inherent aversive nature, contributes to this function. The response of the peripheral neural apparatus via primary sensory neurones (known as nociceptors), to noxious (injurious or potentially injurious) stimuli alerts the organism to injury (potential injury). Acute pain is an important and adaptive element of the normal nervous system. In chronic or persistent pain, the purpose of the pain is lost. Such pain often represents an aberration of neural processing.

Nociceptive or neuropathic pain. 'Pain' is used to describe all sensations that are perceived as hurting; it requires the higher centres. The causes of pain may be many. For example, pain can be nociceptive or neuropathic, with many pains having both a neuropathic and nociceptive component:

- Nociceptive pain is caused by direct stimulation of nociceptors in the periphery; peripheral inflammation may or may not be present. An example of physiological nociceptive pain is when an individual perceives pain due to hot water running over their skin resulting in the individual withdrawing from the stimulus and there is no injury. Pathological nociceptive pain, however, is often associated with tissue damage and inflammation, with inflammation having the effect of increasing the perception of pain associated with peripheral stimulation.
- Neuropathic pain is caused by a lesion to the peripheral or central nervous system.

Acute or chronic pain. Pain may also be described as either acute or chronic pain:

- Acute pathological pain has an acute onset and is short-lived, usually less than a week or so, and is associated with tissue trauma, e.g. following surgery. Transient acute pain may also be caused by acute nerve injury, e.g. local injury to the ulnar nerve from hitting the elbow. Although the mechanisms of acute and chronic pain may overlap, the mechanisms of acute pain resolve quickly in contrast to chronic pain.
- Chronic (also known as persistent) pain occurs for at least 3 months. However, the mechanisms involved are more important than the duration of the pain. Chronic pain is associated with changes in the central nervous system (CNS), which may maintain the perception of pain in the absence of acute injury. These changes may also magnify perception so that non-painful stimuli are perceived as painful (allodynia), while painful stimuli are perceived as more painful than expected (hyperalgesia).

The bladder provides a good example of how changes in the CNS affect sensory perception. An acute pain insult to the bladder can produce functional changes within the CNS, so that pain persists even after removal of the stimulus. These central functional changes may also be associated with a dysaesthetic (unpleasant sensation) response; for instance, mild distension or stimulation of the bladder by urine not normally perceived, may produce the urge to urinate. Furthermore, core muscles, including pelvic muscles, may become hyperalgesic with multiple trigger points, while other organs may also become sensitive, e.g. the uterus with dyspareunia and dysmenorrhoea, the bowel with irritable bowel symptoms. The spread of abnormal sensory responses between the organs and musculoskeletal system is a well-described consequence of the CNS changes and a crucial cause of complex chronic pelvic pains. Functional abnormalities such as urinary retention may also occur.

Chronic pain is associated with various psychological responses, partly due to the long duration of the pain and partly due to neuroplasticity of the CNS. Chronic pain inhibits feelings, emotions, thinking and reactions, while reduced mobility and inhibited physiological functions restrict social interactions and work.

Although there are established management strategies, pain is often undertreated because many clinicians have a poor understanding of the principles of pain therapy. Efforts are needed to improve this situation. When appropriate, management should be both holistic and multidisciplinary.

Deep visceral pain. There are important differences between cutaneous and deep visceral pain. Unlike cutaneous pain, deep visceral pain is diffuse and poorly localized. It may be accompanied by strong autonomic

responses, such as sweating and changes in heart rate, blood pressure and respiration. Deep visceral pain may also be produced by stimuli that are not tissue-damaging, e.g. bowel and bladder distension (2,3), and may be associated with referred pain and cutaneous and deep tissue hyperalgesia.

Modulation of pain. Pain transmission from the periphery to the higher brain centres via the spinal cord is not a simple, passive process involving exclusive pathways. The relationship between a stimulus causing pain and the way it is perceived by an individual is dramatically affected by circuitry within the spinal cord and the brain. The sensation of pain is modulated as it is transmitted upwards from the periphery to the cortex. It is modulated at a segmental level and by descending control from higher centres, with the main neurotransmitters involved being serotonin, noradrenaline and the endogenous opioids.

1.1.2 Innervation of the urogenital system

Studies on the response properties of visceral afferents from the urinary tract have highlighted the differences between nociception in the skin and viscera. Most visceral primary afferents from the bladder, urethra, reproductive and other pelvic organs are encoded for both noxious and non-noxious stimuli (4-6). Increasing afferent traffic results in a change from non-noxious sensation to noxious.

Ureter. Ureteric afferents are thinly myelinated or unmyelinated and respond to direct probing of a limited area of tissue. They can be differentiated into two groups (7):

- The first group responds to ureteral contractions and is excited by low levels of distension (average threshold 8 mmHg). They appear to encode levels of distension throughout and beyond the physiological range.
- The second group does not respond to peristaltic contractions of the ureter, but can be excited by distension with a wide range of thresholds.

Urinary bladder. Two groups of afferent fibres signal noxious stimuli in the urinary bladder, mostly non-myelinated fibres, with some myelinated fibres (4).

Graded distension of the healthy urinary bladder in humans initially gives rise to a sensation of fullness and eventually pain, as the volume of urine increases and the intravesical pressure exceeds about 25-35 mmHg (8-11). In the inflamed bladder, the sensations during bladder emptying become unpleasant and painful.

Nearly all visceral primary afferents from the bladder are small myelinated or unmyelinated fibres. Some afferents exhibit a low level of ongoing discharge when the bladder is empty. Distension excites mainly thin myelinated afferents, with pressure thresholds corresponding to levels at which humans report the first sensation of fullness. Nearly all afferents are activated by the intraluminal pressures reached during normal, non-painful micturition. The activation of a large number of initially unresponsive afferents indicates that peripheral afferent mechanisms encoding pain from pelvic viscera are highly malleable and are strongly affected by tissue state. These changes are important for signalling pain and discomfort in inflammatory conditions where there is a group of afferents that become activated by the inflammation.

Male reproductive organs. More than 95% of fibres of the superior spermatic nerve are unmyelinated, with most showing polymodal properties (i.e. responses to mechanical, chemical and thermal stimuli) (12). Myelinated and unmyelinated afferent fibres form a homogeneous group with polymodal receptors in the testis and/or epididymis. Prostaglandins sensitize the afferents to other stimuli (13).

1.1.3 REFERENCES

1. Foley KM, Posner J.B. Pain and its management. In: *Cecil Textbook of Medicine*. 18th edn. Philadelphia: WB Saunders, 1988:104-112.
2. Dubner R. Basic mechanisms of pain associated with deep tissues. *Can J Physiol Pharmacol* 1991;69(5):607-609.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1863910&dopt=Abstract&itool=iconabstr
3. Ness TJ, Gebhart GF. Visceral pain: A review of experimental studies. *Pain* 1990;41(2):167-234.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2195438&dopt=Abstract&itool=iconabstr
4. Häbler H-J, Jänig W, Koltzenburg M. Activation of unmyelinated afferent fibres by mechanical stimuli and inflammation of the urinary bladder in the cat. *J Physiol* 1990;425:545-562.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2213588&dopt=Abstract&itool=iconabstr

5. Bahns E, Ernsberger U, Jänig W, Nelke A. Functional characteristics of lumbar visceral afferent fibres from the urinary bladder and the urethra in the cat. *Pflügers Arch* 1986;407(5):510-518.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3786110&dopt=Abstract&itool=iconabstr
6. Bahns E, Halsband U, Jänig W. Responses of sacral visceral afferent fibres from the lower urinary tract, colon, and anus to mechanical stimulation. *Pflügers Arch* 1987;410(3):296-303.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3684516&dopt=Abstract
7. Cervero F, Jänig W. Visceral nociceptors: A new world order? *Trend Neurosci* 1992;15(10):374-378.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1279857&dopt=Abstract&itool=iconabstr
8. Roberts WJ, Elardo SM. Sympathetic activation of A-delta nociceptors. *Somato Res* 1985;3(1):33-44.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2999942&dopt=Abstract&itool=iconabstr
9. Seltzer Z, Devor M. Ephaptic transmission in chronically damaged peripheral nerves. *Neurology* 1979;29(7):1061-1064.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=224343&dopt=Abstract&itool=iconabstr
10. Kruger L, Perl ER, Sedivec MJ. Fine structure of myelinated mechanical nociceptor endings in cathayr skin. *J Comp Neurol* 1981;198(1):137-154.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7229137&dopt=Abstract&itool=iconabstr
11. Treede R-D, Meyer RA, Raja S N, Campbell JN. Peripheral and central mechanisms of cutaneous hyperalgesia. *Prog Neurobiol* 1992;38(4):397-421.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1574584&dopt=Abstract&itool=iconabstr
12. Kumazawa T. Sensory innervation of reproductive organs. *Prog Brain Res* 1986;67:115-131.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3823468&dopt=Abstract&itool=iconnoabstr
13. Meyer RA, Campbell JN, Raja SN. Peripheral neural mechanisms of nociception In: Wall PD, Melzack R, eds. *Textbook of Pain*. 3rd edn. Edinburgh: Churchill Livingstone, 1994.

1.2 Pain evaluation and measurement

1.2.1 Pain evaluation

The symptom of pain must be fully evaluated. As pain is subjective, the history provides the main evaluation. Examination and investigations provide further understanding of the pain syndrome and exclude other conditions. Pain rating(s) are essential in patient and treatment evaluation.

Pain evaluation includes:

- Baseline and ongoing regular evaluation of severity.
- An initial detailed history to include: chronology of onset and progression, character, site pain perceived and radiation, aggravating and relieving factors, associated symptoms.
- Questions about thoughts, emotions and behaviour associated with the pain.
- Detailed examination, not only of the painful area but of the whole patient, particularly the musculoskeletal and nervous systems.
- Investigations to identify well-defined/confusable/non-pain syndromes.
- Regular review of the condition as appropriate and its response to interventions.

1.2.2 Pain measurement

Pain can only be measured subjectively. The most reliable and well-understood method is a numerical rating scale, from 0 (no pain) to 10 (extreme pain), with half-points marked. This is superior to the widely used visual analogue scale (VAS), which is a 10-cm line with the same labels at the ends. Alternatively, a simple verbal rating scale can be used, e.g. 'none', 'mild', 'moderate', 'severe'. Both numerical and verbal scales can be used by patients without the need for paper and pen, unlike the visual analogue scale.

0	1	2	3	4	5	6	7	8	9	10
No pain										Extreme pain

Because pain is multidimensional, a single rating scale combines these dimensions in unknown quantities. Depending on the clinical question, treatment, patient and setting, it can be helpful to assess separately pain intensity, pain distress, and interference of pain with activities of daily life. It can also be helpful

to ask about average pain, worst pain (as even if this only occurs rarely, it can still reveal what patients should avoid) and pain on, for example, bladder voiding. Pain reduction or relief is measured directly using a percentage, from 0% = no relief up to 100% = total relief.

See www.britishpainsociety.org/members_pain_scales.htm for pain scales in English and other languages. The Brief Pain Inventory (1) consists of four 0 to 10 numerical scales for pain (current, average, worst, and least) and seven scales for interference with aspects of daily life: general activity, mood, walking ability, normal work, relationships with other people, sleep and enjoyment of life. The EuroQoL is a quality-of-life scale (2) available in several European languages and free for non-commercial use. It asks about mobility, self-care, pain, usual activities, and psychological status (www.euroqol.org).

1.2.3 REFERENCES

1. Tan G, Jensen MP, Thornby JI, Shanti BF. Validation of the Brief Pain Inventory for chronic nonmalignant pain. *J Pain* 2004;5(2):133-137.
http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=15042521&ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
www.mdanderson.org/pdf/bpif.pdf
2. Rabin R, de Charro F. EQ-5D: a measure of health status from the EuroQol group. *Ann Med* 2001;33(5):337-343.
http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=11491192&ordinalpos=5&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

2. CHRONIC PELVIC PAIN

2.1 Background

2.1.1 Introduction to chronic urogenital pain syndromes

Pain perceived within the pelvis may arise from a range of different mechanisms, many of which remain poorly understood. Some conditions have become 'well-defined' over the years and it is very important that these are identified and treated by an evidence-based approach, e.g. pudendal neuralgia.

Basic investigations must therefore be undertaken to rule out 'well-defined' pathologies. If the results are negative, a 'well-defined' pathology is unlikely. Any further investigations should be done only for specific indications, e.g. for subdivision of a pain syndrome.

As well as 'well-defined' conditions, conditions associated with chronic urogenital pain are becoming better understood. In many cases, the mechanisms involved are the neural-axial central sensitization described above and that are so familiar in other fields of chronic pain. There is now no doubt that these central changes can produce states of visceral and/or muscle hypersensitivity with long-term pain, sensory dysaesthesia and functional abnormalities. These need to be addressed, as well as the cognitive, behavioural, emotional and sexual consequences of the underlying disease process and long-term pain. This is why the assessment and management of these areas are also expanded in these guidelines and for the emphasis on a multidisciplinary approach.

Earlier EAU guidelines on chronic pelvic pain (CPP) introduced a classification system aimed at replacing old-fashioned terminology based on spurious assumptions of cause. The main emphasis was to make clear that it should not be presumed that pathology would be found where the pain is perceived. The EAU guidelines moved away from using 'prostatitis' and 'interstitial cystitis' in the absence of proven inflammation or infection, while the suffixes 'algia' and 'dynia' have often been used to provide a tangible diagnosis, which in itself may have a therapeutic benefit. In this edition of the guidelines, however, we have decided to avoid such terms completely. Instead, our definitions are based on the recommendation for terminology laid down by the International Continence Society (ICS) (1) and follow the axial structure of the International Association for the Study of Pain (IASP) classification (Table 1) (2).

Pain syndrome terms were introduced to indicate the multiple mechanisms involved, both physical and psychological. This approach has been reviewed on many occasions over the past few years and has been found to be robust. The EAU guidelines expand this approach, so avoiding spurious diagnostic terms, which are associated with inappropriate investigations, inappropriate treatments, inappropriate patient expectations and ultimately a worse prognostic outlook.

Table 1: EAU classification of chronic urogenital pain syndromes. This classification represents the efforts of many groups, as indicated in the main text. The work is in progress and further changes in this classification system are likely.*

Axis I Region	Axis II System	Axis III End organ as identified from Hx, Ex and Ix		Axis IV Referral characteristics	Axis V Temporal characteristics	Axis VI Character	Axis VII Associated symptoms	Axis VIII Psychological symptoms
Chronic pelvic pain syndrome	Urological	Bladder pain syndrome	(See Table 3 on ESSIC classification)	Suprapubic Inguinal Urethral	ONSET Acute Chronic	Aching Burning Stabbing Electric	URINARY Frequency Nocturia Hesitance Poor flow Piss en deux Urge Urgency Incontinence Other	Cognitive Behavioural Emotional
		Prostate pain syndrome	Type A inflammatory Type B non-inflammatory	Penile/clitoral Perineal Rectal Back Buttocks	ONGOING Sporadic Cyclical Continuous	Other		
		Scrotal pain syndrome	Testicular pain syndrome		TIME Filling Emptying Immediate post Late post		GYNAECOLOGICAL e.g Menstrual	
			Epididymal pain syndrome				SEXUAL e.g. Female dyspareunia impotence	
		Penile pain syndrome	Post-vasectomy pain syndrome		PROVOKED		GI	
Non pelvic pain syndrome	Gynaecological	Endometriosis associated pain syndrome						MUSCULAR Hyperalgesia CUTANEOUS Allodynia
		Vaginal pain syndrome						
		Vulvar pain syndrome	Generalised vulvar pain syndrome					
			Localised vulvar pain syndrome					
			Vestibular pain syndrome					
	Anorectal Neurological Muscular		Clitoral pain syndrome					
		e.g. Pudendal pain syndrome						
		e.g. Pudendal neuralgia						

Hx = History; Ex = Examination; Ix = Investigation.

*The table presented is not comprehensive; for the purpose of this document the main emphasis has been on the urological pain syndromes.

2.2 Definitions of chronic pelvic pain terminology

Although this latest EAU CPP guideline retains the basic terminology used in previous EAU CPP guidelines, older terminology has been removed (Table 2).

Table 2: Definitions of chronic pelvic pain terminology

Terminology	Description
<i>Chronic pelvic pain</i>	Non-malignant pain perceived in structures related to the pelvis of either men or women. In the case of documented nociceptive pain that becomes chronic, pain must have been continuous or recurrent for at least 6 months. If non-acute and central sensitization pain mechanisms are well documented, then the pain may be regarded as chronic, irrespective of the time period. In all cases, there often are associated negative cognitive, behavioural, sexual and emotional consequences (5,6)
<i>Pelvic pain syndrome</i>	Persistent or recurrent episodic pelvic pain associated with symptoms suggesting lower urinary tract, sexual, bowel or gynaecological dysfunction. No proven infection or other obvious pathology (adopted from ICS 2002 report) (1)
<i>Bladder pain syndrome</i>	Suprapubic pain is related to bladder filling, accompanied by other symptoms such as increased daytime and night-time frequency. There is an absence of proven urinary infection or other obvious pathology. This term has been adopted from the ICS 2002 report (1), where the term painful bladder syndrome was used; the name has been changed to bladder pain syndrome to be consistent with other pain syndrome terminology (5,6). The European Society for the Study of IC/PBS (ESSIC) publication places greater emphasis on the pain being perceived in the bladder (4)
<i>Urethral pain syndrome</i>	Recurrent episodic urethral pain, usually on voiding, with daytime frequency and nocturia. Absence of proven infection or other obvious pathology (1)
<i>Penile pain syndrome</i>	Pain within the penis that is not primarily in the urethra. Absence of proven infection or other obvious pathology (5,6)
<i>Prostate pain syndrome</i>	Persistent or recurrent episodic prostate pain, associated with symptoms suggestive of urinary tract and/or sexual dysfunction. No proven infection or other obvious pathology (5,6) Definition adapted from the National Institutes of Health (NIH) consensus definition and classification of prostatitis (7) and includes conditions described as 'chronic pelvic pain syndrome'. Using the NIH classification system, prostate pain syndrome may be subdivided into type A (inflammatory) and type B (non-inflammatory)
<i>Scrotal pain syndrome</i>	Persistent or recurrent episodic scrotal pain associated with symptoms suggestive of urinary tract or sexual dysfunction. No proven epididymo-orchitis or other obvious pathology (1)
<i>Testicular pain syndrome</i>	Persistent or recurrent episodic pain localized to the testis on examination, which is associated with symptoms suggestive of urinary tract or sexual dysfunction. No proven epididymo-orchitis or other obvious pathology. This is a more specific definition than scrotal pain syndrome (1)
<i>Post-vasectomy pain syndrome</i>	Scrotal pain syndrome that follows vasectomy (1)
<i>Epididymal pain syndrome</i>	Persistent or recurrent episodic pain localized to the epididymis on examination. Associated with symptoms suggestive of urinary tract or sexual dysfunction. No proven epididymo-orchitis or other obvious pathology (a more specific definition than scrotal pain syndrome (5,6)
<i>Endometriosis-associated pain syndrome</i>	Chronic or recurrent pelvic pain where endometriosis is present but does not fully explain all the symptoms (5,6)
<i>Vaginal pain syndrome</i>	Persistent or recurrent episodic vaginal pain associated with symptoms suggestive of urinary tract or sexual dysfunction. No proven vaginal infection or other obvious pathology (1)
<i>Vulvar pain syndrome</i>	Persistent or recurrent episodic vulvar pain either related to the micturition cycle or associated with symptoms suggestive of urinary tract or sexual dysfunction. There is no proven infection or other obvious pathology (1)

<i>Generalized vulvar pain syndrome</i> (formally dysaesthetic vulvodynia)	Vulval burning or pain that cannot be consistently and tightly localized by point-pressure 'mapping' by probing with a cotton-tipped applicator or similar instrument. The vulvar vestibule may be involved but the discomfort is not limited to the vestibule. Clinically, the pain may occur with or without provocation (touch, pressure or friction) (8)
<i>Localized vulvar pain syndrome</i>	Pain consistently and tightly localized by point-pressure mapping to one or more portions of the vulva. Clinically, pain usually occurs as a result of provocation (touch, pressure or friction) (8)
<i>Vestibular pain syndrome</i> (formerly vulval vestibulitis)	Pain localized by point-pressure mapping to one or more portions of the vulval vestibule (8)
<i>Clitoral pain syndrome</i>	Pain localized by point-pressure mapping to the clitoris (8)
<i>Anorectal pain syndrome</i>	Persistent or recurrent, episodic rectal pain with associated rectal trigger points/tenderness related to symptoms of bowel dysfunction. No proven infection or other obvious pathology (5,6)
<i>Pudendal pain syndrome</i>	Neuropathic-type pain arising in the distribution of the pudendal nerve with symptoms and signs of rectal, urinary tract or sexual dysfunction. No proven obvious pathology (5,6). (This is not the same as the well-defined pudendal neuralgia)
<i>Perineal pain syndrome</i>	Persistent or recurrent, episodic, perineal pain either related to the micturition cycle or associated with symptoms suggestive of urinary tract or sexual dysfunction. No proven infection or other obvious pathology (1)
<i>Pelvic floor muscle pain syndrome</i>	Persistent or recurrent, episodic, pelvic floor pain with associated trigger points, which is either related to the micturition cycle or associated with symptoms suggestive of urinary tract, bowel or sexual dysfunction. No proven infection or other obvious pathology (5,6)

2.3 Classification of chronic pelvic pain syndromes

The EAU classification of 2004 has been updated to provide a classification related to investigation and further management of the pain syndromes. This allows for a possible overlap of mechanisms between different conditions. It also encourages recognition of overlapping symptoms and their treatment by a multidisciplinary approach (Table 1) (3). Axes VII and VIII are very important, because they emphasize the importance of interdisciplinary, multidisciplinary assessment and management. This includes assessment and management of psychological symptoms and early involvement of the pain management centre.

Currently, there is no ideal classification for conditions considered to be chronic pain syndromes. The axes used in Table 1 are based on the IASP classification (2). Much of the terminology comes from the ICS classification of chronic pain (1) with input from the International Society for the Study of Vulvovaginal Disease (ISSVD), the IASP special interest group, Pain of Urogenital Origin (PUGO) group and Specialists in Pain International Network (SPIN). The major controversy in classifying chronic pain is that a pain may involve multiple sites, aetiologies and mechanisms. At a consensus meeting led by A.P. Baranowski and a PUGO working group in November 2006, it was suggested that a patient should be described as having one or two pain syndromes in the case of pain perceived at one or two sites, respectively. If the patient's pain was poorly localized or perceived in three or more sites, the patient would be diagnosed with chronic pelvic pain syndrome (CPPS), with no need for further system or end-organ subdivision. This decision recognized that poor localization in pain suggests overlapping mechanisms. This approach continues to be discussed and publications from a further consensus meeting are planned.

A physician using the classification in Table 1 should start on the left of Table 1 and proceed to the right only if they can truly and confidently confirm the pain to be perceived in the appropriate system and organ. In many cases, it may not be possible to go further than labelling a condition as a pelvic pain syndrome. For example, in many cases previously described as 'prostadynia', it may not be possible to state categorically that the pain stems from the prostate and not other sites, e.g. pelvic floor muscles. Such cases are therefore labelled pelvic pain syndrome.

Although bladder pain syndrome (BPS)/interstitial cystitis (IC) is well defined (see ESSIC reference [4]), many patients, who have been previously labelled as suffering from IC, would not meet the research criteria. These patients would therefore be labelled using Table 1 at some point to the left of IC, possibly as bladder pain syndrome (previously known as the painful bladder syndrome [1]). The term 'pain syndrome' is used when the primary pathology, which may be well defined and at one site to start with, progresses to produce a more complicated picture of pain, involving multiple sites and mechanisms. The condition has therefore become a complex of symptoms and signs, i.e. a syndrome.

The axial classification includes referral, temporal and character axes. These descriptors should also be collected for audit and research purposes. It should also be noted whether or not the pain is provoked, e.g. by pressure.

This EAU classification system aims to draw together the expertise of specialist groups and continues to undergo revision. The European Society for the Study of IC/PBS (ESSIC) have recently shown a way forward by defining the bladder pain syndrome/IC syndrome, supported by an international consensus editorial (3,4). As with the EAU system, ESSIC excluded well-defined (confusable) conditions. ESSIC has further divided the bladder pain syndrome/IC syndrome according to the results of cystoscopy and biopsy (Table 3) (4).

Table 3: ESSIC classification of types of bladder pain syndrome according to the results of cystoscopy with hydrodistension and of biopsies (4).

	Cystoscopy with hydrodistension			
	Not done	Normal	Glomerulations ^a	Hunner's lesion ^b
Biopsy				
Not done	XX	1X	2X	3X
Normal	XA	1A	2A	3A
Inconclusive	XB	1B	2B	3B
Positive ^c	XC	1C	2C	3C

^a Cystoscopy: glomerulations grade 2-3.

^b With or without glomerulations.

^c Histology showing inflammatory infiltrates and/or detrusor mastocytosis and/or granulation tissue and/or intrafascicular fibrosis.

2.4 REFERENCES

- Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, van Kerrebroeck P, Victor A, Wein A. The standardisation of terminology of lower urinary tract function: report from the Standardisation Subcommittee of the International Continence Society. *Am J Obstet Gynecol* 2002;187(1):116-126. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=12114899&dopt=Abstract
- Merskey H, Bogduk N. Classification of Chronic Pain. Descriptions of Chronic Pain Syndromes and Definitions of Pain Terms. IASP Press, 2002.
- Baranowski AP, Abrams P, Berger RE, Buffington CA, de C Williams AC, Hanno P, Loeser JD, Nickel JC, Wessellmann U. Urogenital pain—time to accept a new approach to phenotyping and, as a consequence, management. *Eur Urol* 2008;53(1):33-36. http://www.ncbi.nlm.nih.gov/pubmed/17961909?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- van de Merwe JP, Nordling J, Bouchelouche P, Bouchelouche K, Cervigni M, Dahi LK, Elneil S, Fall M, Hohlbrugger G, Irwin P, Mortensen S, van Ophoven A, Osborne JL, Pecker R, Richter B, Riedl C, Sairanen J, Tinzl M, Wyndaele JJ. Diagnostic criteria, classification, and nomenclature for painful bladder syndrome/interstitial cystitis: an ESSIC proposal. *Eur Urol* 2008;53(1):60-67. http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=17900797&ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- Fall M, Baranowski AP, Fowler CJ, Lepinard V, Malone-Lee JG, Messelink EJ, Oberpenning F, Osborne JL, Schumacher S. EAU Guidelines on Chronic Pelvic Pain. In: *EAU Guidelines*, edition published at the 18th Annual EAU Congress, Madrid, 2003, ISBN 90-70244-06-3.
- Fall M, Baranowski AP, Elneil S, Engeler D, Hughes J, Messelink EJ, Oberpenning F, Williams A C de C. EAU Guidelines on Chronic Pelvic Pain. In: *EAU Guidelines*. 23rd Annual EAU Congress, Milan, ISBN 978-90-70244-91-0. <http://www.uroweb.org/nc/professional-resources/guidelines/online/>
- Krieger JN, Nyberg L Jr, Nickel JC. NIH consensus definition and classification of prostatitis. *JAMA* 1999;282(3):236-237. http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10422990&dopt=Abstract
- Proceedings of the XVth World Congress. International Society for the Study of Vulvovaginal Disease, Santa Fe, NM, September 26-30, 1999. International Society for the Study of Vulvovaginal Disease Newsletter, Summer 2000.


2.5 An algorithm for chronic pelvic pain diagnosis and treatment

2.5.1 How to use the algorithm

The algorithm for diagnosing and treating chronic pelvic pain (CPP) (Figure 1) has been written to guide a physician through the process from diagnosis to management. A physician should follow steps 1 to 6 (Table 4), while referring to the correct column in the algorithm. Further guidance on which diagnostic tools should be

used in specific pain locations is provided in different chapters of this guideline.

Figure 1: An algorithm for diagnosing and managing chronic pelvic pain (CPP)



DRE = digital rectal examination; US = ultrasound; PFM = pelvic floor muscles.

Table 4: Step-by-step guidance on using the algorithm for diagnosis and treatment of CPP

Step	Action	Algorithm
1	Start by considering the organ system where the symptoms appear to be primarily perceived	First column
2	'Well-defined' conditions, such as cystitis, should be diagnosed and treated according to national or international guidelines	Second column and upper part third column
3	When treatment has no effect on the pain, additional tests (e.g. cystoscopy or ultrasound) should be performed	Lower part third column
4	When these tests reveal any pathology, this should be treated appropriately	Fourth column
5	If treatment has no effect, the patient should be referred to a pain team	Fifth column
6	If no well-defined condition is present or when no pathology is found by additional tests, the patient should also be referred to a pain team	Fifth column


The only aspect of diagnosis that is specific for CPP is where the pain is localized. However, because pain is perceived in structures related to the pelvis, this has led to many organ-specific, but often not well-defined, local disease syndromes.

Because CPP is pain perceived in structures related to the pelvis, it is necessary to approach diagnosis of a patient with CPP as a chronic pain patient. Confining the diagnosis to a specific organ may overlook multisystem functional abnormalities requiring individual treatment and general aspects of pain in planning investigation and treatment.

For the above reasons, we advocate early involvement of a multidisciplinary pain team. In practice, this should mean that well-known diseases, e.g. 'true' cystitis and endometriosis, will be diagnosed and treated early. If treating such conditions does not reduce symptoms, or such well-defined conditions are not found, then further investigation may be necessary, depending on where the pain is localized.

It should be noted, however, that over-investigation may be as harmful as not performing appropriate investigations. The EAU algorithms introduce the concept of the 'minimum investigations' required to exclude a well-defined condition.

2.6 Chronic prostate pain/chronic prostatitis associated with chronic pelvic pain syndrome (CP/CPPS)

Figure 2: General diagnostic and treatment algorithm for chronic prostate pain

NIH-CPSI = National Institute of Health chronic prostatitis symptom index; I-PSS = international prostate symptom score; DRE = digital rectal examination; PSA = prostate-specific antigen; PVR = post-void residual urine; PPMT = pre-post-massage test; CP/CPPS = chronic prostatitis/chronic pelvic pain syndrome; PCA = prostate cancer; NSAID = non-steroidal anti-inflammatory drug.

2.6.1 Introduction

Chronic prostatitis is an obscure and poorly understood disease. Restricted physical access has made it difficult to study the prostate gland, resulting in a lack of certainty about the aetiology, a lack of distinguishing clinical features, non-uniform diagnostic criteria and a protracted treatment course.

In about 5-10% of cases, clinical prostatitis has a proven bacterial aetiology. The remaining 90% of cases, in which laboratory methods have not found a bacterial cause, are classified as 'chronic non-bacterial prostatitis' or 'prostatodynia' (1-3). An appreciation of the fact that symptoms do not necessarily indicate isolated prostatic disease has led to a renaming of the condition: 'Chronic prostatitis associated with chronic pelvic pain syndrome' (CP/CPPS). This is now the term used by the NIH for patients with symptomatic prostatitis of non-bacterial origin (4).

2.6.2 Definition

Chronic prostatitis associated with chronic pelvic pain syndrome (CP/CPPS) is discomfort or pain in the pelvic region over a minimum of 3 months, with sterile specimen cultures and either significant, or insignificant, white blood cell counts in prostate-specific specimens (i.e. semen, expressed prostatic secretions and urine collected after prostate massage) (4). According to the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) classification, CP/CPPS is prostatitis category III (5) (Table 5). At present, there are no clinically relevant diagnostic or therapeutic results arising from differentiating inflammatory (NIH Cat. IIIA) from non-inflammatory (NIH Cat. IIIB) CP/CPPS. CP/CPPS Cat. III is therefore considered as one entity. According to the more general definition described in Section 2.2 (see Table 2), the disease is referred to as 'prostate pain syndrome (CP/CPPS)' throughout the rest of this chapter.

Table 5: Classification of prostatitis according to NIDDK/NIH

I.	Acute bacterial prostatitis (ABP)
II.	Chronic bacterial prostatitis (CBP)
III.	Chronic pelvic pain syndrome (CPPS)
	A. Inflammatory CPPS: WBC in semen/EPS/VB3
	B. Non-inflammatory CPPS: no WBC semen/EPS/VB3
IV.	Asymptomatic inflammatory prostatitis (histological prostatitis)

WBC = white blood cells; EPS = expressed prostatic secretions; VB3 = voided bladder urine-3.

2.6.3 Pathogenesis

The aetiology and pathophysiology of prostate pain syndrome (CP/CPPS) remains a mystery. Acute bacterial prostatitis is a different disease to chronic prostatitis syndromes. Patients with CPPS show no evidence of inflammation; they do not have urethritis, urogenital cancer, urethral stricture, or neurological disease involving the bladder nor exhibit any overt renal tract disease (4).

As often occurs with pelvic pain syndromes, there are several, poorly evidenced, hypotheses to explain the aetiology of prostate pain syndrome (CP/CPPS):

- Pain and subsequent irritative and obstructive voiding symptoms may be caused by lower urinary tract obstruction (LUTS), due to bladder neck problems, detrusor sphincter dysfunction, urethral stricture or dysfunctional voiding, resulting in high-pressure voiding (6-11).
- Intraprostatic ductal reflux caused by high-pressure turbulent voiding due to an anatomical abnormality (12-15).
- Microbiological cause, due to apparently harmless lower urinary tract commensals which require more sensitive isolation methods to be identified (4).
- Immunological processes precipitated by an unrecognized antigen or an autoimmune process (16-18). Urinary reflux into the prostatic ducts and acini might stimulate a sterile inflammatory response (13).
- A neuromuscular aetiology (19-21), in which symptoms represent a type of reflex sympathetic dystrophy of the perineum and pelvic floor.
- An interstitial cystitis-like pathogenic mechanism based on a significant overlap of symptomatology (pain, voiding symptoms) and cystoscopic or urodynamic findings. In patients diagnosed with prostate pain syndrome (CP/CPPS), a bladder-oriented interstitial cystitis mechanism accounts for the symptoms and the prostate is involved only indirectly (22).

2.6.4 Diagnosis

Despite its name, prostate pain syndrome (CP/CPPS) is a symptomatic diagnosis, which is diagnosed from a 3-month history of genitourinary pain and an absence of other lower urinary tract pathologies (see above). Determination of the severity of disease, its progression and treatment response can be assessed only by means of a validated symptom-scoring instrument (23,24). Quality of life should also be measured because it can be as poor as in acute myocardial infarction, unstable angina pectoris or Crohn's disease (25,26). Reliable, valid indexes of symptoms and quality of life are the NIH Prostatitis Symptom Index (NIH-CPSI) (27) and the International Prostate Symptom Score (I-PSS) (28). These subjective outcome measures are recommended for the basic evaluation and therapeutic monitoring of patients in urological practice and have been translated and validated for many European languages.

In prostate pain syndrome (CP/CPPS), urodynamic studies demonstrate decreased urinary flow rates, incomplete relaxation of the bladder neck and prostatic urethra, as well as abnormally high urethral closure pressure at rest. The external urethral sphincter is normal during urination (6,29).

Laboratory diagnosis has been classically based on the four-glass test for bacterial localization ('gold standard') (30). Besides a sterile pre-massage urine (voided bladder urine-2 [VB2]), CP/CPPS shows less than 10,000 colony-forming units of uropathogenic bacteria in expressed prostatic secretions (EPS) and insignificant numbers of leucocytes or bacterial growth in ejaculate. However, this test is too complex for use by practising urologists (4). Diagnostic efficiency may be enhanced cost-effectively by a simple screening procedure, i.e. the two-glass test or pre-post-massage test (PPMT) (31). In an extensive analysis of both tests, PPMT was able to indicate the correct diagnosis in more than 96% of patients (32).

A general algorithm for diagnosis and treatment of chronic prostatic pain is shown in Figure 2.

2.6.5 Treatment

Because of the unknown cause of prostate pain syndrome (CP/CPPS), many therapies used are based on anecdote. Most patients require multimodal treatment aimed at the main symptoms and taking comorbidities into account. In the past few years, results from randomized controlled trials (RCTs) have led to advances in standard and novel treatment options. Graded recommendations are given in Table 6.

2.6.5.1 Alpha-blockers. Increasing evidence from recent RCTs has shown that alpha-blockers, e.g. terazosin (33), alfuzosin (34), doxazosin (35), tamsulosin, reduce urinary symptoms and pain in prostate pain syndrome (CP/CPPS). The effects of alpha-antagonists may include improved outflow performance by blocking the alpha-receptors of the bladder neck and prostate and by direct action on alpha1A/1D-receptors in the CNS (36). Some symptoms may be relieved in roughly 50-60% of patients. Meta-analysis of nine trials (n = 734) showed that NIH-CPSI or I-PSS was reduced significantly with treatment duration of at least 3 months. However, in contrast to some individual studies (33-36), the meta-analysis showed no beneficial effect on pain (37). In addition, treatment-naïve and/or newly diagnosed patients appeared more likely to respond to alpha-blockers (38), although there is not sufficient evidence to conclude that they are effective (39). Because patients with heavily re-treated disease may not significantly improve in the short term compared to placebo (40), an alpha-blocker should be given for at least 3-6 months before assessing treatment (36).

2.6.5.2 Antibiotic therapy. Empirical antibiotic therapy is widely used because some patients have improved with antimicrobial therapy. Patients responding to antibiotics should be maintained on medication for 4-6 weeks or even longer. Unfortunately, culture, leucocyte and antibody status of prostate-specific specimens do not predict antibiotic response in patients with prostate pain syndrome (CP/CPPS) (41), and prostate biopsy culture findings do not differ from those of healthy controls (42). Long-term results with trimethoprim-sulphamethoxazole have remained poor (43-45). More encouraging results have been obtained with quinolones, including ciprofloxacin (46) and ofloxacin (41,47), but overall, antibiotic treatment of the prostate pain syndrome (CP/CPPS) is based only on weak evidence. After one unsuccessful course of a quinolone antibiotic over 4-6 weeks, other therapeutic options should be offered.

2.6.5.3 Non-steroidal anti-inflammatory drugs. Non-steroidal anti-inflammatory drugs may have favourable results in some patients. Immunomodulation using cytokine inhibitors or other approaches may be helpful, but proper trials are needed before this type of therapy can be recommended (48,49). Only one RCT has been published. This was for rofecoxib, which is no longer on the market; statistical significance was achieved in some of the outcome measures (50).

2.6.5.4 Corticosteroids are not recommended. A few anecdotal case reports have shown some improvement. However, no significant benefits were shown in a low-power, placebo-controlled, randomized pilot study of a short course of oral prednisolone (51).

2.6.5.5 Opioids produce modest pain relief in some patients with refractory prostate pain syndrome (CP/CPPS), though there is limited data on the long-term efficacy of opioids in non-cancer pain. Opioid treatment carries the risks of side effects, reduced quality of life, addiction, opioid tolerance and opioid-induced hyperalgesia (52). Urologists should use opioids for prostate pain syndrome (CP/CPPS) in collaboration with pain clinics and with other treatments.

2.6.5.6 5-alpha-reductase inhibitors. A few small pilot studies with 5-alpha-reductase inhibitors supported the view that finasteride may improve voiding and pain (53-56). In a randomized trial, finasteride provided better amelioration of symptoms compared to saw palmetto over a 1-year period, but lacked a placebo-control arm (57). A 6-month placebo-controlled study showed a tendency towards better outcome in favour of finasteride without statistical significance, possibly because of a lack of power (58).

2.6.5.7 Allopurinol. An RCT of allopurinol was conducted based on the hypothesis that urine reflux into prostatic ducts causes prostatic inflammation via high concentrations of purine and pyrimidine base-containing metabolites in prostatic secretions (59). However, positive results were not considered to be sufficient for recommendation by reviewers of the Cochrane Database (60). In addition, a recent randomized placebo-controlled trial of allopurinol as an adjunct to ofloxacin has not shown any benefit (61).

2.6.5.8 Phytotherapy. Positive effects of phytotherapy have been documented. Although a validated symptom score was not used, an RCT of a pollen extract (Prostat/Poltit) showed significant symptom improvement in the pollen-treated group (62). Another pollen extract, Cernilton N, provided only weak improvement. For 'uncomplicated' cases, a 36% cure rate could be shown over a 6-month period in a prospective study (63). Quercetin, a polyphenolic bioflavonoid with documented antioxidant and anti-inflammatory properties, improved NIH-CPSI scores significantly in a small RCT (64). In addition, high-dose oral PPS, as for interstitial cystitis, is able to ameliorate symptoms and improve quality of life significantly in men with prostate pain syndrome (CP/CPPS), suggesting a possible common aetiology (65). In contrast, treatment with saw palmetto, most commonly used for benign prostatic hyperplasia, did not improve symptoms over a 1-year period (57).

2.6.5.9 Muscle relaxants (diazepam, baclofen) are claimed to be helpful in sphincter dysfunction or pelvic floor/perineal muscle spasm, but there have been only few prospective clinical trials to support these claims (21). In a recent RCT, a triple combination of a muscle relaxant (tiocolchicoside), an anti-inflammatory drug (ibuprofen) and an alpha-blocker (doxazosin) was effective in treatment-naïve patients, but not superior to an alpha-blocker alone (66).

2.6.5.10 Supportive therapies, such as biofeedback, relaxation exercises, lifestyle changes (i.e. diet, discontinuing bike riding), acupuncture, massage therapy, chiropractic therapy or meditation, have all been claimed to improve symptoms (4,67). In a small, sham-controlled, double-blind study, 4-week electromagnetic therapy showed a significant, sustained effect over a 1-year period (68). Some patients have reported favourable effects from heat therapy, e.g. transrectal hyperthermia (69-72) and transurethral thermotherapy (73-77).

2.6.5.11 Surgical management, including transurethral incision of the bladder neck (9), radical transurethral resection of the prostate (78,79) or in particular radical prostatectomy, has a very limited role and requires an additional, specific indication (67). In addition, the treatment effect of transurethral needle ablation of the prostate (TUNA) was only comparable to sham treatment (80).

Table 6: Treatment of prostate pain syndrome (CP/CPPS)

	Level of evidence	Grade of recommendation	Comment
• Alpha-blockers	1a	A	Effect on total NIH-CPSI
• Muscle relaxants	3	C	Only very limited data
• Antimicrobial therapy	3	B	Quinolones If previously untreated (naïve) only, reassess after 2-3 weeks. Duration 4-6 weeks

• Opioids	3	C	As part of multimodal therapy for treatment-refractory pain in collaboration with pain clinics
• Non-steroidal anti-inflammatory drugs	1b	B	Long-term side effects have to be considered
• Steroids	3	Not recommended	Not outside clinical trials
• Immunosuppressive agents			
• 5-alpha-reductase inhibitors	1b	B	If benign prostatic hyperplasia is present
• Phytotherapy	1b-3	B	
• Biofeedback, relaxation exercise	2a-3	B	As supportive, second-line therapies
• Lifestyle changes			
• Massage therapy			
• Chiropractor therapy			
• Acupuncture			
• Meditation			
• Electromagnetic therapy	1b	C	Not outside clinical trials
• Transrectal hyperthermia	3	C	
• Transurethral thermotherapy			
• Transurethral incision of the bladder neck	3	Not recommended general	Specific additional indication required
• Transurethral resection of the prostate			
• Radical prostatectomy			

2.6.6 References

- de la Rosette JJ, Hubregtse MR, Meuleman EJ, Stolk-Engelaar MV, Debruyne FM. Diagnosis and treatment of 409 patients with prostatitis syndromes. *Urology* 1993;41(4):301-307.
http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=8470312&ordinalpos=47&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- Meares EM Jr. Prostatitis. *Med Clin North Am* 1991;75(2):405-424.
http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=1996042&ordinalpos=86&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- Brunner H, Weidner W, Schiefer HG. Studies on the role of *Ureaplasma urealyticum* and *Mycoplasma hominis* in prostatitis. *J Infect Dis* 1983;147(5):807-813.
http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=6842018&ordinalpos=32&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- Nickel JC, Weidner W. Chronic prostatitis: current concepts and antimicrobial therapy. *Infect Urol* 2000;13:S22-S28.
- Nickel JC. Prostatitis: myths and realities. *Urology* 1998;51(3):362-326.
http://www.ncbi.nlm.nih.gov/pubmed/9510337?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- Barbalias GA, Meares EM Jr, Sant GR. Prostatodynia: clinical and urodynamic characteristics. *J Urol* 1983;130:514-517.
http://www.ncbi.nlm.nih.gov/pubmed/6887365?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- Blacklock NJ. Urodynamic and psychometric observations and their implication in the management of prostatodynia. In: Weidner W, Brunner H, Krause W et al. (eds.) *Therapy of Prostatitis*. Munich: Zuckschwerdt Verlag, 1986:201.
- Hellstrom WJ, Schmidt RA, Lue TF, Tanagho EA. Neuromuscular dysfunction in nonbacterial prostatitis. *Urology* 1987;30(2):183-188.
http://www.ncbi.nlm.nih.gov/pubmed/3497475?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

9. Kaplan SA, Te AE, Jacobs BZ. Urodynamic evidence of vesical neck obstruction in men with misdiagnosed chronic nonbacterial prostatitis and the therapeutic role of endoscopic incision of the bladder neck. *J Urol* 1994;152(6 Pt 1):2063-2065.
http://www.ncbi.nlm.nih.gov/pubmed/7966675?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
10. Kaplan SA, Santarosa RP, D'Alisera PM, Fay BJ, Ikeguchi EF, Hendricks J, Klein L, Te AE. Pseudodyssynergia (contraction of the external sphincter during voiding) misdiagnosed as chronic nonbacterial prostatitis and the role of biofeedback as a therapeutic option. *J Urol* 1997;157(6):2234-2237.
http://www.ncbi.nlm.nih.gov/pubmed/9146624?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
11. Murnaghan GF, Millard RJ. Urodynamic evaluation of bladder neck obstruction in chronic prostatitis. *Br J Urol* 1984;56(6):713-716.
http://www.ncbi.nlm.nih.gov/pubmed/6534495?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
12. Blacklock NJ. The anatomy of the prostate: relationship with prostatic infection. *Infection* 1991;19(Suppl 3):S111-S114.
http://www.ncbi.nlm.nih.gov/pubmed/2055644?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
13. Persson BE, Ronquist G. Evidence for a mechanistic association between nonbacterial prostatitis and levels of urate and creatinine in expressed prostatic secretion. *J Urol* 1996;155(3):958-960.
http://www.ncbi.nlm.nih.gov/pubmed/8583617?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
14. Blacklock NJ. Anatomical factors in prostatitis. *Br J Urol* 1974;46(1):47-54.
http://www.ncbi.nlm.nih.gov/pubmed/4406038?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
15. Kirby RS, Lowe D, Bultitude MI, Shuttleworth KE. Intra-prostatic urinary reflux: an aetiological factor in abacterial prostatitis. *Br J Urol* 1982;54(6):729-731.
http://www.ncbi.nlm.nih.gov/pubmed/7150931?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
16. Doble A, Walker MM, Harris JR, Taylor-Robinson D, Witherow RO. Intraprostatic antibody deposition in chronic abacterial prostatitis. *Br J Urol* 1990;65(6):598-605.
http://www.ncbi.nlm.nih.gov/pubmed/2196972?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
17. Nickel JC, Olson ME, Barabas A, Benediktsson H, Dasgupta MK, Costerton JW. Pathogenesis of chronic bacterial prostatitis in an animal model. *Br J Urol* 1990;66(1):47-54.
http://www.ncbi.nlm.nih.gov/pubmed/2203502?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
18. Shortliffe LM, Wehner N. The characterization of bacterial and nonbacterial prostatitis by prostatic immunoglobulins. *Medicine (Baltimore)* 1986;65(6):399-414.
http://www.ncbi.nlm.nih.gov/pubmed/3537628?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
19. Andersen JT. Treatment of prostatodynia. In: Nickel JC (ed). *Textbook of Prostatitis*. London: ISIS, 1999.
20. Egan KJ, Krieger JL. Chronic abacterial prostatitis—a urological chronic pain syndrome? *Pain* 1997;69(3):213-218.
http://www.ncbi.nlm.nih.gov/pubmed/9085294?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
21. Osborn DE, George NJ, Rao PN, Barnard RJ, Reading C, Marklow C, Blacklock NJ. Prostatodynia—physiological characteristics and rational management with muscle relaxants. *Br J Urol* 1981;53(6):621-623.
http://www.ncbi.nlm.nih.gov/pubmed/7032641?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
22. Sant GR, Nickel JC. Interstitial cystitis in chronic prostatitis: The same syndrome? In: Nickle JC (ed). *Textbook of Prostatitis*. Oxford (UK): Medical Media, 1999: 169-176.
23. Barry MJ, Fowler FJ Jr, O'Leary MP, Bruskewitz RC, Holtgrewe HL, Mebust WK, Cockett AT. The American Urological Association symptom index for benign prostatic hyperplasia. The Measurement Committee of the American Urological Association. *J Urol* 1992;148(5):1549-1557; discussion 1564.
http://www.ncbi.nlm.nih.gov/pubmed/1279218?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

24. Nickel JC. Effective office management of chronic prostatitis. *Urol Clin North Am* 1998;25(4):677-684.
http://www.ncbi.nlm.nih.gov/pubmed/10026774?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
25. Wenninger K, Heiman JR, Rothman I, Berghuis JP, Berger RE. Sickness impact of chronic nonbacterial prostatitis and its correlates. *J Urol* 1996;155(3):965-968.
http://www.ncbi.nlm.nih.gov/pubmed/8583619?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
26. McNaughton Collins M, Pontari MA, O'Leary MP, Calhoun EA, Santanna J, Landis JR, Kusek JW, Litwin MS; Chronic Prostatitis Collaborative Research Network. Quality of life is impaired in men with chronic prostatitis: the Chronic Prostatitis Collaborative Research Network. *J Gen Intern Med* 2001;16(10):656-662.
http://www.ncbi.nlm.nih.gov/pubmed/11679032?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
27. Litwin MS, McNaughton-Collins M, Fowler FJ Jr, Nickel JC, Calhoun EA, Pontari MA, Alexander RB, Farrar JT, O'Leary MP. The National Institutes of Health chronic prostatitis symptom index: development and validation of a new outcome measure. Chronic Prostatitis Collaborative Research Network. *J Urol* 1999;162(2):369-375.
http://www.ncbi.nlm.nih.gov/pubmed/10411041?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
28. Mebust WK, Bosch R, Donovan J et al. Symptom evaluation, quality of life and sexuality. In: Cockett ATK, Khoury S, Aso Y et al. *Proceedings of the 2nd Consultation on Benign Prostatic Hyperplasia (BPH)*, Paris. Channel Islands: Scientific Communication International Ltd, 1993:129. www.congress-urology.org
29. Meares EMJ, Minich W. Prostatodynia: clinical findings and rationale for treatment. In: Weidner W, Brunner H, Krause W et al. *Therapy of Prostatitis*. Munich: Zuckschwerdt Verlag, 1986:207.
30. Meares EM, Stamey TA. Bacteriologic localization patterns in bacterial prostatitis and urethritis. *Invest Urol* 1968;5(5):492-518.
http://www.ncbi.nlm.nih.gov/pubmed/4870505?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
31. Nickel JC. The Pre and Post Massage Test (PPMT): a simple screen for prostatitis. *Tech Urol* 1997;3(1):38-43.
http://www.ncbi.nlm.nih.gov/pubmed/9170224?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
32. Nickel JC, Shoskes D, Wang Y, Alexander RB, Fowler JE Jr, Zeitlin S, O'Leary MP, Pontari MA, Schaeffer AJ, Landis JR, Nyberg L, Kusek JW, Probert KJ. How does the pre-massage and post-massage 2-glass test compare to the Meares-Stamey 4-glass test in men with chronic prostatitis/chronic pelvic pain syndrome? *J Urol* 2006;176(1):119-124.
http://www.ncbi.nlm.nih.gov/pubmed/16753385?ordinalpos=6&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
33. Cheah PY, Liong ML, Yuen KH, Teh CL, Khor T, Yang JR, Yap HW, Krieger JN. Terazosin therapy for chronic prostatitis/chronic pelvic pain syndrome: a randomized, placebo controlled trial. *J Urol* 2003;169(2):592-596.
http://www.ncbi.nlm.nih.gov/pubmed/12544314?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
34. Mehik A, Alas P, Nickel JC, Sarpola A, Helstrom PJ. Alfuzosin treatment for chronic prostatitis/chronic pelvic pain syndrome: a prospective, randomized, double-blind, placebo-controlled, pilot study. *Urology* 2003;62(3):425-429.
http://www.ncbi.nlm.nih.gov/pubmed/12946740?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
35. Evliyaoglu Y, Burgut R. Lower urinary tract symptoms, pain and quality of life assessment in chronic non-bacterial prostatitis patients treated with alpha-blocking agent doxazosin; versus placebo. *Int Urol Nephrol* 2002;34(3):351-356.
http://www.ncbi.nlm.nih.gov/pubmed/12899226?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
36. Nickel JC, Narayan P, McKay J, Doyle C. Treatment of chronic prostatitis/chronic pelvic pain syndrome with tamsulosin: a randomized double blind trial. *J Urol* 2004;171(4):1594-1597.
http://www.ncbi.nlm.nih.gov/pubmed/15017228?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

37. Yang G, Wei Q, Li H, Yang Y, Zhang S, Dong Q. The effect of alpha-adrenergic antagonists in chronic prostatitis/chronic pelvic pain syndrome: a meta-analysis of randomized controlled trials. *J Androl* 2006;27(6):847-852.
http://www.ncbi.nlm.nih.gov/pubmed/16870951?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
38. Lee SW, Liong ML, Yuen KH, Liong YV, Krieger JN. Chronic prostatitis/chronic pelvic pain syndrome: role of alpha blocker therapy. *Urol Int* 2007;78(2):97-105.
http://www.ncbi.nlm.nih.gov/pubmed/17293646?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
39. Mishra VC, Browne J, Emberton M. Role of alpha-blockers in type III prostatitis: a systematic review of the literature. *J Urol* 2007;177(1):25-30.
http://www.ncbi.nlm.nih.gov/pubmed/17161995?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
40. Alexander RB, Probert KJ, Schaeffer AJ, Landis JR, Nickel JC, O'Leary MP, Pontari MA, McNaughton-Collins M, Shoskes DA, Comiter CV, Datta NS, Flower JE Jr, Nadler RB, Zeitlin SI, Knauss JS, Wang Y, Kusek JW, Nyberg LM Jr, Litwin MS; Chronic Prostatitis Collaborative Research Network. Ciprofloxacin or tamsulosin in men with chronic prostatitis/chronic pelvic pain syndrome: a randomized, double-blind trial. *Ann Intern Med* 2004;141(8):581-589.
http://www.ncbi.nlm.nih.gov/pubmed/15492337?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
41. Nickel JC, Downey J, Johnston B, Clark J; Canadian Prostatitis Research Group. Predictors of patient response to antibiotic therapy for the chronic prostatitis/chronic pelvic pain syndrome: a prospective multicenter clinical trial. *J Urol* 2001;165(5):1539-1544.
http://www.ncbi.nlm.nih.gov/pubmed/11342913?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
42. Lee JC, Muller CH, Rothman I, Agnew KJ, Eschenbach D, Ciol MA, Turner JA, Berger RE. Prostate biopsy culture findings of men with chronic pelvic pain syndrome do not differ from those of healthy controls. *J Urol* 2003;169(2):587-588.
http://www.ncbi.nlm.nih.gov/pubmed/12544312?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
43. Drach GW. Trimethoprim sulfamethoxazole therapy of chronic bacterial prostatitis. *J Urol* 1974;111(5):637-639.
http://www.ncbi.nlm.nih.gov/pubmed/4274697?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
44. McGuire EJ, Lytton B. Bacterial prostatitis: treatment with trimethoprim-sulfamethoxazole. *Urology* 1976;7(5):499-500.
http://www.ncbi.nlm.nih.gov/pubmed/1274009?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
45. Meares EM. Long-term therapy of chronic bacterial prostatitis with trimethoprim-sulfamethoxazole. *Can Med Assoc J* 1975;112(13 Spec No):22-25.
http://www.ncbi.nlm.nih.gov/pubmed/236820?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
46. Weidner W, Schiefer HG, Brahler E. Refractory chronic bacterial prostatitis: a re-evaluation of ciprofloxacin treatment after a median followup of 30 months. *J Urol* 1991;146(2):350-352.
http://www.ncbi.nlm.nih.gov/pubmed/1856930?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
47. Cox CE. Ofloxacin in the management of complicated urinary tract infections, including prostatitis. *Am J Med* 1989;87(6C):61S-68S.
http://www.ncbi.nlm.nih.gov/pubmed/2690622?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
48. Canale D, Scaricabarozzi I, Giorgi P, Turchi P, Ducci M, Menchini-Fabris GF. Use of a novel non-steroidal anti-inflammatory drug, nimesulide, in the treatment of abacterial prostatovesiculitis. *Andrologia* 1993;25(3):163-166.
http://www.ncbi.nlm.nih.gov/pubmed/14550427?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
49. Canale D, Turchi P, Giorgi PM, Scaricabarozzi I, Menchini-Fabris GF. Treatment of abacterial prostatovesiculitis with nimesulide. *Drugs* 1993;46 (Suppl 1):147-150.
http://www.ncbi.nlm.nih.gov/pubmed/7506156?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

50. Nickel JC, Pontari M, Moon T, Gittelman M, Malek G, Farrington J, Pearson J, Krupa D, Bach M, Drisko J; Rofecoxib Prostatitis Investigator Team. A randomized, placebo controlled, multicenter study to evaluate the safety and efficacy of rofecoxib in the treatment of chronic nonbacterial prostatitis. *J Urol* 2003;169(4):1401-1405.
http://www.ncbi.nlm.nih.gov/pubmed/12629372?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
51. Bates SM, Hill VA, Anderson JB, Chapple CR, Spence R, Ryan C, Talbot MD. A prospective, randomized, double-blind trial to evaluate the role of a short reducing course of oral corticosteroid therapy in the treatment of chronic prostatitis/chronic pelvic pain syndrome. *BJU Int* 2007;99(2):355-359.
http://www.ncbi.nlm.nih.gov/pubmed/17313424?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
52. Nickel JC. Opioids for chronic prostatitis and interstitial cystitis: lessons learned from the 11th World Congress on Pain. *Urology* 2006;68(4):697-701.
http://www.ncbi.nlm.nih.gov/pubmed/17070334?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
53. Olavi L, Make L, Imo M. Effects of finasteride in patients with chronic idiopathic prostatitis: a doubleblind, placebo-controlled pilot study. *Eur Urol* 1998;33(Suppl. 1):33.
54. Golio G. The use of finasteride in the treatment to chronic nonbacterial prostatitis. In: *Abstracts of the 49th Annual Meeting of the Northeastern Section of the American Urological Association*. Phoenix, 1997: 128.
55. Holm M, Meyhoff HH. Chronic prostatic pain. A new treatment option with finasteride? *Scand J Urol Nephrol* 1997;31(2):213-215.
http://www.ncbi.nlm.nih.gov/pubmed/9165592?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
56. Leskinen M, Lukkarinen O, Marttila T. Effects of finasteride in patients with inflammatory chronic pelvic pain syndrome: a double-blind, placebo-controlled, pilot study. *Urology* 1999;53(3):502-505.
http://www.ncbi.nlm.nih.gov/pubmed/10096374?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
57. Kaplan SA, Volpe MA, Te AE. A prospective, 1-year trial using saw palmetto versus finasteride in the treatment of category III prostatitis/chronic pelvic pain syndrome. *J Urol* 2004;171:284-288.
http://www.ncbi.nlm.nih.gov/pubmed/14665895?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
58. Nickel JC, Downey J, Pontari MA, Shoskes DA, Zeitlin SI. A randomized placebo-controlled multicentre study to evaluate the safety and efficacy of finasteride for male chronic pelvic pain syndrome (category IIIA chronic nonbacterial prostatitis). *BJU Int* 2004;93(7):991-995.
http://www.ncbi.nlm.nih.gov/pubmed/15142149?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
59. Persson BE, Ronquist G, Ekblom M. Ameliorative effect of allopurinol on nonbacterial prostatitis: a parallel double-blind controlled study. *J Urol* 1996;155(3):961-964.
http://www.ncbi.nlm.nih.gov/pubmed/8583618?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
60. McNaughton CO, Wilt T. Allopurinol for chronic prostatitis. *Cochrane Database Syst Rev* 2002;(4):CD001041.
http://www.ncbi.nlm.nih.gov/pubmed/12519549?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
61. Ziaee AM, Akhavadegan H, Karbakhsh M. Effect of allopurinol in chronic nonbacterial prostatitis: a double blind randomized clinical trial. *Int Braz J Urol* 2006;32(2):181-186.
http://www.ncbi.nlm.nih.gov/pubmed/16650295?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
62. Elist J. Effects of pollen extract preparation Prostat/Poltit on lower urinary tract symptoms in patients with chronic nonbacterial prostatitis/chronic pelvic pain syndrome: a randomized, double-blind, placebo-controlled study. *Urology* 2006;67(1):60-63.
http://www.ncbi.nlm.nih.gov/pubmed/16413333?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
63. Rugendorff EW, Weidner W, Ebeling L, Buck AC. Results of treatment with pollen extract (Cernilton N) in chronic prostatitis and prostatodynia. *Br J Urol* 1993;71(4):433-448.
http://www.ncbi.nlm.nih.gov/pubmed/8499988?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

64. Shoskes DA, Zeitlin SI, Shahed A, Rajfer J. Quercetin in men with category III chronic prostatitis: a preliminary prospective, double-blind, placebo-controlled trial. *Urology* 1999;54(6):960-963.
http://www.ncbi.nlm.nih.gov/pubmed/10604689?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
65. Nickel JC, Forrest JB, Tomera K, Hernandez-Graulau J, Moon TD, Schaeffer AJ, Krieger JN, Zeitlin SI, Evans RJ, Lama DJ, Neal DE Jr, Sant GR. Pentosan polysulfate sodium therapy for men with chronic pelvic pain syndrome: a multicenter, randomized, placebo controlled study. *J Urol* 2005;173(4):1252-1255.
http://www.ncbi.nlm.nih.gov/pubmed/15758763?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
66. Tugcu V, Tasci AI, Fazlioglu, Gurbuz G, Ozbek E, Sahin S, Kurtulus F, Cek M. A placebo-controlled comparison of the efficiency of triple- and monotherapy in category III B chronic pelvic pain syndrome (CPPS). *Eur Urol* 2007;51(4):1113-1137; discussion 1118.
http://www.ncbi.nlm.nih.gov/pubmed/17084960?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
67. Nickel JC. Prostatitis: evolving management strategies. *Urol Clin North Am* 1999;26(4):737-751.
http://www.ncbi.nlm.nih.gov/pubmed/10584615?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
68. Rowe E, Smith C, Laverick L, Elkabir J, Witherow RO, Patel A. A prospective, randomized, placebo controlled, double-blind study of pelvic electromagnetic therapy for the treatment of chronic pelvic pain syndrome with 1 year of followup. *J Urol* 2005;173(6):2044-2047.
http://www.ncbi.nlm.nih.gov/pubmed/15879822?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
69. Kamihira O, Sahashi M, Yamada S et al. [Transrectal hyperthermia for chronic prostatitis.] *Nippon Hinyokika Gakkai Zasshi* 1993;84(6):1095-1098. [Japanese]
http://www.ncbi.nlm.nih.gov/pubmed/8345726?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
70. Kumon H, Ono N, Uno S et al. [Transrectal hyperthermia for the treatment of chronic prostatitis.] *Nippon Hinyokika Gakkai Zasshi* 1993;84(2):265-271. [Japanese]
http://www.ncbi.nlm.nih.gov/pubmed/8464182?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
71. Montorsi F, Guazzoni G, Bergamaschi F, Galli L, Consonni P, Matozzo V, Barbieri L, Rigatti P. Is there a role for transrectal microwave hyperthermia of the prostate in the treatment of abacterial prostatitis and prostatodynia? *Prostate* 1993;22(2):139-146.
http://www.ncbi.nlm.nih.gov/pubmed/8456052?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
72. Shaw TK, Watson GM, Barnes DG. Microwave hyperthermia in the treatment of chronic abacterial prostatitis and prostatodynia: results of a double-blind placebo controlled trial. *J Urol* 1993;149:405A.
73. Choi NG, Soh SH, Yoon TH, Song MH. Clinical experience with transurethral microwave thermotherapy for chronic nonbacterial prostatitis and prostatodynia. *J Endourol* 1994;8(1):61-64.
http://www.ncbi.nlm.nih.gov/pubmed/7514470?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
74. Michielsen D, Van Camp K, Wyndaele JJ, Verheyden B. Transurethral microwave thermotherapy in the treatment of chronic abacterial prostatitis: a 2 years follow-up. *Acta Urol Belg* 1995;63(4):1-4.
http://www.ncbi.nlm.nih.gov/pubmed/8644548?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
75. Nickel JC, Sorenson R. Transurethral microwave thermotherapy of nonbacterial prostatitis and prostatodynia: initial experience. *Urology* 1994;44(3):458-460.
http://www.ncbi.nlm.nih.gov/pubmed/8073567?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
76. Nickel JC, Sorensen R. Transurethral microwave thermotherapy for nonbacterial prostatitis: a randomized double-blind sham controlled study using new prostatitis specific assessment questionnaires. *J Urol* 1996;155(6):1950-1954; discussion 4-5.
http://www.ncbi.nlm.nih.gov/pubmed/8618295?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVAbstractPlusDrugs1
77. Kastner C, Hochreiter W, Huidobro C, Cabezas J, Miller P. Cooled transurethral microwave thermotherapy for intractable chronic prostatitis—results of a pilot study after 1 year. *Urology* 2004;64(6):1149-1154.
http://www.ncbi.nlm.nih.gov/pubmed/15596188?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

78. Barnes RW, Hadley HL, O'Donoghue EP. Transurethral resection of the prostate for chronic bacterial prostatitis. *Prostate* 1982;3(3):215-219.
http://www.ncbi.nlm.nih.gov/pubmed/7100001?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
79. Sant GR, Heaney JA, Meares EM. Radical transurethral prostatic resection in the management of chronic bacterial prostatitis. *J Urol* 1984;131:184A.
80. Leskinen MJ, Kilponen A, Lukkarinen O, Tammela TL. Transurethral needle ablation for the treatment of chronic pelvic pain syndrome (category III prostatitis): a randomized, sham-controlled study. *Urology* 2002;60(2):300-304.
http://www.ncbi.nlm.nih.gov/pubmed/12137830?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

2.7 Bladder pain syndrome/interstitial cystitis (BPS/IC)

2.7.1 Introduction

Interstitial cystitis describes a chronic, distressing bladder condition (1). The so-called 'ulcer', which is a typical cystoscopic finding in 10-50% of IC patients, was first described by Guy L Hunner at the beginning of the last century (2,3). Subsequent research (4-6) showed that IC was not a single entity, but had different endoscopic and histopathological presentations.

It is very important to realise that IC is a heterogeneous spectrum of disorders, which are still poorly defined, and that inflammation is an important feature in only a subset of patients. To embrace all patients suffering from bladder pain, the terms painful bladder syndrome (PBS) or bladder pain syndrome (BPS) have been suggested as more accurate terminology (7,8). This terminology assumes that IC represents a special type of chronic inflammation of the bladder, while PBS or BPS refers to pain in the bladder region. The term bladder pain syndrome or BPS will be used in these guidelines.

2.7.2 Definition

An extremely wide variety of diagnostic criteria have been used because of the difficulty in defining IC. In the late 1980s, NIDDK consensus criteria were established to ensure that scientific studies would be relatively comparable (Table 7) (9). The NIDDK criteria produce a diagnosis of IC by exclusion. Bladder pain, urgency and the finding of submucosal haemorrhages, called glomerulations, are the only positive elements. Identification of circumscribed Hunner-type lesions is an automatic inclusion criterion. Although generally accepted, the NIDDK criteria provide only a minimum framework to establish the diagnosis and some have felt them to be too restrictive for clinical use (10). Whatever the method used, heterogeneity seems currently unavoidable (6,11,12).

Table 7: NIDDK workshop research definition* of interstitial cystitis (IC) (9)**

Automatic inclusions
<ul style="list-style-type: none"> • Hunner's ulcer
Positive factors
<ul style="list-style-type: none"> • Pain on bladder filling relieved by emptying • Pain (suprapubic, pelvic, urethral, vaginal or perineal) • Glomerulations on endoscopy • Decreased compliance on cystometrogram
Automatic exclusions
<ul style="list-style-type: none"> • < 18 years old • Benign or malignant bladder tumours • Radiation cystitis • Tuberculous cystitis • Bacterial cystitis • Vaginitis • Cyclophosphamide cystitis • Symptomatic urethral diverticulum • Uterine, cervical, vaginal or urethral cancer • Active herpes • Bladder or lower ureteral calculi • Waking frequency < five times in 12 hours • Nocturia < two times

- Symptoms relieved by antibiotics, urinary antiseptics, urinary analgesics, e.g. phenazopyridine hydrochloride
- Duration < 12 months
- Involuntary bladder contractions (urodynamics)
- Capacity > 400 mL, absence of sensory urgency

** Bladder distension was defined arbitrarily as 80 cm water pressure for 1 minute (sic). Two positive factors were necessary for inclusion in the study population. Under anaesthesia, patients were sub-stratified at the end of the study into two groups according to bladder capacity < 350 mL and > 350 mL.

Recently, the European Society for the Study of IC/PBS (ESSIC) has suggested a standardized scheme of diagnostic criteria (13) to make it easier to compare different studies. In a consensus statement, the diagnosis of bladder pain syndrome (BPS) was preferred as the general term to match the current taxonomy of chronic pain syndromes.

Bladder pain syndrome should be diagnosed on the basis of symptoms of pain associated with the urinary bladder, accompanied by at least one other symptom, such as day-time and/or night-time urinary frequency, the exclusion of confusable diseases as the cause of symptoms, and if indicated, cystoscopy with hydrodistension and biopsy (Table 8) (8).

Table 8: ESSIC classification of types of bladder pain syndrome according to the results of cystoscopy with hydrodistension and of biopsies (8).

	Not done	Cystoscopy with hydrodistension		
		Normal	Glomerulations ^a	Hunner's lesion ^b
Biopsy				
Not done	XX	1X	2X	3X
Normal	XA	1A	2A	3A
Inconclusive	XB	1B	2B	3B
Positive ^c	XC	1C	2C	3C

^a Cystoscopy: glomerulations grade 2-3.

^b With or without glomerulations.

^c Histology showing inflammatory infiltrates and/or detrusor mastocytosis and/or granulation tissue and/or intrafascicular fibrosis.

2.7.3 Pathogenesis

There are many different hypotheses about the causes of BPS/IC.

Infection. No micro-organism has been found to be the cause despite the extensive use of sophisticated microbiological detection methods. Although it has been suggested that fastidious bacteria may be responsible (14), no immunological evidence of recent or remote bacterial infection has been found (15). The results of viral culture (16,17) and polymerase chain reaction methods (18,19) have been just as disappointing. Although urine culture from a few IC patients has contained bacteria, antibiotic treatment has been ineffective. Nevertheless, the possibility of a microbiological contribution has not been looked upon as a 'closed book', including *Helicobacter pylori*, though no *H. pylori* DNA has been detected in bladder biopsies (20).

Inflammation is an essential part of classic IC, with pancystitis and perineural inflammatory infiltrates of lymphocytes and plasma cells (17). Inflammation is scant in non-ulcer IC (6).

Mast cell activation. Mast cells are multifunctional immune cells that contain highly potent inflammatory mediators, such as histamine, leukotrienes, serotonin and cytokines (21). Many of the symptoms and findings in classic IC, such as pain, frequency, oedema, fibrosis and neovascularization in the lamina propria, may be due to the release of mast cell-derived factors. There is a ten-fold increase in the mast cell count in bladder tissue from patients with classic IC compared with controls. In non-ulcer IC, however, the mast cell count is normal or only slightly increased (6,21,22).

Urothelial dysfunction/glycosaminoglycan (GAG)-layer defects. All patients with IC present with fragility of the bladder mucosa, expressed as fissures or rupture of the bladder urothelium on distension (mucosal cracking). In classic IC, the presence of granulation tissue indicates a reparative process (23). In patients with classic IC, urothelial detachment and gross defects of the urothelial lining are characteristic findings. However, in some non-ulcer IC patients, multiple superficial defects are seen after bladder distension (23), including widened tight

junctions and increased permeability (24,25). These changes could be consistent with defects in the GAG-layer that expose the submucosal nerve filaments to noxious chemicals in urine (26,27). Urinary uronate and sulphated GAG levels are increased in patients with severe BPS/IC, suggesting that such substances may become useful markers for monitoring (28).

Autoimmune mechanisms. Numerous studies of autoantibodies have been performed since the 1970s in patients with IC (29), but the findings have been far from specific. Some of the clinical and histopathological characteristics are similar to other autoimmune phenomena. Antinuclear antibodies have been described (30,31), which has led to the hypothesis of a lupus-like reaction (32,33). In fact, only some BPS patients demonstrate autoantibodies and the proposal that autoantibody titres could reflect disease severity is untested (34).

Immune deposits in bladder wall vasculature were found by Mattila (35), while other studies by the same group have implicated complement activation (36). Immunohistochemical and cytofluorometric analyses of the bladder mucosa have highlighted differences between classic and non-ulcer IC patients. In classic IC, intense T-cell infiltrates and B-cell nodules were seen, whereas only some T-cell infiltration was observed in non-ulcer IC (37). The inadequate description of patients in many studies, particularly when it comes to subtyping IC patients, has made it difficult to interpret data. Systemic aspects, especially the potential association with Sjögren's syndrome, are interesting features of BPS/IC (38).

Nitric oxide metabolism. Inevitably, nitric oxide synthetase activity has been scrutinized (39). Oral administration of L-arginine (40) has been shown to increase nitric oxide-related enzymes and metabolites in the urine of patients with BPS/IC (41). However, the relevance of this finding is not clear. An intriguing fact is that evaporation of nitric oxide from the urine is dramatically increased in patients with classic disease, as well as during periods of symptoms decreased by treatment; however, patients with non-ulcer IC, have similar nitric oxide levels to controls (42). To further illustrate the complexity involved, it has been suggested that inducible nitric oxide synthetase-dependent nitric oxide production may have a role in epithelial barrier dysfunction in cats with feline interstitial cystitis (43).

Neurobiology. An increase in the sympathetic innervation and activation of purinergic neurotransmission has been reported in BPS patients. The S-100 family of proteins appears in Schwann cells of the peripheral nervous system (44). Decreased levels of S-100 protein were found in non-ulcer BPS patients compared with controls (45). However, this finding conflicts with that of Hohenfellner et al. (46), who used 'polyclonal antihuman protein gene product 9.5 antibody' and found that the overall nerve content increased in IC patients compared with controls. They did not subtype their patients into classic and non-ulcer forms.

Tyrosine hydroxylase is the rate-limiting enzyme for all catecholamine synthesis. An increase in tyrosine hydroxylase immunoreactivity has been described in bladder tissue from IC patients but not in controls (47); this could be interpreted as a sign of increased sympathetic outflow. Recent reports have suggested that autonomic responses and CNS processing of afferent stimuli are altered in patients with CPP/BPS/IC (48,49). The distinctive ultrastructural appearance of specimens from patients with non-ulcer IC prompted Elbadawi and Light to hypothesize neurogenic inflammation as a trigger to a cascade of events (50).

Toxic agents. Toxic constituents in the urine may cause injury to the bladder in BPS. One hypothesis is that heat labile, cationic urine components of low molecular weight may exert a cytotoxic effect (51). Defective constitutive cytokine production may decrease mucosal defences to toxic agents (52). Tamm-Horsfall protein is a factor whose protective function may be due to its sialic acid content, which is compromised in BPS/IC individuals (53).

Hypoxia. A decrease in the microvascular density in the suburothelium has been observed (54). A recent study, found that bladder perfusion decreased with bladder filling in IC patients, but that the opposite occurred in controls (55). Hyperbaric oxygen therapy has been on trial in an RCT; a total of 30 treatment sessions of hyperbaric oxygenation appeared to be safe, effective and feasible (56).

Complex pathogenic interactions. In recent years, more complex, multifaceted mechanisms have been proposed. Theoharides et al. have shown that activation of mast cells in close proximity to nerve terminals can be influenced by oestradiol as well as corticotrophin-releasing hormone (57). Okragly et al. found elevated levels of tryptase, nerve growth factor, neurotrophin-3 and glial cell line-derived neurotrophic factor in IC compared with controls (58). These findings prompted suggestions that IC may result from interactions between the nervous, immune and endocrine systems. Recently, it was proposed that the epithelial distribution of mast cells in classic IC could be explained by the epithelial co-expression of stem cell factor and interleukin-6 (IL-6). According to Abdel-Mageed et al., IC patients showed an increased expression of p65, a nuclear factor

kappa B subunit (59). Subsequent data has shown a five-fold increase in the expression of the gene for IL-6 after activation of nuclear factor-kappa B (61), although IL-6 is a ubiquitous cytokine.

2.7.4 Epidemiology

Reports of the prevalence of BPS/IC have varied tremendously. However, it should be remembered when comparing studies that most of them have used only symptomatic diagnostic criteria and/or have different study populations. The first systematic study by Oravisto et al. in 1975 found that IC affected approximately 10/100,000 (18/100,000 in women) of the population in Finland (62), with rather similar findings reported 15 years later in the USA (although figures were demonstrated to be dependent on the method of evaluation) (63), as well as in 1995 in the Netherlands with a prevalence of 8-16/100,000 (64). However, other reports claim that the prevalence of IC is underestimated and may exceed 0.5% among adults in the USA (65), with recent US reports suggesting that 50-60/100,000 may be affected (66). Thirty years after the Oravisto study, Leppilahti et al. (67) reported higher figures in Finland of 239/100,000 clinically confirmed probable IC and 530/100,000 possible/probable IC. These figures suggested that BPS/IC was much more common than previously thought.

Recent reports generally claim higher figures than earlier ones. A recent Austrian study reported the overall prevalence of IC as 306/100,000 women, with the highest value (464/100,000) in middle-aged women (aged 40-59 years) (68). A review has claimed that 20% of women may be affected (69). In contrast, however, the incidence of physician-diagnosed incidence in Olmsted County (MN, USA) was extremely low at 1.1/100,000 (70).

There is a female predominance of about 10:1 (4,62,71,72) and it seems that the disease is more common among Caucasians (72).

The relative proportions of classic and non-ulcer disease are unclear. Messing and Stamey reported that classic IC accounted for about half of all patients with IC (5). The same rate has been reported from Sweden (12). Centres in the USA with large patient databases have found that the classic Hunner-type accounts for 5-10% of cases of BPS (74). Koziol et al. recently presented a very large study from the USA, in which classic IC accounted for approximately 20% of cases (75).

Evidence that BPS may have a genetic component is increasing. According to Parsons (76), 35% of 466 patients with BPS and 33% of 166 patients with urethral syndrome reported urgency/frequency problems in female relatives. Warren et al. (77) surveyed 2,058 patients of the Interstitial Cystitis Association (ICA) for first-degree relatives with IC and found a higher prevalence than in the general population. The authors also determined the concordance of IC among ICA twins (78); among the co-twins of eight monozygotic twin respondents, five had probable or confirmed IC, while none of 26 dizygotic co-twins were affected.

BPS/IC has significant economic costs. Excluding indirect costs, the incremental medical cost attributable to this symptom complex in the USA has been estimated to more than \$100 million/year (63).

2.7.5 Association with other diseases

An association between BPS and inflammatory bowel disease, systemic lupus erythematosus, irritable bowel syndrome, fibromyalgia and panic disorders has been reported (79-82). An excellent review has explored comorbidities of BPS/IC with other unexplained clinical conditions presented in the literature (83). The review found a significant overlap of symptoms, suggesting a common stress response pattern, including an increased sympathetic nervous system activity in a subset of patients with many associated conditions. In this context it is interesting to note that feline interstitial cystitis leads to increased corticotrophin-releasing factor activity and decreased adrenocortical reserve.

2.7.6 Diagnosis

The diagnosis of BPS is made using symptoms, examination, urine analysis, cystoscopy with hydrodistension and biopsy (see Figure 3). Patients present with characteristic pain and urinary frequency, which is sometimes extreme and always includes nocturia.

The character of the pain is the key symptom of the disease:

- Pain is related to the degree of bladder filling, typically increasing with increasing bladder content.
- It is located suprapubically, sometimes radiating to the groins, vagina, rectum or sacrum.
- Pain is relieved by voiding but soon returns (6,84-86).

The differences between the two IC subtypes include clinical presentation and age distribution (12), and they may be discriminated non-invasively (75). The two subtypes respond differently to treatment (87-90) and express different histopathological, immunological and neurobiological features (22,23,37,45,47,91,92).

Classic IC is a destructive inflammation with some patients eventually developing a small-capacity fibrotic bladder or upper urinary tract outflow obstruction. There is no such progression in non-ulcer disease (6,93). Endoscopically, classic IC displays reddened mucosal areas often associated with small vessels radiating towards a central scar, sometimes covered by a small clot or fibrin deposit (6). The scar ruptures with increasing bladder distension, producing a characteristic waterfall-type of bleeding. There is a strong

association between classic IC and reduced bladder capacity under anaesthesia (6,12,94).

Cystoscopy. Non-ulcer IC displays a normal bladder mucosa at initial cystoscopy. The development of glomerulations after hydrodistension is considered to be a positive diagnostic sign. A recent report showed that there was no difference in cystoscopic appearance between patients with non-ulcer IC and women without bladder symptoms about to undergo tubal ligation (95). It has also been noted that glomerulations are not always constant when observed over time (96).

Some maintain that cystoscopy with hydrodistension provide little useful information above the history and physical examination findings (97,98). On the other hand, others have found a strong correlation between pain and cystoscopic findings in patients with untreated IC, with the difference in results compared to other studies possibly due to treatment effects (99). Glomerulations may be involved in the disease mechanism, as such findings are highly associated with overexpression of angiogenetic growth factors in the bladder and neovascularization (100).

The European Society for the Study of IC/PBS (ESSIC) believes objective findings are important and that a standardized scheme of diagnostic criteria would help improve the uniformity and comparability of different studies (13).

Biopsies are helpful in establishing or supporting the clinical diagnosis of both classic and non-ulcer types of disease (13,23,101). Important differential diagnoses to exclude by histological examination are carcinoma in situ and tuberculous cystitis.

Potassium chloride bladder permeability test has been used in the diagnosis of IC (102), but recent reports have suggested that it lacks discriminating power (103,104). A modified test using less concentrated solution has been suggested. This test, though painless in contrast to the original procedure, decreased the maximum cystometric volume in 90% of patients with BPS/IC, but not in controls (105). Furthermore, it has been suggested that the potassium sensitivity test can help to predict the response to GAG treatment (106).

Symptom scores may help to describe symptoms in an individual patient and as outcome measures. The O'Leary-Sant Symptom Index, also known as the Interstitial Cystitis Symptom Index (ICSI) has recently been validated successfully in a large study (107).

2.7.7 Biological markers. It is an attractive idea to support or even better to confirm the clinical diagnosis using a biological marker. Finding a universally helpful one is hampered by heterogeneity within the diagnostic group of BPS and by usually making a diagnosis merely on symptoms. Many candidate markers have been suggested. One of the most interesting is antiproliferative factor, which is present in BPS/IC and is associated with downregulation of heparin-binding epidermal growth factor-like growth factor (108). Nitric oxide is interesting because of its ability to discriminate classic from non-ulcer disease with minimal invasiveness (42).

2.7.8 IC in children and males

According to NIDDK criteria, children aged under 18 years is an exclusion criterion. However, occasional cases of BPS of both subtypes have been identified in younger patients (109). There is increasing evidence that very young individuals and children may also be affected, though prevalence figures are low (110). Thus, BPS/IC cannot be excluded on the basis of age.

There is a marked female predominance with a female to male ratio of 10:1. However, the diagnosis must also be considered in men presenting with relevant symptoms (111). It has been argued that many men diagnosed with chronic prostatitis may present with signs consistent with NIDDK criteria of BPS/IC and that these diagnoses are inter-related (112,113). However, differences in urinary markers suggest that BPS/IC and CPP/CP may be different disorders with distinct pathophysiologies (114).

2.7.9 Medical treatment

Analgesics. Since pain is often a dominant symptom, many patients will try commonly used analgesics at some stage of disease. However, pain relief is disappointing because the visceral pain experienced in BPS/IC responds poorly to analgesic drugs. No systematic studies have been presented on conventional analgesics. Short-term opioids may be indicated for breakthrough or exacerbated pain and periodic flare-ups.

Long-term opioids may be considered after all other available therapeutic options have been exhausted. Urologists should obtain informed consent, arrange for regular follow-up, and be prepared to recognize opioid-induced side effects (115). Because BPS/IC is a chronic disease, long-term opioids should be used only exceptionally and under close surveillance.

Corticosteroids. Reports on outcome with corticosteroid therapy have been both promising (116) and

discouraging (117). Soucy et al. (118) suggest a trial of prednisone (25 mg daily for 1 to 2 months, afterwards reduced to the minimum required for symptom relief) in patients with severe ulcerative IC, which is otherwise unresponsive to conventional treatment. The side effects of steroids can be very serious, making it very difficult to justify their use.

Antiallergics. Mast cells may play a role in IC. Among the substances released by mast cells is histamine. Histamine receptor antagonists have been used to block the H₁ receptor subtype (119) as well as the H₂ receptor (120), with variable results.

Hydroxyzine is a histamine H₁-receptor antagonist, which blocks neuronal activation of mast cells by inhibiting serotonin secretion from thalamic mast cells and neurons (121). Hydroxyzine hydrochloride (Atarax) is usually given, starting with 25 mg at bedtime, increasing to 50 mg/day or if tolerated 75 mg. The most common side effects are sedation and generalized weakness that usually resolve after a period of treatment. In the first series using hydroxyzine, > 90% of patients showed an improvement across the whole range of symptoms. Interestingly, an improvement was noted in associated symptoms including migraine, irritable bowel syndrome and allergies (119).

Although these initial results were supported by a further uncontrolled study (119,122), a prospective RCT of hydroxyzine or sodium pentosanpolysulphate (PPS) compared to placebo failed to show a statistically significant effect (123). However, the study was underpowered, which may be why it failed to demonstrate a statistically significant outcome for either drug compared to placebo. Combination therapy showed the highest response rate of 40%, with a placebo response rate of 13%.

Amitriptyline. The tricyclic antidepressant, amitriptyline, has alleviated symptoms in BPS/IC, probably via mechanisms such as blockade of acetylcholine receptors, inhibition of reuptake of released serotonin and norepinephrine, and blockade of the histamine H₁ receptor. It is also an anxiolytic (124). Several reports have indicated amelioration after oral amitriptyline (4,125,126).

In a prospective RCT study, 48 patients (127) were treated for 4 months with amitriptyline. Drug dosages were escalated in 25 mg increments at 1-week intervals up to a maximum dosage of 100 mg. Amitriptyline significantly improved the mean symptom score, pain and urgency intensity, while frequency and functional bladder capacity improved but were not statistically significant.

In a subsequent, prospective, open-label study (128), a response rate of 64% with an overall mean dose of 55 mg was seen with long-term amitriptyline for 20 months. Patient overall satisfaction was good to excellent in 46%, with significant improvement in symptoms. A therapeutic response was observed in all patients fulfilling NIDDK criteria and those with a clinical diagnosis of IC. Anticholinergic side effects (mouth dryness, weight gain) were common and considered to be a drawback of amitriptyline.

Pentosanpolysulphate sodium (PPS, Elmiron) has been evaluated in double-blind, placebo-controlled studies. PPS is thought to substitute for a defect in the GAG layer. Subjective improvement of pain, urgency, frequency, but not nocturia, was reported in patients taking the drug compared to placebo (129,130). In an open multicentre study, PPS had a more favourable effect in classic IC than in non-ulcer disease (90).

The normal dose is 150-200 mg twice daily between meals. However, absorption is incomplete. An RCT compares 300 mg of PPS with evaluated dosages of 600 and 900 mg in 380 IC patients. Mean ICSI scores improved significantly for all dosages (131). However, treatment response was not dose-dependent but related more to treatment duration. At 32 weeks, about half of all patients were responders. Most adverse events were mild and resolved without intervention.

In contrast, a prospective RCT comparing PPS and hydroxyzine against placebo failed to demonstrate a statistically significant outcome for either drug, though PPS approached statistical significance ($p = 0.064$) (123). Combination therapy showed the highest response rate of 40% compared to 13% with placebo. For patients with an initial minor response to PPS, additional subcutaneous administration of heparin appeared helpful (132).

Antibiotics have a limited role in the treatment of BPS/IC. A prospective RCT pilot study of sequential oral antibiotics in 50 patients found that overall improvement occurred in 12/25 patients in the antibiotic group and 6/25 in the placebo group, while 10 and 5 patients reported an improvement in pain and urgency, respectively. Antibiotics alone or in combination may be associated with decreased symptoms in some patients, but do not represent a major advance in therapy for BPS/IC (133).

Immunosuppressants. Azathioprine, 50-100 mg daily, was given to 38 patients, resulting in disappearance of pain in 22 and urinary frequency in 20 (134). Cyclosporin A (CyA) (135) and methotrexate (136) were initially

evaluated in open studies, with a good effect on pain, but a limited effect on urgency-frequency. More recent studies of CyA have reported promising results (137,138). In 23 patients, daily voidings, maximal bladder capacity and voided volume improved significantly after 1 year of treatment. The effect was maintained throughout 5 years' follow-up, with 20/23 patients reporting no bladder pain. However, symptoms recurred within months of discontinuing CyA.

In a subsequent randomized study (138), 64 patients fulfilling the NIH-criteria were randomized to 1.5 mg/kg CyA twice daily or low-dose (3 x 100 mg) PPS for 6 months. CyA was superior to PPS in all clinical outcome parameters, with the frequency of micturition significantly reduced in CyA-treated patients, and clinical global response rates of 75% (CyA) and 19% (PPS) ($p < 0.001$). However, there were more adverse events in the CyA arm (including induced hair growth, gingival pain and hyperplasia, paresthesias in extremities, abdominal pain, flushing, muscle pain and shaking) and only 29 patients completed the 6-month follow-up in both groups. During CyA therapy, careful follow-up is mandatory, including regular blood pressure measurement and serum creatinine.

Gabapentin / Pregabalin

Gabapentin is an antiepileptic drug, which is used as adjunctive treatment in painful disorders. Gabapentin may reduce the use of co-therapeutics, such as opioids. Two patients with IC showed improved functional capacity and received adequate pain control when gabapentin was added to their medication regimen (139). In an uncontrolled dose-escalation protocol with 21 chronic genitourinary pain patients (140), 10 had improved with gabapentin at 6 months. The study included eight IC patients, of whom five responded to gabapentin.

Pregabalin is an alpha(2)-delta ligand that binds to and modulates voltage-gated calcium channels, exerting its intended effect to reduce neuropathic pain (141). Pregabalin is the second of only two medications that are US FDA-approved for the treatment of neuropathic pain associated with diabetic peripheral neuropathy; it is used for the treatment of post-herpetic neuralgia. Studies on IC are still lacking.

Suplatast tosilate (IPD-1151T) is an oral immunoregulator that suppresses helper T-cell mediated allergic processes. Fourteen women with IC treated with suplatast tosilate reported significantly increased bladder capacity and decreased symptoms after 1 year of treatment. No major side effects occurred and therapeutic effects correlated with a reduction in blood eosinophils, immunoglobulin E and urinary T-cells (142). Comparative controlled data are unavailable.

Quercetin is a bioflavonoid that may be effective in male pelvic pain syndrome. It was first tested in a limited, open-label study with hopeful results (143). Theoharides et al. (144) reported on the dietary supplement CystoProtek formulated from quercetin and the natural GAG components, chondroitin sulphate and sodium hyaluronate. In an uncontrolled study, symptoms were significantly improved in 37 IC patients (NIH-criteria), who had failed all forms of therapy and who took six capsules per day for 6 months. Larger controlled studies are warranted by this result.

Recombinant human nerve growth factor. A small randomized study (145) was performed on 30 patients (NIH-criteria), who received either placebo or one of two dosages (0.1 or 0.3 mg/kg) of recombinant human nerve growth factor weekly for 3 months. Significant improvement was seen after 3 months as measured by subjective improvement and ICSI score, while mast cells were significantly reduced and nerve cells elevated ($p < 0.05$) in a dose-dependent manner. Side effects were arthralgias (5%), myalgias (4%), and myasthenia and asthenia (2%).

The results suggest that recombinant human nerve growth factor is safe and shows preliminary evidence of efficacy in patients with BPS/IC, but further studies are needed to define its role.

2.7.10 Intravesical treatment

Intravesical application of medications establishes high concentrations at the target with few systemic side effects. Disadvantages include the need for intermittent catheterization, which can be painful in IC patients, the cost, and the risk of infection.

Local anaesthetics. There are sporadic reports of successful treatment of IC with intravesical lidocaine (146,147). Alkalinization of lidocaine prior to intravesical application improved pharmacokinetics (148). In an uncontrolled study, significant immediate symptom relief was reported in 94% of patients and sustained relief after 2 weeks in 80%, using instillations of combined heparin and alkalinized lidocaine (40,000 U heparin, 2% lidocaine (160 mg), and 3 mL 8.4% sodium bicarbonate) (149).

Pentosanpolysulphate sodium (PPS) is a glycoprotein aimed at replenishing the GAG layer, which is applied

intravesically due to poor bioavailability following oral administration. A double-blind placebo-controlled study (150) was performed in 20 patients, of whom 10 received intravesical PPS (300 mg in 50 mL of 0.9% saline) twice a week for 3 months and 10 received placebo. At 3 months, four patients in the PPS group and two patients in the placebo group gained significant symptomatic relief. Bladder capacities showed a statistically significant increase only in patients treated with PPS. At 18 months, symptoms were relieved in eight patients, who were still receiving PPS instillations, and in four patients not receiving PPS.

Intravesical heparin was proposed as a coating agent. In an open, prospective, uncontrolled trial (151), 48 IC patients received instillations of 10,000 units in 10 mL sterile water three times per week for 3 months. In over half of the patients studied, intravesical heparin controlled the symptoms, with continued improvement after 1 year of therapy. Kuo et al. (152) reported another uncontrolled trial of intravesical heparin (25,000 units twice a week for 3 months) in women with frequency-urgency syndrome and a positive potassium test. The study included 10 patients with IC, of whom eight reported symptomatic improvement. Baykal et al. (153) evaluated intravesical heparin plus dorsal tibial nerve stimulation in 10 refractory IC patients. Voiding frequency, pain scores and maximum cystometric capacity were significantly better after 2 and 12 months compared to pretreatment values.

Hyaluronic acid (hyaluronan) is a natural proteoglycan aimed at repairing defects in the GAG layer. A response rate of 56% at week 4 and 71% at week 7 was reported in 25 patients treated with hyaluronic acid (154). After week 24, effectiveness decreased, but there was no significant toxicity. Nordling et al. (155) and Kallestrup (156) reported a 3-year follow-up of a 3-month, prospective, non-randomized study evaluating the effect of intravesical hyaluronic acid on BPS/IC symptoms. Of the 20 patients, 11 chose to continue treatment beyond the initial trial, and modest beneficial long-term effects were noted in about two-thirds of patients. Reduction in urinary frequency was less effective and mostly due to an improvement in night-time voids.

Another study (157) demonstrated a similar favourable effect of hyaluronic acid on pain reduction. Forty-eight patients were treated with typical symptoms and a positive potassium (0.4 M) sensitivity test with weekly instillations of 40 mg hyaluronic acid for 10 weeks. Visual analogue scale scores showed symptom relief due to hyaluronic acid therapy, irrespective of bladder capacity. The improvement was particularly evident in patients with a reduction in C(max) \leq 30% compared to patients with a reduction of $<$ 30% with 0.2 M KCl solution ($p = 0.003$).

Chondroitin sulphate. Intravesical chondroitin sulphate (158) demonstrated beneficial effects in patients, who had given a positive potassium stimulation test, in two non-randomized, uncontrolled, open-label pilot studies. Steinhoff (159) treated 18 patients with 40 mL instilled intravesically once a week for 4 weeks and then once a month for 12 months. Thirteen of 18 patients were followed for the entire 13-month study. Twelve of these patients responded to treatment within 3-12 weeks. A total of 6/13 (46.2%) showed a good response, 2/13 (15.4%) had a fair response, 4/13 (30.8%) had a partial response, and 1/13 (7.7%) showed no response.

In a second trial (160), 24 refractory patients with BPS/IC were treated with high-dose (2.0%) chondroitin sulphate instillations twice weekly for 2 weeks, then weekly with 0.2% solution for 4 weeks, and monthly thereafter for 1 year. The average symptom improvement reported in 20 patients completing the trial was 73.1% (range 50–95%). The time to optimum response was 4-6 months. More concentrated 2.0% solution was needed in eight patients to maintain results. A Canadian phase II/III non-randomized, uncontrolled, community-based, open-label efficacy and safety study is underway.

Dimethyl sulphoxide (DMSO) is a chemical solvent and water-soluble liquid that penetrates cell membranes. It is claimed to have analgesic, anti-inflammatory, collagenolytic and muscle relaxant effects. It is also a scavenger of the intracellular OH radical believed to be an important trigger of the inflammatory process. It has been tested empirically and found to alleviate symptoms in IC. DMSO is now a standard treatment.

In a controlled, crossover trial (161), 33 patients received instillations of 50% DMSO solution and placebo (saline). All patients received both regimens, which were administered intravesically every 2 weeks for two sessions of four treatments each. Subjective improvement was noted in 53% of patients receiving DMSO versus 18% receiving placebo, and objective improvement in 93% and 35%, respectively. Other uncontrolled trials with DMSO have reported response rates of 50-70% for a period of between 1 and 2 months (162). Rossberger et al. (163) evaluated the discomfort and long-term of DMSO instillations in a total of 28 patients. Side effects were not more common or pronounced in patients with classic compared to non-ulcer disease. After DMSO instillations, a residual treatment effect lasting 16-72 months could be seen.

DMSO is contraindicated during urinary tract infections or shortly after bladder biopsy. It temporarily causes a garlic-like odour. Because there has been a case report in which DMSO treatment may have caused pigmented eye lens deposits (164), ophthalmic review should be considered during treatment.

Bacillus Calmette Guérin. The tuberculosis vaccine, *Bacillus Calmette-Guérin* (BCG), is used for its immunomodulatory properties in the intravesical treatment of superficial bladder carcinoma. In 1997, a small prospective, double-blind pilot study on intravesical BCG demonstrated a 60% BCG versus 27% placebo response rate in 30 patients who received six weekly instillations of Tice strain BCG or placebo (165). In a subsequent 24-33 months' follow-up report, eight of the nine responders reported favourably. BCG did not worsen symptoms in non-responders (166). However, these results are at variance with two controlled trials.

In a prospective, double-blind crossover trial of BCG and DMSO (89), BCG treatment failed to demonstrate any benefit. Another randomized, placebo-controlled, double-blind trial on 260 refractory IC patients (167) reported global response rates of 12% for placebo and 21% for BCG ($p = 0.062$). Small improvements were observed for all secondary outcomes (voiding diary, pain, urgency, symptom indexes and adverse events), some of which were greater with BCG, but with only borderline statistical significance. In a subsequent study (168), 156 non-responders from both groups were offered treatment with open-label BCG. The low response rate (18%) for BCG in this series is a further argument against the routine use of BCG as treatment for BPS/IC.

Vanilloids disrupt sensory neurones (169). Resiniferatoxin (RTX) is an ultrapotent analogue of the chilli pepper extract capsaicin, causing less pain on instillation and therefore no anaesthesia. Chen et al. (170) investigated RTX tolerability (0.05 μ M or 0.10 μ M) in 22 BPS/IC patients versus placebo. The most commonly reported adverse event was pain during instillation (RTX > 80.0%, placebo 25.0%) but no serious adverse events were reported.

In a small RCT on 18 patients with hypersensitive bladder disorder and pain (171), RTX significantly reduced mean frequency, nocturia and pain scores by about 50%. In another study of seven patients with detrusor hyperreflexia, RTX improved urinary frequency, incontinence and bladder capacity (172). In a small open-label study with single-dose RTX in patients with frequency and urgency (173), RTX significantly improved lower urinary tract symptoms, urodynamic parameters, and quality of life for up to 6 months.

These results are in contrast with an RCT in 163 BPS/IC patients randomly assigned to receive a single intravesical dose of 50 mL of either placebo or RTX (in the dosages 0.01, 0.05, or 0.10 μ M) (174). RTX resulted in a dose-dependent increase in instillation pain, but otherwise was well tolerated. It did not improve overall symptoms, pain, urgency, frequency, nocturia, or average void volume during 12 weeks' follow-up.

More favourable results were reported from a prospective study on multiple intravesical instillations of RTX (175) (0.01 μ M once weekly for 4 weeks). Among 12 patients (one drop-out for severe pain), the overall satisfactory rate was 58.3%, with several scales of symptom and quality of life significantly improved after RTX treatment. There was no significant increase in functional bladder capacity or change in urodynamic parameters.

Modification of urine pH. A prospective, randomized, double-blind cross-over study was performed in 26 women, who received instillations of various pH values. There was no evidence that changes in urinary pH affected the pain associated with IC (176).

2.7.11 *Interventional treatments*

Bladder distension. A frequently cited report by Bumpus et al. (177) claims that hydrodistension achieved symptom improvement in 100 patients over several months. However, the study did not define either patient population or symptoms and the methods used were inadequately described. Reports by Ormond (178) and Longacre (179) were just as vague during the 1930s. In 1957, an uncontrolled retrospective study was presented by Franksson (180), who treated 33 patients with repeated, up to 10-fold, distensions. Twelve patients had improved symptoms for up to 4 weeks, in 14 patients for up to 6 months, and in seven patients for up to 1 year. British studies from the 1970s reported contradictory results. Dunn et al. (181) claimed to have achieved complete absence of symptoms in 16/25 patients during a mean follow-up of 14 months using the Helmstein method (182), where an intravesical balloon is distended at the level of systolic blood pressure for 3 hours. Bladder rupture occurred in two cases. These results disagree with those of Badenoch (116), who failed to note any improvement in 44/56 patients after hydrodistension. Twenty years later, McCahy (183) rejected balloon hydrodistension because of inefficacy and a complication rate of 20%. In the recent literature, bladder necrosis following hydrodistension is extremely rare (184).

In 2002, Glemain et al. (185) reported an uncontrolled study on 65 IC patients treated by 3-hour balloon hydrodistension. Treatment efficacy in the 33 retrospectively and 32 prospectively studied patients was 38% and 60% at 6 months, and 22% and 43% at 1 year, respectively. Results were superior for bladder capacities above 150 mL.

Ottem and Teichmann (2006) reported a retrospective study of 84 BPS/IC patients (97), among which 56% reported short-lived improvement from hydrodistension. Rose et al. investigated bladder distension using electromotive drug administration (EMDA) (186,187), as an alternative to general anaesthesia. Among 11 patients, the distention capacity achieved by EMDA was nearly identical to that in the operating room and

cystoscopic findings were similar. Yamada et al. (188) reported on repeated hydrodistension in 52 IC patients (NIH-criteria). Under epidural anaesthesia, the bladder was repeatedly distended to maximal capacity and distension was repeated on the following day for 30 minutes. Five patients were classified as good responders, 30 as moderate responders and 17 as poor responders. Overall, hydrodistension was effective for about 70% of patients for more than 3 months without serious complications.

According to a study by Erickson et al. (189), the median symptom score for newly diagnosed patients decreased after distention, but only a few patients had at least 30% symptom improvement. Bladder distention altered levels of urine antiproliferative factor and heparin-binding epidermal growth factor-like growth factor towards normal. However, the mechanism of symptom relief after distention remains unknown.

In a retrospective review of 185 patients who underwent hydrodistension (190), results failed to identify any statistically significant differences in objective findings (anaesthetic capacity, glomerulations) following distension, or any therapeutic benefits, when patients were categorized according to presenting symptoms.

Although bladder hydrodistension is a common treatment for BPS/IC, scientific justification is scarce. It represents a diagnostic tool, but has a limited therapeutic role.

Electromotive drug administration (EMDA) enhances tissue penetration of ionized drugs by iontophoresis. Adapted for the bladder, it uses a transurethral anode and a suprapubic skin cathode. EMDA is expensive and the subject of uncontrolled studies only.

Six IC patients were treated with EMDA using lidocaine (1.5%) and 1:100,000 epinephrine in aqueous solution, while the bladder was dilated to maximum tolerance (191). Significant bladder enlargement was achieved and voiding symptoms and pain decreased. In four patients, the results were reported as 'durable'. Rosamilia et al. (192) treated 21 women using EMDA with lidocaine and dexamethasone, followed by cystodistension. A good response was seen in 85% of patients at 2 weeks, with 63% still responding at 2 months. Complete resolution of pain was achieved in 25% of patients reviewed at 6 months. Using a similar technique, Riedl et al. (193) noted complete resolution of bladder symptoms in 8/13 patients lasting 1-17 months. Partial or short-term improvement was observed in three patients. Two patients experienced aggravated pain for several days after therapy. A 66% increase in bladder capacity was observed. Upon symptom recurrence, treatments were repeated with equal efficacy in 11 patients.

Transurethral resection (TUR) coagulation and LASER. Endourological ablation of bladder tissue aims to eliminate urothelial, mostly Hunner, lesions. In a case report, Kerr (194) described a transurethral resection of a 1-cm ulcer in a woman who experienced symptom resolution for 1 year. Subsequently, Greenberg et al. (71) reported on 77 patients with Hunner ulcers treated over a 40-year period: 42 were managed conservatively, seven underwent fulguration and 28 were treated by TUR in a non-randomized fashion. Fulguration improved symptoms in 5/7 patients. All patients experienced symptom recurrence in less than 1 year and efficacy was not superior to non-surgical treatment.

In another series of 30 classic IC patients (195), complete TUR of visible lesions resulted in an initial disappearance of pain in all patients and a decrease in frequency in 21 patients. A relapse was noted in one-third of patients after 2-20 months, while the remaining two-thirds were still pain-free after 2-42 months. The same group recently reported the largest series of patients with classic IC treated by complete TUR of all visible ulcers (196). A total of 259 TURs were performed on 103 patients. Ninety-two patients experienced amelioration, with symptom relief lasting longer than 3 years in 40%, while most of the remaining patients responded well to subsequent TUR.

Transurethral application of the neodymium-yttrium-aluminium-garnet (Nd-YAG) laser is suggested as an alternative to TUR for endoscopic treatment in IC. Shanberg et al. (197) treated five refractory IC patients, four of whom demonstrated cessation of pain and frequency within several days. Follow-up at 3-15 months revealed no relapse except mild recurrent voiding symptoms. This series was extended to 76 patients treated at two institutions (198). Although 21 of 27 patients with Hunner ulcers noted symptom improvement, 12 experienced relapse within 18 months. In the group without ulcers, only 20 of 49 patients improved, of whom 10 required further therapy within 1 year.

In a later study, 24 patients with refractory classic IC underwent ablative Nd-YAG laser ablation of Hunner's ulcers (199). All patients showed symptom improvement within days without complications. At 23 months, mean pain and urgency scores, nocturia and voiding intervals had improved significantly. However, relapse in 11 patients required up to four additional treatments. Endourological resections are not applicable to non-ulcer IC. These techniques may provide long-term alleviation of symptoms, but none are a cure for the disease. Controlled studies are still lacking. Endourological resections are not applicable to non-ulcer BPS/IC. These techniques may provide long-term alleviation of symptoms, but none of them cure the disease. Controlled studies are still lacking.

Botulinum toxin A (BTX-A) may have an anti-nociceptive effect on bladder afferent pathways, producing both

symptomatic and urodynamic improvements (200). Thirteen BPS/IC patients were injected with 100-200 IU of Dysport or BTX into 20 to 30 sites submucosally in the trigone and floor of the bladder. Overall, 9 (69%) patients noted a subjective improvement and ICSI scores improved by 70% ($p < 0.05$). There were significant decreases in daytime frequency, nocturia, and pain, and a significant increase in first desire to void and maximal cystometric capacity. However, these results are in contrast with another study of BTX in 10 patients with BPS/IC (201). One hundred units were injected suburothelially into 20 sites in five patients, while 100 units were injected into the trigone in the remaining five patients. None of the patients became symptom-free; two patients showed only limited improvement in bladder capacity and pain score.

Hyperbaric oxygen (HBO). In a prospective pilot study, six patients underwent 30 sessions of 100% hyperbaric oxygen inhalation and were followed up over 15 months. Four patients rated the therapeutic result as excellent or good, while two showed only short-term amelioration (202).

In a subsequent double-blind, sham-controlled study (203), 3/14 patients on HBO and no control patients were identified as responders ($p < 0.05$). At 12 months, three patients (21.4%) still reported a treatment response. Hyperbaric oxygenation resulted in a decrease of baseline urgency and pain ($p < 0.05$). ICSI scores decreased from 26 to 20 points in patients on HBO, while sham treatment did not result in any improvement.

These results suggest that HBO is a safe and feasible therapeutic approach, with moderate effects on a small subgroup of BPS/IC patients. Disadvantages include high costs, limited availability of treatment sites and time-consuming treatment.

2.7.12 *Treatments of limited efficacy and absence of recent publications*

Cimetidine. The H₂-blocker cimetidine has been reported to improve symptoms in bladder pain syndrome (204). Thirty-six patients were enrolled in a double-blind clinical study with oral cimetidine versus placebo for 3 months. Patients receiving cimetidine showed a significant improvement in symptom scores, pain and nocturia, although histologically the bladder mucosa showed no qualitative changes in either group (205).

Prostaglandins. Misoprostol is a prostaglandin that regulates various immunological cascades. Twenty-five IC patients received 600 µg of misoprostol daily for 3 months, with responders treated for a further 6 months. At 3 months, 14 had significantly improved, with 12 showing a sustained response after a further 6 months. However, the incidence of adverse drug effects was 64% (206).

L-arginine. Oral treatment with L-arginine, the substrate for nitric oxide synthase, has been reported to decrease BPS/IC-related symptoms (207-209). Nitric oxide has been shown to be elevated in patients with IC (210). However, others could not demonstrate either symptomatic relief or change in nitric oxide production after treatment (211,212).

Anticholinergics. Oxybutynin is an anticholinergic drug used in overactive detrusor dysfunction. Intravesically administered oxybutynin was combined with bladder training in one study, with improvement of functional bladder capacity, volume at first sensation and cystometric bladder capacity (213). However, the effect on pain was not reported.

Duloxetine inhibits both serotonin and noradrenaline reuptake. In an observational study, 48 women were prospectively treated with duloxetine for 2 months following an up-titration protocol to the target dose of 2 x 40 mg duloxetine per day over 8 weeks (214). Duloxetine did not result in significant improvement of symptoms. Administration was safe, but tolerability was poor due to nausea. Based on these preliminary data, duloxetine cannot be recommended as a therapeutic approach for BPS/IC.

Clorpactin is a detergent of hypochloric acid previously used to treat IC (215-219). Due to high complication rates (217,220), clorpactin instillations can no longer be recommended.

2.7.13 *Alternative and complementary treatments*

Behavioural bladder training techniques are attractive for BPS/IC patients with predominant symptoms of frequency/urgency but hardly any pain. Parsons et al. (221) included 21 selected BPS/IC patients on a protocol, which focused on progressively increasing micturition intervals. Fifteen patients reported a 50% decrease in urgency, frequency and nocturia, and there was a moderate increase in bladder capacity. Chaiken et al. (222) retrospectively analysed 42 patients, who had been instructed in diary keeping, timed voiding, controlled fluid intake and pelvic floor muscle training. After 12 weeks, voiding intervals increased by a mean of 93 minutes and daily micturitions were reduced by an average of nine voids. Overall, 88% of the patients reported markedly improved or improved symptoms.

Diet. Dietary restrictions are among the many physical self-care strategies found among BPS/IC patients (223).

In an analysis of the Interstitial Cystitis Data Base (ICDB) cohort study, special diets were among the five most commonly used therapies (224). Bade et al. (225) found that IC patients consumed significantly less calories, fat and coffee, but more fibre. Scientific data on a rationale for such diets are unavailable.

The concentration of some metabolites and amino acids appears to be changed in IC (226). A study of the metabolism of the arylalkylamines (tryptophan, tyrosine, tyramine, phenylalanine) in 250 patients revealed an inability to synthesize normal amounts of serotonin and a noradrenaline metabolite. In this study, dietary restriction of acid foods and arylalkylamines lessened the symptoms, but did not alter specific abnormalities in dopamine metabolism.

In another, non-randomized, prospective study of BPS/IC patients with nutrition-related exacerbations, calcium glycerophosphate was reported to ease food-related flares (227). The observed efficacy seems little better than would be expected with placebo.

Overall, dietary management is a common self-care strategy in BPS/IC and offers a cost-effective therapeutic approach. Comprehensive instructions on how to identify individual trigger foods are given in the IC-Network Patient Handbook (228). However, scientific data are limited and dietary restriction alone does not produce complete symptomatic relief.

Acupuncture. In non-curable and agonizing diseases like BPS/IC, desperate patients often try complementary medicines, such as acupuncture. However, scientific evidence for such treatments is often poor, with contradictory results from a few low-evidence reports on acupuncture, with any effects appearing to be limited and temporary.

A significant increase in capacity occurred after acupuncture in 52 women with 85% reporting an improvement in frequency, urgency and dysuria and symptoms (229). However, at follow-up at 1 and 3 years, these effects were no longer detectable and the authors concluded that repeated acupuncture was necessary to maintain beneficial effects (230).

In a non-randomized comparison in females with urethral syndrome, 128 patients treated by acupuncture and traditional Chinese medicine were compared to 52 patients treated by Western medicine as controls. Efficacy rates and urodynamic parameters were significantly better in the acupuncture group (231). In contrast, in a prospective study on the effect of acupuncture in IC (232), no differences in frequency, voided volumes or symptom scores were noted and only one patient improved for a short period of time.

Hypnosis is a therapeutic adjunct in the management of cancer, surgical disease and chronic pain. Although used in urological patients (233,234), there is no scientific data on its effect on IC symptoms.

Physiotherapy. General body exercise may be beneficial in some BPS/IC patients (235). An uncontrolled trial of transvaginal manual therapy of the pelvic floor musculature (Thiele massage) in 21 BPS/IC patients with high-tone dysfunction of the pelvic floor resulted in statistically significant improvement on several assessment scales (236). Langford (237) prospectively examined the role of specific levator ani trigger point injections in 18 females with CPP. Each trigger point was identified by intravaginal palpation and injected with 5 mL of a mixture of 10 mL of 0.25% bupivacaine, 10 mL of 2% lidocaine and 1 mL (40 mg) of triamcinolone. A total of 13 out of 18 (72%) women improved with the first trigger point injection, with 6 out of 18 (33%) women completely pain-free.

Intravaginal electrical stimulation was applied to 24 women with CPP in the form of ten 30-minute applications, two or three times per week. Stimulation was effective in alleviating pain, as evaluated at the end of treatment and 2 weeks, 4 weeks and 7 months after completion of treatment ($p < 0.05$). There were significantly fewer complaints of dyspareunia following treatment ($p = 0.0005$) (238).

2.7.14 Surgical treatment

When all efforts fail to relieve disabling symptoms, surgical removal of the diseased bladder is the ultimate option (239-242). Three major techniques of bladder resection are common:

- supratrigonal (i.e. trigone-sparing) cystectomy
- subtrigonal cystectomy
- radical cystectomy including excision of the urethra.

All techniques require substitution of the excised bladder tissue, mostly performed with bowel segments.

Techniques without bladder removal. As early as 1967, Turner-Warwick reported that mere bladder augmentation without removal of the diseased tissue was not appropriate (243). Sporadic reports that unresected IC bladders cease to cause symptoms when excluded from the flow or urine are scarce (5,244).

Supratrigonal cystectomy with subsequent bladder augmentation represents the most favoured continence-

preserving surgical technique. Various intestinal segments have been used for trigonal augmentation, including ileum (116,245-252), ileocecum (251-258), right colon (116,252,259), and sigma (246,248,249,254,258). Substituting gastric segments (260,261) seems to be less helpful because the production of gastric acids may maintain dysuria and persistent pain.

The therapeutic success of supratrigonal cystectomy has been reported in many studies. In 1966, von Garrelts reported excellent results in 8/13 patients with a follow-up of 12-72 months (248). In 1977, Bruce et al. achieved satisfactory relief of IC symptoms by ileocystoplasty and colcystoplasty in eight patients (246). Dounis and Gow reported seven IC patients whose pain and frequency were considerably improved after supratrigonal cystectomy with ileocecal augmentation (262).

In 1991, Kontturi et al. employed segments of colon and sigmoid colon in 12 cases (258). All five patients augmented with sigmoid colon remained symptom-free over 4.7 years of follow-up. Two of seven cases augmented with colon required secondary cystectomy with formation of an ileal conduit. Nielsen et al. reported a series of eight patients undergoing supratrigonal cystectomy with ileocaecocystoplasty. While symptoms resolved in two patients, treatment failure in another six patients necessitated secondary cystectomy and ileal conduit formation (253).

Linn et al. (263) followed six BPS/IC patients after supratrigonal cystectomy with an ileocecal augmentation for a period of 30 months and reported that all patients were symptom-free and voided spontaneously.

In 2002, Van Ophoven et al. (239) reported the long-term results of trigone-preserving cystectomy and consecutive orthotopic substitution enteroplasty in 18 women with IC, using ileocaecal (n = 10) or ileal (n = 8) segments. At a mean follow-up of nearly 5 years, 14 patients were completely pain-free, 12 voided spontaneously and 15 had complete resolution of dysuria. Ileocaecal bowel segments showed superior functional results, since in the group augmented with ileum, three patients required self-catheterization and one a suprapubic catheter. Overall, surgery achieved a significant improvement in diurnal and nocturnal frequencies, functional bladder capacity and symptom scores, with only two treatment failures.

In more recent reports with longer follow-ups, the debate on the outcome of BPS/IC patients undergoing cystectomy continues and results vary greatly between different surgeons and patient populations.

Chakravarti (264) presented a retrospective review of 11 patients, who had undergone a trigone-preserving orthotopic substitution caecocystoplasty for intractable classical IC and were followed up for a mean period of 9 years. All had symptomatic relief and an increase in bladder capacity to normal. There was no mortality and minimal post-operative morbidity, with two patients requiring intermittent self-catheterization due to high residual volumes. No significant urinary reflux or metabolic complications were noted. However, two patients required a cystectomy after 4 and 6 years, respectively, due to recurrent trigonal disease in one patient and urethrotigonal hypersensitivity following intermittent self-catheterization in the other. One patient developed an advanced adenocarcinoma in the caecal segment 7 years after the primary operation.

Blaivas et al. (265) reported less favourable results. Long-term outcomes of augmentation enterocystoplasty or continent urinary diversion were analysed in 76 patients with benign urological disorders, including seven patients with a clinical diagnosis of IC. The BPS/IC patients all failed surgical treatment because of persistent pelvic pain and failure to achieve adequate bladder capacity rather than because of incontinence. The authors currently consider BPS/IC to be a contraindication for enterocystoplasty.

In contrast, Navalon et al. (266) reported a 32-month follow-up of four women suffering refractory IC who underwent supratrigonal cystectomy with orthotopic substitution iliocystoplasty. Suprapubic pain disappeared in all cases, as well as lower urinary tract symptoms, with good control of urinary frequency day and night in the immediate post-operative period. All patients reported high satisfaction with the outcome.

Subtrigonal cystectomy. Although less popular, subtrigonal cystectomy has also been reported (263,267-270). Subtrigonal resection has the potential of removing the trigone as a possible disease site, but at the cost of requiring ureteral reimplantation with associated risks of leakage, stricture and reflux. Nurse et al. reported trigonal disease in 50% within their cohort (13/25) and blamed surgical failures on the trigone left in place (271). In contrast, Linn et al. indicated that the level of resection was not solely responsible for treatment success. While completely curing six patients by supratrigonal resection, there were three failures among 17 subtrigonal resections and half of the successful subtrigonal resections required self-catheterization to support voiding of the ileocecal augmentate (263). A recent report on female sexuality after cystectomy and orthotopic ileal neobladder (272) included eight patients. Pain was relieved in all eight patients, but only one patient regained a normal sexual life post-operatively.

Selecting patients and technique. Bladder pain syndrome /Interstitial cystitis is benign and does not shorten life, so that operative procedures rank last in the therapeutic algorithm. However, severely refractory patients should not have to tolerate unsuccessful conservative treatments for years when surgical options are available.

Detailed counselling and informed consent must precede any irreversible type of major surgery, which

should only be undertaken by experienced surgeons. The choice of technique will be influenced by the experience of the surgeon. The appropriate extent of tissue resection should be based on the endoscopic and histopathological findings. Some surgeons recommend pre-operative cystoscopy and bladder capacity as a prognostic parameter for operative success (244). Responders and failures following orthotopic substitution differed in mean pre-operative bladder capacity (200 mL vs 525 mL, respectively) (253). These findings were supported by Peeker et al. (273), who found that patients with end-stage classic IC had excellent results following ileocystoplasty while patients with non-ulcer disease were not helped. These results have recently been confirmed by another report from the same institution. A retrospective analysis of 47 patients fulfilling the NIH criteria, who underwent reconstructive surgery using various techniques during 1978-2003 (274), resulted in complete symptom resolution in 32/34 patients with classic Hunner-type disease, but only 3/13 patients with non-ulcer disease.

Cystectomy with formation of an ileal conduit stills ranks first in current US practice trends in surgical IC therapy (275). For cosmetic reasons, however, techniques of continent diversion are preferred, particularly in younger patients. After orthotopic bladder augmentation, particularly when removing the trigone, voiding may be incomplete and require intermittent self-catheterization. Patients considering these procedures should be advised and must be considered capable of performing, accepting and tolerating self-catheterization. For patients with BPS/IC who develop recurrent pain in the augmented bladder or continent pouch after enterocystoplasty or continent urinary diversion, Elzawahri (276) recommends retubularization of a previously used bowel segment to form a urinary conduit. For younger patients, it may be important to know that pregnancies with subsequent lower-segment Caesarean section after ileocystoplasty have been reported (277).

Reconstructive surgery for refractory BPS/IC is an appropriate last resort only for well-selected patients with refractory end-stage disease. The decision to embark on major reconstructive surgery should be preceded by a thorough pre-operative evaluation, with an emphasis on assessment to determine the relevant disease location and subtype.

A summary of the treatment options for BPS/IC, including a rating of the level of evidence and grade of recommendation (Table 9), is given in Tables 10 and 11. Figure 3 provides an algorithm for the diagnosis and therapy of BPS/IC based on the information discussed above.

Table 9: Explanations of levels of evidence and grades of recommendation for treatment options for BPS/IC outlined in Tables 10 and 11

Level	Type of evidence
1a	Meta-analysis of randomized trials
1b	At least one randomized trial
2a	One well-designed controlled study w/o randomization
2b	One other type of well-designed quasi-experimental study
3	Non-experimental study (comparative study, correlation study, case reports)
4	Expert committee, expert opinion
Grade	Nature of recommendation based on
A	Clinical studies of good quality and consistency including at least one randomized trial
B	Well-conducted clinical studies without randomized trials
C	Absence of directly applicable clinical studies of good quality

Table 10: Medical treatment of BPS/IC

	Level of evidence	Grade of recommendation	Comment
Analgesics	2b	C	Indications limited to cases awaiting further treatment
Corticosteroids	3	C	Corticosteroids not recommended as long-term treatment
Hydroxyzine	1b	A	Standard treatment, even though limited efficacy shown in RCT
Cimetidine	1b	B	Insufficient data
Amitriptyline	1b	A	Standard treatment
Sodium pentosanpolysulphate	1a	A	Standard treatment Data contradictory
Antibiotics	1b	A	Limited role in the treatment of IC

Prostaglandins	3	C	Insufficient data on IC, adverse effects
L-arginine	1b	C	Effect in IC uncertain
Cyclosporin A	1b	A	RCT: superior to PPS but more adverse effects
Duloxetine	2b	C	No effect, tolerability poor
Oxybutynin/tolterodine	3	C	Limited indication in IC
Gabapentin	3	C	Preliminary data so far
Suplatast tosilate	3	C	Preliminary data so far
Quercetin	3	C	Preliminary data so far


RCT = randomized controlled trial; IC = interstitial cystitis; PPS = pentosanpolysulphate sodium.

Table 11: Intravesical, interventional, alternative and surgical treatment of BPS/IC

	Level of evidence	Grade of recommendation	Comment
Intravesical anaesthesia	3	C	
Intravesical PPS	1b	A	
Intravesical heparin	3	C	
Intravesical hyaluronic acid	2b	B	
Intravesical chondroitin sulphate	2b	B	
Intravesical DMSO	1b	A	
Intravesical Bacillus Calmette Guérin	1b	Not recommended	
Intravesical clorpactin	3	Not recommended	Obsolete
Intravesical vanilloids	1b	C	Data contradictory
Bladder distension	3	C	
Electromotive drug administration	3	B	
Transurethral resection (coagulation and LASER)	NA	A/B	Hunner's lesions only
Nerve blockade/epidural pain pumps	3	C	For crisis intervention; affects pain only
Sacral neuromodulation	3	B	Not recommended beyond clinical trials
Bladder training	3	B	Patients with little pain
Manual and physical therapy		3	B
Diet	3	C	
Acupuncture	3	C	Data contradictory
Hypnosis		No data	
Psychological therapy	3	B	
Surgical treatment	NA	A	Largely varying data ultima ratio, experienced surgeons

PPS = pentosanpolysulphate sodium; DMSO = dimethyl sulphoxide; NA = type of evidence not applicable, since RCTs are unethical in such surgical procedures.

Figure 3: Flowchart for the diagnosis and therapy of bladder pain syndrome/interstitial cystitis



2.7.15 References

1. Skene AJC. Diseases of the bladder and urethra in women. New York: William Wood 1887;167.
2. Hunner GL. A rare type of bladder ulcer in women: report of cases. Boston Med Surg J 1915;172:660-664.
3. Hunner G. Elusive ulcer of the bladder: further notes on a rare type of bladder ulcer with report of 25 cases. Am J Obstet 1918;78:374-395.
4. Hand JR. Interstitial cystitis: report of 223 cases (204 women and 19 men). J Urol 1949;61:291.
5. Messing EM, Stamey TA. Interstitial cystitis: early diagnosis, pathology, and treatment. Urology 1978;12(4):381-392.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=213864&dopt=Abstract
6. Fall M, Johansson SL, Aldenborg F. Chronic interstitial cystitis: a heterogeneous syndrome. J Urol 1987;137(1):35-38.
7. Abrams P, Cardozo L, Fall M, Griffiths D, Rosier P, Ulmsten U, van Kerrebroeck P, Victor A, Wein A. The standardisation of terminology of lower urinary tract function: report from the Standardisation Subcommittee of the International Continence Society. Am J Obstet Gynecol 2002;187(1):116-126.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=12114899&dopt=Abstract
8. van de Merwe JP, Nordling J, Bouchelouche P, Bouchelouche K, Cervigni M, Daha LK, Elneil S, Fall M, Hohlbrugger G, Irwin P, Mortensen S, van Ophoven A, Osborne JL, Pecker R, Richter B, Riedl C, Sairanen J, Tinzi M, Wyndaele JJ. Diagnostic criteria, classification, and nomenclature for painful bladder syndrome/interstitial cystitis: an ESSIC proposal. Eur Urol 2008;53(1):60-67.
http://www.ncbi.nlm.nih.gov/pubmed/17900797?ordinalpos=17&itool=EntrezSystem2.PEntrez.Pubmed_d.Pubmed_ResultsPanel.Pubmed_RVDocSum
9. Gillenwater JY, Wein AJ. Summary of the National Institute of Arthritis, Diabetes, Digestive and Kidney Diseases Workshop on Interstitial Cystitis, National Institutes of Health, Bethesda, Maryland, August 28-29, 1987. J Urol 1988;140(1):203-206.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3379688&dopt=Abstract

10. Hanno PM, Landis JR, Matthews-Cook Y, Kusek J, Nyberg L Jr. The diagnosis of interstitial cystitis revisited: lessons learned from the National Institutes of Health Interstitial Cystitis Database study. *J Urol* 1999;161(2):553-557.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9915447&dopt=Abstract
11. Erickson DR, Belchis DA, Dabbs DJ. Inflammatory cell types and clinical features of interstitial cystitis. *J Urol* 1997;158(3 Pt 1):790-793.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9258082&dopt=Abstract
12. Peeker R, Fall M. Toward a precise definition of interstitial cystitis: further evidence of differences in classic and nonulcer disease. *J Urol* 2002;167(6):2470-2472.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11992059&dopt=Abstract
13. Nordling J, Anjum FH, Bade JJ, Bouchelouche K, Bouchelouche P, Cervigni M, Elneil S, Fall M, Hald T, Hanus T, Hedlund H, Hohlbrugger G, Horn T, Larsen S, Leppilahti M, Mortensen S, Nagendra M, Oliveira PD, Osborne J, Riedl C, Sairanen J, Tinzl M, Wyndaele JJ. Primary evaluation of patients suspected of having interstitial cystitis (IC). *Eur Urol* 2004;45(5):662-669.
http://www.ncbi.nlm.nih.gov/pubmed/15082211?ordinalpos=20&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
14. Domingue GJ, Ghoniem GM, Bost KL, Fermin C, Human LG. Dormant microbes in interstitial cystitis. Erratum in: *J Urol* 1996;155:298. *J Urol* 1995;153(4):1321-1326.
http://www.ncbi.nlm.nih.gov/80/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7869536&dopt=Abstract
15. Lynes WL, Sellers RG, Dairiki Shortliffe LM. The evidence for occult bacterial infections as a cause for interstitial cystitis. *J Urol* 1989;141:268A (Abstract 393).
16. Hukkanen V, Haarala M, Nurmi M, Kleini P, Kiilholma P. Viruses and interstitial cystitis: adenovirus genomes cannot be demonstrated in urinary bladder biopsies. *Urol Res* 1996;24(4):235-238.
http://www.ncbi.nlm.nih.gov/pubmed/8873382?ordinalpos=11&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
17. Fall M, Johansson SL, Vahlne A. A clinicopathological and virological study of interstitial cystitis. *J Urol* 1985;133(5):771-773.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2985831&dopt=Abstract
18. Duncan JL, Schaeffer AJ. Do infectious agents cause interstitial cystitis? *Urology* 1997;49(5A Suppl):48-51.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9146001&dopt=Abstract
19. Al-Hadithi HN, Williams H, Hart CA, Frazer M, Adams EJ, Richmond DH, Tincello DG. Absence of bacterial and viral DNA in bladder biopsies from patients with interstitial cystitis/chronic pelvic pain syndrome. *J Urol* 2005;174(1):151-154.
http://www.ncbi.nlm.nih.gov/pubmed/15947607?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
20. Agarwal M, Dixon RA. A study to detect *Helicobacter pylori* in fresh and archival specimens from patients with interstitial cystitis, using amplification methods. *BJU Int* 2003;91(9):814-816.
http://www.ncbi.nlm.nih.gov/pubmed/12780839?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
21. Peeker R, Enerback L, Fall M, Aldenborg F. Recruitment, distribution and phenotypes of mast cells in interstitial cystitis. *J Urol* 2000;163(3):1009-1015.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10688040&dopt=Abstract
22. Dundore PA, Schwartz AM, Semerjian H. Mast cell counts are not useful in the diagnosis of nonulcerative interstitial cystitis. *J Urol* 1996;155(3):885-887.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8583599&dopt=Abstract
23. Johansson SL, Fall M. Clinical features and spectrum of light microscopic changes in interstitial cystitis. *J Urol* 1990;143(6):1118-1124.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2342171&dopt=Abstract

24. Anderström CR, Fall M, Johansson SL. Scanning electron microscopic findings in interstitial cystitis. *Br J Urol* 1989;63(3):270-275.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2702424&dopt=Abstract
25. Fellows GJ, Marshall DH. The permeability of human bladder epithelium to water and sodium. *Invest Urol* 1972;9(4):339-944.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5058772&dopt=Abstract
26. Parsons CL, Mulholland SG. Successful therapy of interstitial cystitis with pentosanpolysulfate. *J Urol* 1987;138(3):513-516.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2442417&dopt=Abstract
27. Parsons CL, Lilly JD, Stein P. Epithelial dysfunction in nonbacterial cystitis (interstitial cystitis). *J Urol* 1991;145(4):732-735.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2005689&dopt=Abstract
28. Lokeshwar VB, Selzer MG, Cerwinka WH, Gomez MF, Kester RR, Bejany DE, Gousse AE. Urinary uronate and sulfated glycosaminoglycan levels: markers for interstitial cystitis severity. *J Urol* 2005;174(1):344-349.
http://www.ncbi.nlm.nih.gov/pubmed/15947687?ordinalpos=7&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
29. Oravisto KJ, Alfthan OS, Jokinen EJ. Interstitial cystitis. Clinical and immunological findings. *Scand J Urol Nephrol* 1970;4(1):37-42.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5314306&dopt=Abstract
30. Jokinen EJ, Alfthan OS, Oravisto KJ. Antitissue antibodies in interstitial cystitis. *Clin Exp Immunol* 1972;11(3):333-339.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=4114472&dopt=Abstract
31. Ochs RL, Stein TW Jr, Peebles CL, Gittes RF, Tan EM. Autoantibodies in interstitial cystitis. *J Urol* 1994;151(3):587-592.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8308964&dopt=Abstract
32. Tan EM. Antinuclear antibodies: diagnostic markers for autoimmune diseases and probes for cell biology. *Adv Immunol* 1989;44:93-151.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2646863&dopt=Abstract
33. von Muhlen CA, Tan EM. Autoantibodies in the diagnosis of systemic rheumatic diseases. *Semin Arthritis Rheum* 1995;24(5):323-358.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7604300&dopt=Abstract
34. Ochs RL. Autoantibodies and interstitial cystitis. *Clin Lab Med* 1997;17(3):571-579.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9316774&dopt=Abstract
35. Mattila J. Vascular immunopathology in interstitial cystitis. *Clin Immunol Immunopathol* 1982;23(3):648-655.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6981479&dopt=Abstract
36. Mattila J, Linder E. Immunoglobulin deposits in bladder epithelium and vessels in interstitial cystitis: possible relationship to circulating anti-intermediate filament autoantibodies. *Clin Immunol Immunopathol* 1984;32(1):81-89.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6733983&dopt=Abstract
37. Harrington DS, Fall M, Johansson SL. Interstitial cystitis: bladder mucosa lymphocyte immunophenotyping and peripheral blood flow cytometry analysis. *J Urol* 1990;144(4):868-871.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2204728&dopt=Abstract
38. van de Merwe JP. Sjögren's syndrome in patients with interstitial cystitis. Preliminary results in 100 patients. *Int J Urol* 2003; 10 (Suppl):S69.

39. Ehren I, Hosseini A, Lundberg JO, Wiklund NP. Nitric oxide: a useful gas in the detection of lower urinary tract inflammation. *J Urol* 1999;162(2):327-329.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10411031&dopt=Abstract
40. Moncada S, Higgs A. The L-arginine-nitric oxide pathway. *N Engl J Med* 1993;329(27):2002-2012.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7504210&dopt=Abstract
41. Smith SD, Wheeler MA, Foster HE Jr, Weiss RM. Urinary nitric oxide synthase activity and cyclic GMP levels are decreased with interstitial cystitis and increased with urinary tract infections. *J Urol* 1996;155(4):1432-1435.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8632605&dopt=Abstract
42. Logadottir YR, Ehren I, Fall M, Wiklund NP, Peeker R, Hanno PM. Intravesical nitric oxide production discriminates between classic and nonulcer interstitial cystitis. *J Urol* 2004;171(3):1148-50; discussion 50-1.
<http://www.ncbi.nlm.nih.gov/pubmed/14767289?ordinalpos=11&itool=EntrezSystem2.PEntrez.Pubmed.ResultsPanel.Pubmed.RVDocSum>
43. Birder LA, Wolf-Johnston A, Buffington CA, Roppolo JR, de Groat WC, Kanai AJ. Altered inducible nitric oxide synthase expression and nitric oxide production in the bladder of cats with feline interstitial cystitis. *J Urol* 2005;173(2):625-629.
<http://www.ncbi.nlm.nih.gov/pubmed/15643277?ordinalpos=10&itool=EntrezSystem2.PEntrez.Pubmed.ResultsPanel.Pubmed.RVDocSum>
44. Sugimura K, Haimoto H, Nagura H, Kato K, Takahashi A. Immunohistochemical differential distribution of S-100 alpha and S-100 beta in the peripheral nervous system of the rat. *Muscle Nerve* 1989;12(11):929-935.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2608087&dopt=Abstract
45. Peeker R, Aldenborg F, Haglid K, Johansson SL, Rosengren L, Fall M. Decreased levels of S-100 protein in non-ulcer interstitial cystitis. *Scand J Urol Nephrol* 1998;32(6):395-398.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9925003&dopt=Abstract
46. Hohenfellner M, Nunes L, Schmidt RA, Lampel A, Thuroff JW, Tanagho EA. Interstitial cystitis: increased sympathetic innervation and related neuropeptide synthesis. *J Urol* 1992;147(3):587-591.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1538434&dopt=Abstract
47. Twiss CO, Kilpatrick L, Triaca V, Arboleda V, Craske M, Ibrahimovic H, Raz S, Mayer EA, Ornitz E, Rodriguez LV, Naliboff BD. Evidence for central hyperexcitability in patients with interstitial cystitis. *J Urol* 2007;177(4):49.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10737477&dopt=Abstract
48. Yilmaz U, Liu YW, Berger RE, Yang CC. Autonomic nervous system changes in men with chronic pelvic pain syndrome. *J Urol* 2007;177(6):2170-2174.
<http://www.ncbi.nlm.nih.gov/pubmed/17509311?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.ResultsPanel.Pubmed.RVDocSum>
49. Twiss CO, Kilpatrick L, Triaca V, Arboleda V, Craske M, Ibrahimovic H, Raz S, Mayer EA, Ornitz E, Rodriguez LV, Naliboff BD. Evidence for central hyperexcitability in patients with interstitial cystitis. *J Urol* 2007;177(4):49.
50. Elbadawi AE, Light JK. Distinctive ultrastructural pathology of nonulcerative interstitial cystitis: new observations and their potential significance in pathogenesis. *Urol Int* 1996;56(3):137-162.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Search&itool=pubmed_AbstractPlus&term=%22Elbadawi+AE%22%5BAuthor%5D
51. Parsons CL, Bautista SL, Stein PC, Zupkas P. Cyto-injury factors in urine: a possible mechanism for the development of interstitial cystitis. *J Urol* 2000;164(4):1381-1384.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10992419&dopt=Abstract
52. Hang L, Wullt B, Shen Z, Karpman D, Svanborg C. Cytokine repertoire of epithelial cells lining the human urinary tract. *J Urol* 1998;159(6):2185-2192.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9598567&dopt=Abstract

53. Parsons CL, Stein P, Zupkas P, Chenoweth M, Argade SP, Proctor JG, Datta A, Trotter RN. Defective Tamm-Horsfall protein in patients with interstitial cystitis. *J Urol* 2007;178(6):2665-2670.
http://www.ncbi.nlm.nih.gov/pubmed/17945284?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
54. Rosamilia A, Cann L, Dwyer P, Scurry J, Rogers P. Bladder microvasculature in women with interstitial cystitis. *J Urol* 1999;161(6):1865-1870.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10332455&dopt=Abstract
55. Pontari MA, Hanno PM, Ruggieri MR. Comparison of bladder blood flow in patients with and without interstitial cystitis. *J Urol* 1999;162(2):330-334.
56. van Ophoven A, Rossbach G, Pajonk F, Hertle L. Safety and efficacy of hyperbaric oxygen therapy for the treatment of interstitial cystitis: a randomized, sham controlled, double-blind trial. *J Urol* 2006;176(4 Pt 1):1442-1446
http://www.ncbi.nlm.nih.gov/pubmed/16952654?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
57. Theoharides TC, Pang X, Letourneau R, Sant GR. Interstitial cystitis: a neuroimmunoendocrine disorder. *Ann N Y Acad Sci* 1998;840:619-634.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9629289&dopt=Abstract
58. Okragly AJ, Niles AL, Saban R, Schmidt D, Hoffman RL, Warner TF, Moon TD, Uehling DT, Haak-Frendscho M. Elevated tryptase, nerve growth factor, neurotrophin-3 and glial cell line-derived neurotrophic factor levels in the urine of interstitial cystitis and bladder cancer patients. *J Urol* 1999;161(2):438-441; discussion 441-442.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9915421&dopt=Abstract
59. Abdel-Mageed AB, Ghoniem GM. Potential role of rel/nuclear factor-kappaB in the pathogenesis of interstitial cystitis. *J Urol* 1998;160(6 Pt 1):2000-2003.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9817309&dopt=Abstract
60. Abdel-Mageed AB, Bajwa A, Shenassa BB, Human L, Ghoniem GM. NF-kappaB-dependent gene expression of proinflammatory cytokines in T24 cells: possible role in interstitial cystitis. *Urol Res* 2003;31(5):300-305.
http://www.ncbi.nlm.nih.gov/pubmed/14574533?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
61. Abdel-Mageed A, Ghoniem G, Human I, Agrawal KD. Induction of proinflammatory cytokine gene expression by NF-kappaB in human bladder epithelial (T-24) cells: possible mechanism for interstitial cystitis. *J Urol* 1999;161(Suppl):28.
62. Oravisto KJ. Epidemiology of interstitial cystitis. *Ann Chir Gynaecol Fenn* 1975;64(2):75-77.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1137336&dopt=Abstract
63. Held PJ, Hanno PM, Wein AJ. In: Hanno PM, Staskin DR, Krane RJ, Wein AJ eds. *Interstitial Cystitis. Epidemiology of interstitial cystitis*. London: Springer Verlag, 1990, pp. 29-48.
64. Bade JJ, Rijcken B, Mensink HJ. Interstitial cystitis in The Netherlands: prevalence, diagnostic criteria and therapeutic preferences. *J Urol* 1995;154(6):2035-2037; discussion 2037-2038.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7500452&dopt=Abstract
65. Jones CA, Harris MA, Nyberg L. Prevalence of interstitial cystitis in the United States, *Proc Am Urol Ass J Urol* 1994;151(Suppl):423A.
66. Curhan GC, Speizer FE, Hunter DJ, Curhan SG, Stampfer MJ. Epidemiology of interstitial cystitis: a population based study. *J Urol* 1999;161(2):549-552.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9915446&dopt=Abstract
67. Leppilahti M, Sairanen J, Tammela TL, Aaltomaa S, Lehtoranta K, Auvinen A; Finnish Interstitial Cystitis-Pelvic Pain Syndrome Study Group. Prevalence of clinically confirmed interstitial cystitis in women: a population based study in Finland. *J Urol* 2005;174(2):581-583.
http://www.ncbi.nlm.nih.gov/pubmed/16006902?ordinalpos=5&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

68. Temml C, Wehrberger C, Riedl C, Ponholzer A, Marszalek M, Madersbacher S. Prevalence and correlates for interstitial cystitis symptoms in women participating in a health screening project. *Eur Urol* 2007;51(3):803-808; discussion 809.
http://www.ncbi.nlm.nih.gov/pubmed/16979286?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
69. Burkman RT. Chronic pelvic pain of bladder origin: epidemiology, pathogenesis and quality of life. *J Reprod Med* 2004;49(3 Suppl):225-229.
http://www.ncbi.nlm.nih.gov/pubmed/15088860?ordinalpos=6&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
70. Roberts RO, Bergstralh EJ, Bass SE, Lightner DJ, Lieber MM, Jacobsen SJ. Incidence of physician-diagnosed interstitial cystitis in Olmsted County: a community-based study. *BJU Int* 2003;91(3):181-185.
http://www.ncbi.nlm.nih.gov/pubmed/12581000?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
71. Greenberg E, Barnes R, Stewart S, Furnish T. Transurethral resection of Hunner's ulcer. *J Urol* 1974;111(6):764-766.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=4830879&dopt=Abstract
72. Koziol JA. Epidemiology of interstitial cystitis. *Urol Clin North Am* 1994;21(1):7-20.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8284848&dopt=Abstract
74. Parsons CL. Interstitial cystitis: clinical manifestations and diagnostic criteria in over 200 cases. *Neurourol Urodynam* 1990;9:241.
75. Koziol JA, Adams HP, Frutos A. Discrimination between the ulcerous and the nonulcerous forms of interstitial cystitis by noninvasive findings. *J Urol* 1996;155(1):87-90.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7490906&dopt=Abstract
76. Parsons CL, Zupkas P, Parsons JK. Intravesical potassium sensitivity in patients with interstitial cystitis and urethral syndrome. *Urology* 2001;57(3):428-432.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11248610&dopt=Abstract
77. Warren J, Jackson T, Meyers D, Xu J. Fishbein/interstitial cystitis association (ICA) survey of interstitial cystitis among family members of ICA members: preliminary analysis. *Urology* 2001;57(6 Suppl 1):126-127.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11378121&dopt=Abstract
78. Warren JW, Keay SK, Meyers D, Xu J. Concordance of interstitial cystitis in monozygotic and dizygotic twin pairs. *Urology* 2001;57(6 Suppl 1):22-25.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11378045&dopt=Abstract
79. Alagiri M, Chottiner S, Ratner V, Slade D, Hanno PM. Interstitial cystitis: unexplained associations with other chronic disease and pain syndromes. *Urology* 1997;49(5A Suppl):52-57.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9146002&dopt=Abstract
80. Clauw DJ, Schmidt M, Radulovic D, Singer A, Katz P, Bresette J. The relationship between fibromyalgia and interstitial cystitis. *J Psychiatr Res* 1997;31(1):125-131.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9201654&dopt=Abstract
81. Erickson DR, Morgan KC, Ordille S, Keay SK, Xie SX. Nonbladder related symptoms in patients with interstitial cystitis. *J Urol* 2001;166(2):557-561; discussion 561-562.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11458068&dopt=Abstract
82. Myrna M. Weissman, PhD; Raz Gross, MD, MPH; Abby Fyer, MD; Gary A. Heiman, PhD; Marc J. Gameroff, PhD; Susan E. Hodge, DSc; David Kaufman, MD; Steven A. Kaplan, MD; Priya J. Wickramaratne, PhD. Interstitial Cystitis and Panic Disorder - A Potential Genetic Syndrome. *Arch Gen Psych* 2004;61:273-279.
<http://archpsyc.ama-assn.org/cgi/content/abstract/61/3/273>

83. Buffington CA. Comorbidity of interstitial cystitis with other unexplained clinical conditions. *J Urol* 2004;172(4 Pt 1):1242-1248.
http://www.ncbi.nlm.nih.gov/pubmed/15371816?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
84. Bullock AD, Becich MJ, Klutke CG, Ratliff TL. Experimental autoimmune cystitis: a potential murine model for ulcerative interstitial cystitis. *J Urol* 1992;148(6):1951-1956.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1433651&dopt=Abstract
85. Dodd LG, Tello J. Cytologic examination of urine from patients with interstitial cystitis. *Acta Cytol* 1998;42(4):923-927.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9684578&dopt=Abstract
86. Erickson DR, Davies MF. Interstitial cystitis. *Int Urogynecol J Pelvic Floor Dysfunct* 1998;9(3):174-183.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9745978&dopt=Abstract
87. Hanno PM. Amitriptyline in the treatment of interstitial cystitis. *Urol Clin North Am* 1994;21(1):89-91.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8284851&dopt=Abstract
88. Fall M, Lindstrom S. Transcutaneous electrical nerve stimulation in classic and nonulcer interstitial cystitis. *Urol Clin North Am* 1994;21(1):131-139.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?db=pubmed&cmd=Retrieve&dopt=AbstractPlus&list_uids=8284836&itool=iconabstr&query_hl=129&itool=pubmed_docsum
89. Peeker R, Haghsheno MA, Holmang S, Fall M. Intravesical bacillus Calmette-Guerin and dimethyl sulfoxide for treatment of classic and nonulcer interstitial cystitis: a prospective, randomized double-blind study. *J Urol* 2000;164(6):1912-1915; discussion 1915-1916.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11061879&dopt=Abstract
90. Fritjofsson A, Fall M, Juhlin R, Persson BE, Ruutu M. Treatment of ulcer and nonulcer interstitial cystitis with sodium pentosanpolysulfate: a multicenter trial. *J Urol* 1987;138(3):508-512.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2442416&dopt=Abstract
91. Kozioł JA, Clark DC, Gittes RF, Tan EM. The natural history of interstitial cystitis: a survey of 374 patients. *J Urol* 1993;149(3):465-469.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8437248&dopt=Abstract
92. Enerback L, Fall M, Aldenborg F. Histamine and mucosal mast cells in interstitial cystitis. *Agents & Actions* 1989;27(1-2):113-116.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2750582&dopt=Abstract
93. Lechevallier E. Interstitial cystitis. *Prog Urol* 1995;5(1):21-30.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7719356&dopt=Abstract
94. Messing E, Pauk D, Schaeffer A, Nieweglowski M, Nyberg LM Jr, Landis JR, Cook YL, Simon LJ. Associations among cystoscopic findings and symptoms and physical examination findings in women enrolled in the Interstitial Cystitis Data Base (ICDB) Study. *Urology* 1997;49(5A Suppl):8185.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9146006&dopt=Abstract
95. Waxman JA, Sulak PJ, Kuehl TJ. Cystoscopic findings consistent with interstitial cystitis in normal women undergoing tubal ligation. *J Urol* 1998;160(5):1663-1667.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9783927&dopt=Abstract
96. Shear S, Mayer R. Development of glomerulations in younger women with interstitial cystitis. *Urology* 2006;68(2):253-256.
http://www.ncbi.nlm.nih.gov/pubmed/16904429?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
97. Ottem DP, Teichman JM. What is the value of cystoscopy with hydrodistension for interstitial cystitis? *Urology* 2005;66(3):494-499.
http://www.ncbi.nlm.nih.gov/pubmed/16140064?ordinalpos=15&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

98. Cole EE, Scarpero HM, Dmochowski RR. Are patient symptoms predictive of the diagnostic and/or therapeutic value of hydrodistention? *Neurourol Urodyn* 2005;24(7):638-42.
http://www.ncbi.nlm.nih.gov/pubmed/16208660?ordinalpos=34&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
99. Lamale LM, Lutgendorf SK, Hoffman AN, Kreder KJ. Symptoms and cystoscopic findings in patients with untreated interstitial cystitis. *Urology* 2006;67(2):242-245.
http://www.ncbi.nlm.nih.gov/pubmed/16442603?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
100. Tamaki M, Saito R, Ogawa O, Yoshimura N, Ueda T. Possible mechanisms inducing glomerulations in interstitial cystitis: relationship between endoscopic findings and expression of angiogenic growth factors. *J Urol* 2004;172(3):945-948.
http://www.ncbi.nlm.nih.gov/pubmed/15311005?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
101. Johansson SL, Fall M. Pathology of interstitial cystitis. *Urol Clin North Am* 1994;21(1):55-62.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8284845&dopt=Abstract
102. Parsons CL, Greenberger M, Gabal L, Bidair M, Barme G. The role of urinary potassium in the pathogenesis and diagnosis of interstitial cystitis. *J Urol* 1998;159(6):1862-6;discussion 1866-1867.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9598476&dopt=Abstract
103. Chambers GK, Fenster HN, Cripps S, Jens M, Taylor D. An assessment of the use of intravesical potassium in the diagnosis of interstitial cystitis. *J Urol* 1999;162(3 Pt 1):699-701.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10458346&dopt=Abstract
104. Gregoire M, Liandier F, Naud A, Lacombe L, Fradet Y. Does the potassium stimulation test predict cystometric, cystoscopic outcome in interstitial cystitis? *J Urol* 2002;168(2):556-557.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=12131308&dopt=Abstract
105. Daha LK, Riedl CR, Hohlbrugger G, Knoll M, Engelhardt PF, Pflüger H. Comparative assessment of maximal bladder capacity, 0.9% NaCl versus 0.2 M KCl, for the diagnosis of interstitial cystitis: a prospective controlled study. *J Urol* 2003;170(3):807-809.
http://www.ncbi.nlm.nih.gov/pubmed/12913704?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
106. Gupta SK, Pidcock L, Parr NJ. The potassium sensitivity test: a predictor of treatment response in interstitial cystitis. *BJU Int* 2005;96(7):1063-1066.
http://www.ncbi.nlm.nih.gov/pubmed/16225529?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
107. Lubeck DP, Whitmore K, Sant GR, Alvarez-Horine S, Lai C. Psychometric validation of the O'leary-Sant interstitial cystitis symptom index in a clinical trial of pentosan polysulfate sodium. *Urology* 2001; 57(6 Suppl 1):62-66.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11378052&dopt=Abstract&itool=iconabstr
108. Keay S, Kleinberg M, Zhang CO, Hise MK, Warren JW. Bladder epithelial cells from patients with interstitial cystitis produce an inhibitor of heparin-binding epidermal growth factor-like growth factor production. *J Urol* 2000;164(6):2112-2118.
http://www.ncbi.nlm.nih.gov/pubmed/11061938?ordinalpos=13&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
109. Close CE, Carr MC, Burns MW, Miller JL, Bavendam TG, Mayo ME, Mitchell ME. Interstitial cystitis in children. *J Urol* 1996;156(2 Pt 2):860-862.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8683802&dopt=Abstract
110. Mattox TF. Interstitial cystitis in adolescents and children: a review. *J Pediatr Adolesc Gynecol* 2004;17(1):7-11.
http://www.ncbi.nlm.nih.gov/pubmed/15010032?ordinalpos=194&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
111. Novicki DE, Larson TR, Swanson SK. Interstitial cystitis in men. *Urology* 1998;52(4):621-624.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9763081&dopt=Abstract

112. Miller JL, Rothman I, Bavendam TG, Berger RE. Prostatodynia and interstitial cystitis: one and the same? *Urology* 1995;45(4):587-90.
http://www.ncbi.nlm.nih.gov/pubmed/7716839?ordinalpos=6&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
113. Pontari MA. Chronic prostatitis/chronic pelvic pain syndrome and interstitial cystitis: are they related? *Curr Urol Rep* 2006;7(4):329-334.
http://www.ncbi.nlm.nih.gov/pubmed/16930505?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
114. Keay S, Zhang CO, Chai T, Warren J, Koch K, Grkovic D, Colville H, Alexander R. Antiproliferative factor, heparin-binding epidermal growth factor-like growth factor, and epidermal growth factor in men with interstitial cystitis versus chronic pelvic pain syndrome. *Urology* 2004;63(1):22-26.
http://www.ncbi.nlm.nih.gov/pubmed/14751340?ordinalpos=48&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
115. Nickel JC. Opioids for chronic prostatitis and interstitial cystitis: lessons learned from the 11th World Congress on Pain. *Urology* 2006;68(4):697-701.
http://www.ncbi.nlm.nih.gov/pubmed/17070334?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
116. Badenoch AW. Chronic interstitial cystitis. *Br J Urol* 1971;43(6):718-721.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5159574&dopt=Abstract
117. Pool TL. Interstitial cystitis: clinical considerations and treatment. *Clin Obstet Gynecol* 1967;10(1):185-191.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6021011&dopt=Abstract
118. Soucy F, Gregoire M. Efficacy of prednisone for severe refractory ulcerative interstitial cystitis. *J Urol* 2005;173(3):841-3; discussion 843.
http://www.ncbi.nlm.nih.gov/pubmed/15711286?ordinalpos=10&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
119. Theoharides TC. Hydroxyzine in the treatment of interstitial cystitis. *Urol Clin North Am* 1994;21(1):113-119.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8284834&dopt=Abstract
120. Seshadri P, Emerson L, Morales A. Cimetidine in the treatment of interstitial cystitis. *Urology* 1994;44(4):614-616.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7941209&dopt=Abstract
121. Theoharides TC. Hydroxyzine for interstitial cystitis. *J Allergy Clin Immunol* 1993;91(2):686-687.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8436783&dopt=Abstract
122. Theoharides TC, Sant GR. Hydroxyzine therapy for interstitial cystitis. *Urology* 1997;49(5A Suppl):108-110.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9146011&dopt=Abstract
123. Sant GR, Probert KJ, Hanno PM, Burks D, Culkin D, Diokno AC, Hardy C, Landis JR, Mayer R, Madigan R, Messing EM, Peters K, Theoharides TC, Warren J, Wein AJ, Steers W, Kusek JW, Nyberg LM Interstitial Cystitis Clinical Trials Group. A pilot clinical trial of oral pentosan polysulfate and oral hydroxyzine in patients with interstitial cystitis. *J Urol* 2003;170(3):810-815.
http://www.ncbi.nlm.nih.gov/pubmed/12913705?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
124. Baldessarini RJ. Drugs and the treatment of psychiatric disorders. In: Gilman A, Gooman L, Rall T, eds. *The Pharmacological Basis of Therapeutics*. 7th ed. New York: Macmillan, 1985, pp. 387-445.
125. Hanno PM, Buehler J, Wein AJ. Use of amitriptyline in the treatment of interstitial cystitis. *J Urol* 1989;141(4):846-848.
http://www.ncbi.nlm.nih.gov/pubmed/2926877?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
126. Kirkemo AK, Miles BJ, Peters JM. Use of amitriptyline in interstitial cystitis. *J Urol* 1990;143(Suppl):279A.

127. van Ophoven A, Pokupic S, Heinecke A, Hertle L. A prospective, randomized, placebo controlled, double-blind study of amitriptyline for the treatment of interstitial cystitis. *J Urol* 2004;172(2):533-536.
http://www.ncbi.nlm.nih.gov/pubmed/15247722?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
128. van Ophoven A, Hertle L. Long-term results of amitriptyline treatment for interstitial cystitis. *J Urol* 2005;174(5):1837-1840.
http://www.ncbi.nlm.nih.gov/pubmed/16217303?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
129. Mulholland SG, Hanno P, Parsons CL, Sant GR, Staskin DR. Pentosan polysulfate sodium for therapy of interstitial cystitis. A double-blind placebo-controlled clinical study. *Urology* 1990;35(6):552-558.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1693797&dopt=Abstract
130. Hwang P, Auclair B, Beechinor D, Diment M, Einarson TR. Efficacy of pentosan polysulfate in the treatment of interstitial cystitis: a meta-analysis. *Urology* 1997;50(1):39-43.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9218016&dopt=Abstract
131. Nickel JC, Barkin J, Forrest J, Mosbaugh PG, Hernandez-Graulau J, Kaufman D, Lloyd K, Evans RJ, Parsons CL, Atkinson LE; Elmiron Study Group. Randomized, double-blind, dose-ranging study of pentosan polysulfate sodium for interstitial cystitis. *Urology* 2005;65(4):654-658.
http://www.ncbi.nlm.nih.gov/pubmed/15833501?ordinalpos=16&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
132. van Ophoven A, Heinecke A, Hertle L. Safety and efficacy of concurrent application of oral pentosan polysulfate and subcutaneous low-dose heparin for patients with interstitial cystitis. *Urology* 2005;66(4):707-711.
http://www.ncbi.nlm.nih.gov/pubmed/16230121?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
133. Warren JW, Horne LM, Hebel JR, Marvel RP, Keay SK, Chai TC. Pilot study of sequential oral antibiotics for the treatment of interstitial cystitis. *J Urol* 2000;163(6):1685-1688.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10799160&dopt=Abstract
134. Oravisto KJ, Alfthan OS. Treatment of interstitial cystitis with immunosuppression and chloroquine derivatives. *Eur Urol* 1976;2(2):82-84.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=971677&dopt=Abstract
135. Forsell T, Ruutu M, Isoniemi H, Ahonen J, Alfthan O. Cyclosporine in severe interstitial cystitis. *J Urol* 1996;155(5):1591-1593.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8627830&dopt=Abstract
136. Moran PA, Dwyer PL, Carey MP, Maher CF, Radford NJ. Oral methotrexate in the management of refractory interstitial cystitis. *Aust NZ J Obstet Gynaecol* 1999;39(4):468-471.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10687766&dopt=Abstract
137. Sairanen J, Forsell T, Ruutu M. Long-term outcome of patients with interstitial cystitis treated with low dose cyclosporine A. *J Urol* 2004;171(6 Pt 1):2138-2141.
http://www.ncbi.nlm.nih.gov/pubmed/15126772?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
138. Sairanen J, Tammela TL, Leppilahti M, Multanen M, Paananen I, Lehtoranta K, Ruutu M. Cyclosporine A and pentosan polysulfate sodium for the treatment of interstitial cystitis: a randomized comparative study. *J Urol* 2005;174(6):2235-2238.
http://www.ncbi.nlm.nih.gov/pubmed/16280777?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
139. Hansen HC. Interstitial cystitis and the potential role of gabapentin. *South Med J* 2000;93(2):238-242.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10701800&dopt=Abstract
140. Sasaki K, Smith CP, Chuang YC, Lee JY, Kim JC, Chancellor MB. Oral gabapentin (neurontin) treatment of refractory genitourinary tract pain. *Tech Urol* 2001;7(1):47-49.
http://www.ncbi.nlm.nih.gov/pubmed/11272678?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

141. Sonnett TE, Setter SM, Campbell RK. Pregabalin for the treatment of painful neuropathy. *Expert Rev Neurother* 2006;6(11):1629-35.
http://www.ncbi.nlm.nih.gov/pubmed/17144773?ordinalpos=5&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
142. Ueda T, Tamaki M, Ogawa O, Yamauchi T, Yoshimura N. Improvement of interstitial cystitis symptoms and problems that developed during treatment with oral IPD-1151T. *J Urol* 2000;164(6):1917-1920.
http://www.ncbi.nlm.nih.gov/pubmed/11061880?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
143. Katske F, Shoskes DA, Sender M, Poliakin R, Gagliano K, Rajfer J. Treatment of interstitial cystitis with a quercetin supplement. *Tech Urol* 2001;7(1):44-46.
http://www.ncbi.nlm.nih.gov/80/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11272677&dopt=Abstract
144. Theoharides TC, Sant GR. A pilot open label study of Cystoprotek in interstitial cystitis. *Int J Immunopathol Pharmacol* 2005 Jan-Mar;18(1):183-188.
http://www.ncbi.nlm.nih.gov/pubmed/15698523?ordinalpos=8&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
145. Dimitrakov J, Tchitalov J, Zlatanov T, Dikov D, Rawadi G. Recombinant human nerve growth factor in the treatment of interstitial cystitis: preliminary results. *Urology* 2001;57(6 Suppl 1):118-119.
http://www.ncbi.nlm.nih.gov/pubmed/11378099?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
146. Giannakopoulos X, Champilomatos P. Chronic interstitial cystitis. Successful treatment with intravesical lidocaine. *Arch Ital Urol Nefrol Androl* 1992;64(4):337-339.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1462157&dopt=Abstract
147. Asklin B, Cassuto J. Intravesical lidocaine in severe interstitial cystitis. Case report. *Scand J Urol Nephrol* 1989;23(4):311-312.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2595329&dopt=Abstract
148. Henry R, Patterson L, Avery N, Tanzola R, Tod D, Hunter D, Nickel JC, Morales A. Absorption of alkalized intravesical lidocaine in normal and inflamed bladders: a simple method for improving bladder anaesthesia. *J Urol* 2001;165(6 Pt 1):1900-1903.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11371877&dopt=Abstract
149. Parsons CL. Successful downregulation of bladder sensory nerves with combination of heparin and alkalized lidocaine in patients with interstitial cystitis. *Urology* 2005;65(1):45-48.
http://www.ncbi.nlm.nih.gov/pubmed/15667861?ordinalpos=6&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
150. Bade JJ, Laseur M, Nieuwenburg A, van der Weele LT, Mensink HJ. A placebo-controlled study of intravesical pentosanpolysulphate for the treatment of interstitial cystitis. *Br J Urol* 1997;79(2):168-171.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9052464&dopt=Abstract
151. Parsons CL, Housley T, Schmidt JD, Lebow D. Treatment of interstitial cystitis with intravesical heparin. *Br J Urol* 1994;73(5):504-507.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8012771&dopt=Abstract
152. Kuo HC. Urodynamic results of intravesical heparin therapy for women with frequency urgency syndrome and interstitial cystitis. *J Formos Med Assoc* 2001;100(5):309-314.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11432309&dopt=Abstract&itool=iconabstr
153. Baykal K, Senkul T, Sen B, Karademir K, Adayener C, Erden D. Intravesical heparin and peripheral neuromodulation on interstitial cystitis. *Urol Int* 2005;74(4):361-364.
http://www.ncbi.nlm.nih.gov/pubmed/15897705?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
154. Morales A, Emerson L, Nickel JC, Lundie M. Intravesical hyaluronic acid in the treatment of refractory interstitial cystitis. *J Urol* 1996;156(1):45-48.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8648835&dopt=Abstract

155. Nordling J, Jorgensen S, Kallestrup E. Cystistat for the treatment of interstitial cystitis: a 3-year followup study. *Urology* 2001;57(6 Suppl 1):123.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11378112&dopt=Abstract
156. Kallestrup EB, Jorgensen SS, Nordling J, Hald T. Treatment of interstitial cystitis with Cystistat: a hyaluronic acid product. *Scand J Urol Nephrol* 2005;39(2):143-147.
http://www.ncbi.nlm.nih.gov/pubmed/16032779?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
157. Daha LK, Riedl CR, Lazar D, Hohlbrugger G, Pfluger H. Do cystometric findings predict the results of intravesical hyaluronic acid in women with interstitial cystitis? *Eur Urol* 2005;47(3):393-397; discussion 397.
http://www.ncbi.nlm.nih.gov/pubmed/15716206?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
158. Palylyk-Colwell E. Chondroitin sulfate for interstitial cystitis. *Issues Emerg Health Technol* 2006;1-4.
159. Steinhoff G. The efficacy of chondroitin sulfate in treating interstitial cystitis. *Eur Urol* 2003; suppl 2:14-16.
160. Sorensen RB. Chondroitin sulphate in the treatment of interstitial cystitis and chronic inflammatory disease of the urinary bladder. *Eur Urol* 2003;Suppl 2:16-18.
161. Perez-Marrero R, Emerson LE, Feltis JT. A controlled study of dimethyl sulfoxide in interstitial cystitis. *J Urol* 1988;140(1):36-39.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3288775&dopt=Abstract
162. Sant GR, LaRock DR. Standard intravesical therapies for interstitial cystitis. *Urol Clin North Am* 1994;21(1):73-83.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8284849&dopt=Abstract
163. Rössberger J, Fall M, Peecker R. Critical appraisal of dimethyl sulfoxide treatment for interstitial cystitis: discomfort, side-effects and treatment outcome. *Scand J Urol Nephrol* 2005;39(1):73-77
http://www.ncbi.nlm.nih.gov/pubmed/15764276?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
164. Rowley S, Baer R. Lens deposits associated with DMSO-50 (dimethylsulphoxide). *Eye* 2001;15(Pt 3):332-333.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11450733&dopt=Abstract&itool=iconnoabstr
165. Peters K, Diokno A, Steinert B, Yuhico M, Mitchell B, Krohta S, Gillette B, Gonzalez J. The efficacy of intravesical Tice strain bacillus Calmette-Guerin in the treatment of interstitial cystitis: a double-blind, prospective, placebo controlled trial. *J Urol* 1997;157(6):2090-2094.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9146587&dopt=Abstract
166. Peters KM, Diokno AC, Steinert BW, Gonzalez JA. The efficacy of intravesical bacillus Calmette-Guerin in the treatment of interstitial cystitis: long-term followup. *J Urol* 1998;159(5):1483-1486; discussion 1486-1487.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9554338&dopt=Abstract
167. Mayer R, Probert KJ, Peters KM, Payne CK, Zhang Y, Burks D, Culkin DJ, Diokno A, Hanno P, Landis JR, Madigan R, Messing EM, Nickel JC, Sant GR, Warren J, Wein AJ, Kusek JW, Nyberg LM, Foster HE; Interstitial Cystitis Clinical Trials Group. A randomized controlled trial of intravesical bacillus calmette-guerin for treatment refractory interstitial cystitis. *J Urol* 2005;173(4):1186-1191.
http://www.ncbi.nlm.nih.gov/pubmed/15758738?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
168. Probert KJ, Mayer R, Nickel JC, Payne CK, Peters KM, Teal V, Burks D, Kusek JW, Nyberg LM, Foster HE; Interstitial Cystitis Clinical Trials Group. Did patients with interstitial cystitis who failed to respond to initial treatment with bacillus Calmette-Guerin or placebo in a randomized clinical trial benefit from a second course of open label bacillus Calmette-Guerin? *J Urol* 2007;178(3 Pt 1):886-890
http://www.ncbi.nlm.nih.gov/pubmed/18082224?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
169. Chancellor MB. RTX exotoxins. *Urology* 2001;57(6 Suppl 1):106-107.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11378069&dopt=Abstract

170. Chen TY, Corcos J, Camel M, Ponsot Y, Tu le M. Prospective, randomized, double-blind study of safety and tolerability of intravesical resiniferatoxin (RTX) in interstitial cystitis (IC). *Int Urogynecol J Pelvic Floor Dysfunct* 2005;16(4):293-297.
http://www.ncbi.nlm.nih.gov/pubmed/15818465?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
171. Lazzeri M, Beneforti P, Spinelli M, Zanollo A, Barbagli G, Turini D. Intravesical resiniferatoxin for the treatment of hypersensitive disorder: a randomized placebo controlled study. *J Urol* 2000; 164(3 Pt 1):676-679.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10953124&dopt=Abstract
172. Silva C, Avelino A, Souto-Moura C, Cruz F. A light- and electron-microscopic histopathological study of human bladder mucosa after intravesical resiniferatoxin application. *BJU Int* 2001;88(4):355-360.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11564021&dopt=Abstract
173. Apostolidis A, Gonzales GE, Fowler CJ. Effect of intravesical Resiniferatoxin (RTX) on lower urinary tract symptoms, urodynamic parameters, and quality of life of patients with urodynamic increased bladder sensation. *Eur Urol* 2006;50(6):1299-1305.
http://www.ncbi.nlm.nih.gov/pubmed/16697519?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
174. Payne CK, Mosbaugh PG, Forrest JB, Evans RJ, Whitmore KE, Antoci JP, Perez-Marrero R, Jacoby K, Diokno AC, O'Reilly KJ, Griebing TL, Vasavada SP, Yu AS, Frumkin LR; ICOS RTX Study Group (Resiniferatoxin Treatment for Interstitial Cystitis). Intravesical resiniferatoxin for the treatment of interstitial cystitis: a randomized, double-blind, placebo controlled trial. *J Urol* 2005;173(5):1590-1594.
http://www.ncbi.nlm.nih.gov/pubmed/15821499?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
175. Peng CH, Kuo HC. Multiple intravesical instillations of low-dose resiniferatoxin in the treatment of refractory interstitial cystitis. *Urol Int* 2007;78(1):78-81.
http://www.ncbi.nlm.nih.gov/pubmed/17192738?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
176. Ngan C, Franciosi LG, Butterfield NN, Macleod BA, Jens M, Fenster HN. A prospective, double-blind, randomized cross-over study evaluating changes in urinary pH for relieving the symptoms of interstitial cystitis. *BJU Int* 2005;95(1):91-94.
http://www.ncbi.nlm.nih.gov/pubmed/15638902?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
177. Bumpus HCJ. Interstitial cystitis: its treatment by overdistension of the bladder. *Med Clin North Am* 1930;13:1495-1498.
178. Ormond JK. Interstitial cystitis. *J Urol* 1935;33:576-582.
179. Longacre JJ. The treatment of contracted bladder with controlled tidal irrigation. *J Urol* 1936;36:25-33.
180. Franksson C. Interstitial cystitis: a clinical study of fifty-nine cases. *Acta Chir Scand* 1957;113(1):51-62.
http://www.ncbi.nlm.nih.gov:80/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=13443727&dopt=Abstract
181. Dunn M, Ramsden PD, Roberts JB, Smith JC, Smith PJ. Interstitial cystitis, treated by prolonged bladder distension. *Br J Urol* 1977;49(7):641-645.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=597701&dopt=Abstract
182. Helmstein K. Treatment of bladder carcinoma by a hydrostatic pressure technique. Report on 43 cases. *Br J Urol* 1972;44(4):434-450.
http://www.ncbi.nlm.nih.gov:80/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5070147&dopt=Abstract
183. McCahy PJ, Styles RA. Prolonged bladder distension: experience in the treatment of detrusor overactivity and interstitial cystitis. *Eur Urol* 1995;28(4):325-327.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8575501&dopt=Abstract
184. Zabihi N, Allee T, Maher MG, Mourtzinou A, Raz S, Payne CK, Rodriguez LV. Bladder necrosis following hydrodistention in patients with interstitial cystitis. *J Urol* 2007;177(1):149-52; discussion 152.
http://www.ncbi.nlm.nih.gov/pubmed/17162025?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

185. Glemain P, Riviere C, Lenormand L, Karam G, Bouchot O, Buzelin JM. Prolonged hydrodistention of the bladder for symptomatic treatment of interstitial cystitis: efficacy at 6 months and 1 year. *Eur Urol* 2002;41(1):79-84.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11999471&dopt=Abstract
186. Rose AE, Azevedo KJ, Payne CK. Office bladder distention with electromotive drug administration (EMDA) is equivalent to distention under general anesthesia (GA). *BMC Urol* 2005;5:14.
http://www.ncbi.nlm.nih.gov/pubmed/16300684?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
187. Rose AE, Payne CK, Azevedo K. Pilot study of the feasibility of in-office bladder distention using electromotive drug administration (EMDA). *Neurourol Urodyn* 2005;24(3):254-260.
http://www.ncbi.nlm.nih.gov/pubmed/15747341?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
188. Yamada T, Murayama T, Andoh M. Adjuvant hydrodistension under epidural anesthesia for interstitial cystitis. *Int J Urol* 2003;10(9):463-468
http://www.ncbi.nlm.nih.gov/pubmed/12941123?ordinalpos=7&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
189. Erickson DR, Kunselman AR, Bentley CM, Peters KM, Rovner ES, Demers LM, Wheeler MA, Keay SK. Changes in urine markers and symptoms after bladder distention for interstitial cystitis. *J Urol* 2007;177(2):556-560.
http://www.ncbi.nlm.nih.gov/pubmed/17222633?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
190. Cole EE, Scarpero HM, Dmochowski RR. Are patient symptoms predictive of the diagnostic and/or therapeutic value of hydrodistention? *Neurourol Urodyn* 2005;24(7):638-642.
http://www.ncbi.nlm.nih.gov/pubmed/16208660?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
191. Gulpinar T, Wong HY, Griffith DP. Electromotive administration of intravesical lidocaine in patients with interstitial cystitis. *J Endourol* 1996;10(5):443-447.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8905491&dopt=Abstract
192. Rosamilia A, Dwyer PL, Gibson J. Electromotive drug administration of lidocaine and dexamethasone followed by cystodistension in women with interstitial cystitis. *Int Urogynecol J Pelvic Floor Dysfunct* 1997;8(3):142-145.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9449586&dopt=Abstract&itool=iconabstr
193. Riedl CR, Knoll M, Plas E, Pfluger H. Electromotive drug administration and hydrodistention for the treatment of interstitial cystitis. *J Endourol* 1998;12(3):269-272.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9658301&dopt=Abstract
194. Kerr WS Jr. Interstitial cystitis: treatment by transurethral resection. *J Urol* 1971;105(5):664-666.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=4397018&dopt=Abstract
195. Fall M. Conservative management of chronic interstitial cystitis: transcutaneous electrical nerve stimulation and transurethral resection. *J Urol* 1985;133(5):774-778.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3872946&dopt=Abstract
196. Peeker R, Aldenborg F, Fall M. Complete transurethral resection of ulcers in classic interstitial cystitis. *Int Urogynecol J Pelvic Floor Dysfunct* 2000;11(5):290-295.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11052564&dopt=Abstract
197. Shanberg AM, Baghdassarian R, Tansey LA. Treatment of interstitial cystitis with the neodymium-YAG laser. *J Urol* 1985;134(5):885-888.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3840538&dopt=Abstract
198. Malloy TR, Shanberg AM. Laser therapy for interstitial cystitis. *Urol Clin North Am* 1994; 21(1):141-144.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8284837&dopt=Abstract

199. Rofeim O, Hom D, Freid RM, Moldwin RM. Use of the neodymium: yag laser for interstitial cystitis: a prospective study. *J Urol* 2001;166(1):134-136.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11435840&dopt=Abstract
200. Smith CP, Radziszewski P, Borkowski A, Somogyi GT, Boone TB, Chancellor MB. Botulinum toxin a has antinociceptive effects in treating interstitial cystitis. *Urology* 2004;64(5):871-5; discussion 875.
http://www.ncbi.nlm.nih.gov/pubmed/15533466?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
201. Kuo HC. Preliminary results of suburothelial injection of botulinum a toxin in the treatment of chronic interstitial cystitis. *Urol Int* 2005;75(2):170-174.
http://www.ncbi.nlm.nih.gov/pubmed/16123573?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
202. van Ophoven A, Rossbach G, Oberpenning F, Hertle L. Hyperbaric oxygen for the treatment of interstitial cystitis: long-term results of a prospective pilot study. *Eur Urol* 2004 Jul;46(1):108-113.
http://www.ncbi.nlm.nih.gov/pubmed/15183555?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
203. van Ophoven A, Rossbach G, Pajonk F, Hertle L. Safety and efficacy of hyperbaric oxygen therapy for the treatment of interstitial cystitis: a randomized, sham controlled, double-blind trial. *J Urol* 2006;176(4 Pt 1):1442-1446.
http://www.ncbi.nlm.nih.gov/pubmed/16952654?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
204. Dasgupta P, Sharma SD, Womack C, Blackford HN, Dennis P. Cimetidine in painful bladder syndrome: a histopathological study. *BJU Int* 2001;88(3):183-186.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11488726&dopt=Abstract
205. Thilagarajah R, Witherow RO, Walker MM. Oral cimetidine gives effective symptom relief in painful bladder disease: a prospective, randomized, double-blind placebo-controlled trial. *BJU Int* 2001;87(3):207-212.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11167643&dopt=Abstract
206. Kelly JD, Young MR, Johnston SR, Keane PF. Clinical response to an oral prostaglandin analogue in patients with interstitial cystitis. *Eur Urol* 1998;34(1):53-56.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9676414&dopt=Abstract
207. Korting GE, Smith SD, Wheeler MA, Weiss RM, Foster HE Jr. A randomized double-blind trial of oral L-arginine for treatment of interstitial cystitis. *J Urol* 1999;161(2):558-565.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9915448&dopt=Abstract
208. Wheeler MA, Smith SD, Saito N, Foster HE Jr, Weiss RM. Effect of long-term oral L-arginine on the nitric oxide synthase pathway in the urine from patients with interstitial cystitis. *J Urol* 1997;158(6):2045-2050.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9366309&dopt=Abstract
209. Smith SD, Wheeler MA, Foster HE Jr, Weiss RM. Improvement in interstitial cystitis symptom scores during treatment with oral L-arginine. *J Urol* 1997;158(3 Pt 1):703-708.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9258064&dopt=Abstract
210. Lundberg JO, Ehren I, Jansson O, Adolfsson J, Lundberg JM, Weitzberg E, Alving K, Wiklund NP. Elevated nitric oxide in the urinary bladder in infectious and noninfectious cystitis. *Urology* 1996;48(5):700-702.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8911512&dopt=Abstract
211. Ehren I, Lundberg JO, Adolfsson J, Wiklund NP. Effects of L-arginine treatment on symptoms and bladder nitric oxide levels in patients with interstitial cystitis. *Urology* 1998;52(6):1026-1029.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9836549&dopt=Abstract
212. Cartledge JJ, Davies AM, Eardley I. A randomized double-blind placebo-controlled crossover trial of the efficacy of L-arginine in the treatment of interstitial cystitis. *BJU Int* 2000;85(4):421-426.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10691818&dopt=Abstract

213. Barbalias GA, Liatsikos EN, Athanasopoulos A, Nikiforidis G. Interstitial cystitis: bladder training with intravesical oxybutynin. *J Urol* 2000;163(6):1818-1822.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10799190&dopt=Abstract
214. van Ophoven A, Hertle L. The dual serotonin and noradrenaline reuptake inhibitor duloxetine for the treatment of interstitial cystitis: results of an observational study. *J Urol* 2007;177(2):552-555
http://www.ncbi.nlm.nih.gov/pubmed/17222632?ordinalpos=10&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
215. O'Connor VJ. Cloripactin WCS90 in the treatment of interstitial cystitis. *Q Bull Northwest Univ Med Sch* 1955;29(4):292-295.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=13273619&dopt=Abstract&itool=iconnoabstr
216. Wishard WN, Nourse MH, Mertz JHO. Use of Cloripactin WCS90 for relief of symptoms due to interstitial cystitis. *J Urol* 1957;77(3):420-423.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=13417272&dopt=Abstract&itool=iconnoabstr
217. Messing EM, Freiha FS. Complication of Cloripactin WCS90 therapy for interstitial cystitis. *Urology* 1979;13(4):389-392.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=219578&dopt=Abstract
218. Murnaghan GF, Salafeld J, Farnworth RH. Interstitial cystitis - treatment with cloripactin WCS90. *Br J Urol* 1969;42:744.
219. von Heyden B, Schmid HP. [Intravesical therapy of interstitial cystitis.] *Urologe A* 2000;39(6):542-544. [German].
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11138274&dopt=Abstract
220. Hanno P. Interstitial cystitis and related diseases. In: Walsh PC, Retik AB, Stamey TA, Vaughan ED, eds. *Campbell's Urology*. Philadelphia: WB Saunders Co., 1998, pp. 648.
221. Parsons CL, Koprowski PF. Interstitial cystitis: successful management by increasing urinary voiding intervals. *Urology* 1991;37(3):207-212.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2000675&dopt=Abstract
222. Chaiken DC, Blaivas JG, Blaivas ST. Behavioral therapy for the treatment of refractory interstitial cystitis. *J Urol* 1993;149(6):1445-1448.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8501784&dopt=Abstract
223. Webster DC, Brennan T. Use and effectiveness of physical self-care strategies for interstitial cystitis. *Nurse Pract* 1994;19(10):55-61.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7529390&dopt=Abstract
224. Rovner E, Propert KJ, Brensinger C, Wein AJ, Foy M, Kirkemo A, Landis JR, Kusek JW, Nyberg LM. Treatments used in women with interstitial cystitis: the interstitial cystitis data base (ICDB) study experience. The Interstitial Cystitis Data Base Study Group. *Urology* 2000;56(6):940-945.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11113737&dopt=Abstract
225. Bade JJ, Peeters JM, Mensink HJ. Is the diet of patients with interstitial cystitis related to their disease? *Eur Urol* 1997;32(2):179-183.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9286650&dopt=Abstract
226. Gillespie L. Metabolic appraisal of the effects of dietary modification on hypersensitive bladder symptoms. *Br J Urol* 1993;72(3):293-297.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8220989&dopt=Abstract
227. Bologna RA, Gomelsky A, Lukban JC, Tu LM, Holzberg AS, Whitmore KE. The efficacy of calcium glycerophosphate in the prevention of food-related flares in interstitial cystitis. *Urology* 2001;57(6 Suppl 1):119-120.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11378102&dopt=Abstract


228. Osborne JH, Manhattan D, Laumnn B. IC and Diet. In: Osborne JH , ed. *The Interstitial Cystitis Network Patient Handbook. Chapter 5*. Santa Rosa, CA, USA: The Interstitial Cystitis Network (www.ic-network.com), 1999, pp. 43-62 (access date February 2007).
<http://www.ic-network.com/handboo>
229. Chang PL. Urodynamic studies in acupuncture for women with frequency, urgency and dysuria. *J Urol* 1988;140(3):563-566.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3411675&dopt=Abstract
230. Chang PL, Wu CJ, Huang MH. Long-term outcome of acupuncture in women with frequency, urgency and dysuria. *Am J Chin Med* 1993;21(3-4):231-236.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8135166&dopt=Abstract
231. Zheng H, Wang S, Shang J, Chen G, Huang C, Hong H, Chen S. Study on acupuncture and moxibustion therapy for female urethral syndrome. *J Tradit Chin Med* 1998;18(2):122-127.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10437230&dopt=Abstract
232. Geirsson G, Wang YH, Lindstrom S, Fall M. Traditional acupuncture and electrical stimulation of the posterior tibial nerve. A trial in chronic interstitial cystitis. *Scand J Urol Nephrol* 1993;27(1):67-70.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8493470&dopt=Abstract
233. Lynch DF Jr. Empowering the patient: hypnosis in the management of cancer, surgical disease and chronic pain. *Am J Clin Hypn* 1999;42(2):122-130.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10624023&dopt=Abstract
234. Barber J. Incorporating hypnosis in the management of chronic pain. In: Barber J, Adrian C, eds. *Psychological Approaches in the Management of Pain*. New York: Brunner/Mazel, 1982: 60-83.
235. Karper WB. Exercise effects on interstitial cystitis: two case reports. *Urol Nurs* 2004;24(3):202-204.
http://www.ncbi.nlm.nih.gov/pubmed/15311489?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
236. Oyama IA, Rejba A, Lukban JC, Fletcher E, Kellogg-Spadt S, Holzberg AS, Whitmore KE. Modified Thiele massage as therapeutic intervention for female patients with interstitial cystitis and high-tone pelvic floor dysfunction. *Urology* 2004;64(5):862-865.
http://www.ncbi.nlm.nih.gov/pubmed/15533464?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
237. Langford CF, Udvari Nagy S, Ghoniem GM. Levator ani trigger point injections: An underutilized treatment for chronic pelvic pain. *Neurourol Urodyn* 2007;26(1):59-62.
http://www.ncbi.nlm.nih.gov/pubmed/17195176?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
238. de Oliveira Bernardes N, Bahamondes L. Intravaginal electrical stimulation for the treatment of chronic pelvic pain. *J Reprod Med* 2005;50(4):267-272.
http://www.ncbi.nlm.nih.gov/pubmed/15916211?ordinalpos=8&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
239. van Ophoven A, Oberpenning F, Hertle L. Long-term results of trigone-preserving orthotopic substitution enterocystoplasty for interstitial cystitis. *J Urol* 2002;167(2 Pt 1):603-607.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11792927&dopt=Abstract
240. Loch A, Stein U. [Interstitial cystitis. New aspects in diagnosis and therapy]. *Urologe A* 2004;43(9):1135-1146. [German].
http://www.ncbi.nlm.nih.gov/pubmed/15322757?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
241. Oberpenning F, van Ophoven A, Hertle L. [Chronic interstitial cystitis.] *Deutsches Ärzteblatt* 2002, 99:204-208. [German]
242. Oberpenning F, Van Ophoven A, Hertle L. Interstitial cystitis: an update. *Curr Opin Urol* 2002; 12(4):321-332.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=12072654&dopt=Abstract

243. Turner-Warwick R, Ashkan M. The functional results of partial, subtotal and total cystoplasty with special reference to ureterocecocystoplasty, selective sphincterotomy and cystoplasty. *Br J Urol* 1967;39(1):3-12.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5336762&dopt=Abstract
244. Freiha FS, Faysal MH, Stamey TA. The surgical treatment of intractable interstitial cystitis. *J Urol* 1980;123(5):632-634.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7420547&dopt=Abstract
245. Awad SA, Al-Zahrani HM, Gajewski JB, Bourque-Kehoe AA. Long-term results and complications of augmentation ileocystoplasty for idiopathic urge incontinence in women. *Br J Urol* 1998;81(4):569-573.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9598629&dopt=Abstract
246. Bruce PT, Buckham GJ, Carden AB, Salvaris M. The surgical treatment of chronic interstitial cystitis. *Med J Aust* 1977;1(16):581-582.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=875802&dopt=Abstract
247. Christmas TJ, Holmes SA, Hendry WF. Bladder replacement by ileocystoplasty: the final treatment for interstitial cystitis. *Br J Urol* 1996;78(1):69-73.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8795403&dopt=Abstract
248. von Garrelts B. Interstitial cystitis: thirteen patients treated operatively with intestinal bladder substitutes. *Acta Chir Scand* 1966;132(4):436-443.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5972716&dopt=Abstract
249. Guillonneau B, Toussaint B, Bouchot O, Buzelin JM. [Treatment of interstitial cystitis with sub-trigonal cystectomy and enterocystoplasty.] *Prog Urol* 1993;3(1):27-31 [French]
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8485591&dopt=Abstract
250. Koskela E, Kontturi M. Function of the intestinal substituted bladder. *Scand J Urol Nephrol* 1982;16(2):129-133.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7123162&dopt=Abstract
251. Shirley SW, Mirelman S. Experiences with colocytoplasties, cecocystoplasties and ileocystoplasties in urologic surgery: 40 patients. *J Urol* 1978;120(2):165-168.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=671623&dopt=Abstract
252. Webster GD, Maggio MI. The management of chronic interstitial cystitis by substitution cystoplasty. *J Urol* 1989;141(2):287-291.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2913346&dopt=Abstract
253. Nielsen KK, Kromann-Andersen B, Steven K, Hald T. Failure of combined supratrigonal cystectomy and Mainz ileocececystoplasty in intractable interstitial cystitis: is histology and mast cell count a reliable predictor for the outcome of surgery? *J Urol* 1990;144(2 Pt 1):255-258; discussion 258-259.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2374189&dopt=Abstract
254. Hradec EA. Bladder substitution: indications and results in 114 operations. *J Urol* 1965;94(4):406-417.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5320331&dopt=Abstract
255. DeJuana CP, Everett JC Jr. Interstitial cystitis: experience and review of recent literature. *Urology* 1977;10(4):325-329.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=919117&dopt=Abstract
256. Utz DC, Zincke H. The masquerade of bladder cancer in situ as interstitial cystitis. *J Urol* 1974;111(2):160-161.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=4810754&dopt=Abstract

257. Whitmore WF 3rd, Gittes RF. Reconstruction of the urinary tract by cecal and ileocecal cystoplasty: review of a 15-year experience. *J Urol* 1983;129(3):494-498.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6834531&dopt=Abstract
258. Kontturi MJ, Hellstrom PA, Tammela TL, Lukkarinen OA. Colocystoplasty for the treatment of severe interstitial cystitis. *Urol Int* 1991;46(1):50-54.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2024372&dopt=Abstract
259. Seddon JM, Best L, Bruce AW. Intestinocystoplasty in treatment of interstitial cystitis. *Urology* 1977;10(5):431-435.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=919133&dopt=Abstract
260. Leong CH. Use of the stomach for bladder replacement and urinary diversion. *Ann R Coll Surg Engl* 1978;60(4):283-289.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=666231&dopt=Abstract
261. Singla A, Galloway N. Early experience with the use of gastric segment in lower urinary tract reconstruction in adult patient population. *Urology* 1997;50(4):630-635.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9338749&dopt=Abstract
262. Dounis A, Gow JG. Bladder augmentation-a long-term review. *Br J Urol* 1979;51(4):264-268.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=466001&dopt=Abstract
263. Linn JF, Hohenfellner M, Roth S, Dahms SE, Stein R, Hertle L, Thüroff JW, Hohenfellner R. Treatment of interstitial cystitis: comparison of subtrigonal and supratrigonal cystectomy combined with orthotopic bladder substitution. *J Urol* 1998;159(3):774-778.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9474146&dopt=Abstract
264. Chakravarti A, Ganta S, Somani B, Jones MA. Caecocystoplasty for intractable interstitial cystitis: long-term results. *Eur Urol* 2004 Jul;46(1):114-117.
http://www.ncbi.nlm.nih.gov/pubmed/15183556?ordinalpos=8&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
265. Blaivas JG, Weiss JP, Desai P, Flisser AJ, Stember DS, Stahl PJ. Long-term followup of augmentation enterocystoplasty and continent diversion in patients with benign disease. *J Urol* 2005;173(5):1631-1634.
http://www.ncbi.nlm.nih.gov/pubmed/15821519?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
266. Navalón Verdejo P, Ordoño Domínguez F, De la Torre Abril L, Sánchez Ballester F, Escudero JJ, Ramos de Campos M, Ramada Benlloch F. [Orthotopic bladder substitution in the treatment of interstitial cystitis.] *Arch Esp Urol* 2005;58(7):605-610. [Spanish]
http://www.ncbi.nlm.nih.gov/pubmed/16294782?ordinalpos=6&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
267. Bejany DE, Politano VA. Ileocolic neobladder in the woman with interstitial cystitis and a small contracted bladder. *J Urol* 1995;153(1):42-43.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7966787&dopt=Abstract
268. Nurse DE, McCrae P, Stephenson TP, Mundy AR. The problems of substitution cystoplasty. *Br J Urol* 1988;61(5):423-426.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3395801&dopt=Abstract
269. Lotenfoe RR, Christie J, Parsons A, Burkett P, Helal M, Lockhart JL. Absence of neuropathic pelvic pain and favorable psychological profile in the surgical selection of patients with disabling interstitial cystitis. *J Urol* 1995;154(6):2039-2042.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7500453&dopt=Abstract
270. Hughes OD, Kynaston HG, Jenkins BJ, Stephenson TP, Vaughton KC. Substitution cystoplasty for intractable interstitial cystitis. *Br J Urol* 1995;76(2):172-174.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7663907&dopt=Abstract

271. Nurse DE, Parry JR, Mundy AR. Problems in the surgical treatment of interstitial cystitis. Br J Urol 1991;68(2):153-154.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1822961&dopt=Abstract
272. Volkmer BG, Gschwend JE, Herkommer K, Simon J, Küfer R, Hautmann RE. Cystectomy and orthotopic ileal neobladder: the impact on female sexuality. J Urol 2004;172(6 Pt 1):2353-2357.
http://www.ncbi.nlm.nih.gov/pubmed/15538266?ordinalpos=13&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
273. Peeker R, Aldenborg F, Fall M. The treatment of interstitial cystitis with supratrigonal cystectomy and ileocystoplasty: difference in outcome between classic and nonulcer disease. J Urol 1998; 159(5):1479-1482.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9554337&dopt=Abstract
274. Rössberger J, Fall M, Jonsson O, Peeker R. Long-term results of reconstructive surgery in patients with bladder pain syndrome/interstitial cystitis: subtyping is imperative. Urology 2007;70(4):638-642.
http://www.ncbi.nlm.nih.gov/sites/entrez?Db=pubmed&Cmd=ShowDetailView&TermToSearch=17991529&ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
275. Gershbaum D, Moldwin R. Practice trends for the management of interstitial cystitis. Urology 2001;57(6 Suppl 1):119.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11378100&dopt=Abstract
276. Elzawahri A, Bissada NK, Herchorn S, Aboul-Enein H, Ghoneim M, Bissada MA, Finkbeiner A, Glazer AA. Urinary conduit formation using a retubularized bowel from continent urinary diversion or intestinal augmentations: ii. Does it have a role in patients with interstitial cystitis? J Urol 2004;171(4):1559-1562.
http://www.ncbi.nlm.nih.gov/pubmed/15017220?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
277. Shaikh A, Ahsan S, Zaidi Z. Pregnancy after augmentation cystoplasty. J Pak Med Assoc 2006;56(10):465-467.
http://www.ncbi.nlm.nih.gov/pubmed/17144396?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

Figure 4: An algorithm for diagnosing and managing scrotal pain



2.8 Scrotal pain

Acute scrotal pain is a well-known urological emergency, while chronic scrotal pain is a common reason for men to contact their general practitioners. However, although chronic scrotal pain is not life-threatening and does not require immediate action, it has a considerable effect on a patient's quality of life. The prevalence of chronic scrotal pain is unknown.

A physical examination should always be done in patients with scrotal pain. Gentle palpation of each component of the scrotum is performed to search for masses and for painful spots. A DRE is done to look for prostate abnormalities and examine the pelvic floor muscles. Scrotal ultrasound has limited value in finding the cause of the pain (1).

If physical examination is normal, ultrasound is sometimes performed to reassure the patient that there is no tumour in the testis. Ultrasound can be used to diagnose hydroceles, spermatoceles, cysts and varicoceles. The urine should be analysed. Magnetic resonance imaging and computed tomography scans may be used to help with assessment (2).

The ilioinguinal and genitofemoral nerves are the most prominent afferent nerves for the scrotum (3). The pudendal nerve supplies the skin of the perineum.

2.8.1 Management of different conditions

Intra-testicular lesion. Proceed with further analysis. If tumour is suspected, orchiectomy should be performed.

Hydro/spermatocele. Painful fluid-filled spaces in the scrotum can be removed surgically.

Varicocele. When localization of the pain and the pattern of aggravation in standing position are clear, correction can be performed.

Epididymo-orchitis. An infection of the testis or epididymis is usually an acute problem. Chronic epididymitis has been discussed as an entity in literature. The nature of such a chronic inflammation can be infective or based on an obstruction of the vas deferens. For treatment, antibiotics should be tried first for a longer period, up to 3 months, and when needed combined with anti-inflammatory drugs (4).

Post-surgical procedures. Procedures, such as vasectomy and hernia repair, may be complicated by scrotal pain. Scrotal pain after vasectomy occurs in about 15-19% (5,6) and is caused by congestion of the vas deferens and testis. Although antibiotics can be used, the results remain unclear. A spermatic cord blockade is also an option (7).

Surgical options for both vasectomy and hernia repair include removal of the epididymis, with recent results varying from 43-62% (6,8). Results for denervation of the spermatic cord are reported to be as high as 96% for complete pain relief (9). In post-vasectomy pain, a vasovasostomy might help to overcome the obstruction and thereby improve the pain (10).

Pelvic floor muscle dysfunction. At rectal examination, the pelvic floor muscles may be overactive, which means they contract when relaxation is needed, sometimes painfully. An overactive pelvic floor should be treated by physiotherapy (11-13). (See Chapter 6 on Pelvic floor function and dysfunction.)

Myofascial trigger points is a type of end-stage surmenage of muscles. Pain in the scrotum can be the result of trigger points in the pelvic floor, but also in the lower abdominal musculature. Treatment consists of applying pressure to the trigger point and stretching the muscle (14,15). (See Chapter 6 on Pelvic floor function and dysfunction.)

If no pathology is found, or when specific therapy has no effect, the patient should be referred to a multidisciplinary pain team or pain centre (16).

Recommendations for the treatment of scrotal pain syndrome are listed in Table 12.

Table 12: Treatment of scrotal pain syndrome

	Level of evidence	Grade of recommendation	Comment
• Orchiectomy	1a	A	In case of intratesticular tumour
• Excision	3	B	Hydrocele or varicocele
• Antibiotics	3	C	For up to 3 months

•	Surgical intervention	3	C	Epididymectomy, denervation spermatic cord. Vaso-vasostomy
•	Pelvic floor muscle therapy	1b	A	Including trigger point treatment


2.8.2 References

1. van Haast EP, van Andel G, Rijcken TH, Schlatmann TJ, Taconis WK. Value of diagnostic ultrasound in patients with chronic scrotal pain and normal findings on clinical examination. *Urology* 1999;54(6):1068-1072.
http://www.ncbi.nlm.nih.gov/pubmed/10604710?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
2. Lapointe SP, Wei DC, Hricak H, Varghese SL, Kogan BA, Baskin LS. Magnetic resonance imaging in the evaluation of congenital anomalies of the external genitalia. *Urology* 2001;58(3):452-456.
http://www.ncbi.nlm.nih.gov/pubmed/11549498?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
3. Rab M, Ebmer AJ, Dellon AL. Anatomic variability of the ilioinguinal and genitofemoral nerve: implications for the treatment of groin pain. *Plast Reconstr Surg* 2001;108(6):1618-1623.
http://www.ncbi.nlm.nih.gov/pubmed/11711938?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
4. Nickel JC. Chronic epididymitis: a practical approach to understanding and managing a difficult urologic enigma. *Rev Urol* 2003;5(4):209-215.
http://www.ncbi.nlm.nih.gov/pubmed/16985840?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
5. Leslie TA, Illing RO, Cranston DW, Guillebaud J. The incidence of chronic scrotal pain after vasectomy: a prospective audit. *BJU Int* 2007;100(6):1330-1333.
http://www.ncbi.nlm.nih.gov/pubmed/17850378?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
6. Nariculam J, Minhas S, Adeniyi A, Ralph DJ, Freeman A. A review of the efficacy of surgical treatment for and pathological changes in patients with chronic scrotal pain. *BJU Int* 2007;99(5):1091-1093.
http://www.ncbi.nlm.nih.gov/pubmed/17244279?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
7. Yamamoto M, Hibi H, Katsuno S, Miyake K. Management of chronic orchialgia of unknown etiology. *Int J Urol* 1995;2(1):47-49.
http://www.ncbi.nlm.nih.gov/pubmed/7542163?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
8. Granitsiotis P, Kirk D. Chronic testicular pain: an overview. *Eur Urol* 2004;45(4):430-436.
http://www.ncbi.nlm.nih.gov/pubmed/15041105?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
9. Heidenreich A, Olbert P, Engelmann UH. Management of chronic testalgia by microsurgical testicular denervation. *Eur Urol* 2002;41(4):392-397.
http://www.ncbi.nlm.nih.gov/pubmed/12074809?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
10. Nangia AK, Myles JL, Thomas AJ Jr. Vasectomy reversal for the post-vasectomy pain syndrome: a clinical and histological evaluation. *J Urol* 2000;164(6):1939-1942.
http://www.ncbi.nlm.nih.gov/pubmed/11061886?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
11. Cornel EB, van Haast EP, Schaarsberg RW, Geels J. The effect of biofeedback physical therapy in men with Chronic Pelvic Pain Syndrome Type III. *Eur Urol* 2005;47(5):607-611.
http://www.ncbi.nlm.nih.gov/pubmed/15826751?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
12. Hetrick DC, Glazer H, Liu YW, Turner JA, Frest M, Berger RE. Pelvic floor electromyography in men with chronic pelvic pain syndrome: a case-control study. *Neurourol Urodyn* 2006;25(1):46-49.
http://www.ncbi.nlm.nih.gov/pubmed/16167354?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

13. Rowe E, Smith C, Laverick L, Elkabir J, Witherow RO, Patel A. A prospective, randomized, placebo controlled, double-blind study of pelvic electromagnetic therapy for the treatment of chronic pelvic pain syndrome with 1 year of followup. J Urol 2005;173(6):2044-2047.
http://www.ncbi.nlm.nih.gov/pubmed/15879822?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
14. Anderson RU, Wise D, Sawyer T, Chan C. Integration of myofascial trigger point release and paradoxical relaxation training treatment of chronic pelvic pain in men. J Urol 2005;174(1):155-160.
http://www.ncbi.nlm.nih.gov/pubmed/15947608?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
15. Srinivasan AK, Kaye JD, Moldwin R. Myofascial dysfunction associated with chronic pelvic floor pain: management strategies. Curr Pain Headache Rep 2007;11(5):359-364.
http://www.ncbi.nlm.nih.gov/pubmed/17894926?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
16. Messelink EJ. The pelvic pain centre. World J Urol 2001;19(3):208-212.
http://www.ncbi.nlm.nih.gov/pubmed/11469609?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

2.9 Urethral pain syndrome

Figure 5: An algorithm for diagnosing and managing urethral pain syndrome



Urethral pain syndrome is a less well-defined entity and scientific studies are scant. Positive diagnostic signs are urethral tenderness or pain on palpation and a slightly inflamed urethral mucosa found during endoscopy. Hypotheses about the aetiology include concealed infections of the periurethral glands or ducts, according to the anatomical description by Huffman (1), and oestrogen deficiency. Others consider urethral syndrome to be a less severe form of 'early' BPS/IC (2).

In clinical practice, the diagnosis of urethral pain syndrome is commonly given to patients who present with the symptoms of dysuria (with or without frequency, nocturia, urgency and urge incontinence) in the absence of evidence of urinary infection. The 'absence of urinary infection' cause diagnostic problems as the methods typically used to identify urinary infection are extremely insensitive. Dysuria is pain or discomfort experienced in association with micturition. The classical symptom of a burning sensation in the urethra during voiding caused by infection is well known. The external dysuria experienced by women with vaginitis when urine passes over the labia is less appreciated.

Biochemical testing and microbiological culture of urine is important in assessing lower urinary tract symptoms and has been reviewed in some detail in the elderly (3).

There is confusion about the concept of significant bacteriuria. This may be accepted as 10^5 colony-forming units (CFU) of a single species in asymptomatic women. However, it may be as low as 10^2 CFU of a single species of a known urinary pathogen in symptomatic women. Many automated culture systems have a sensitivity of 10^4 CFU, while urinary leucocyte esterase and nitrite tests are correlated only with cultures as high as 10^5 CFU (4). In addition, many laboratory culture systems detect only just over 50% of infections in midstream urine specimens from genuinely infected patients (4).

A narrow spectrum of aetiological agents causes 85-90% of cases of acute, uncomplicated cystitis in women. Nearly one-third of acutely dysuric women with urinary tract infections caused by *Escherichia coli*, *Staphylococcus saprophyticus* or *Proteus spp.* have midstream colony counts in the range of 10^2 - 10^4

bacteria/mL. Investigators have also identified causative organisms by more invasive techniques, such as culturing specimens obtained by catheterization or suprapubic aspiration. Failing to identify an organism does not mean that it is not present.

Although rarely included, proper manual urine microscopy using a haemocytometer should be part of a definitive work-up. Nowadays, most laboratories screen urine in wells using inverted microscopes or rely on robotic detection of pyuria, which are both insensitive methods. This is regrettable because studies have shown that significant pyuria is a nearly universal indicator of urinary tract infection, although it is not specific for differentiating cystitis from urethritis, particularly urethritis due to *Chlamydia trachomatis*. In relation to the latter, dysuria also merits the microscopic examination of a urethral smear after it has been Gram stained. If present, a purulent urethral exudate will be obvious, although a causative micro-organism will be identified in less than 50% of cases. The expression 'non-specific urethritis' is apposite and honestly states our current ignorance.

Urethral trauma arising from intercourse may cause pain and dysuria. This condition used to be called 'honeymoon cystitis', and friction and trauma to the urethra may be the cause in the absence of infection. Women with pelvic floor dysfunction sometimes describe similar symptoms, as do post-menopausal women, in whom trauma is associated with oestrogen deficiency, loss of lubrication and vaginal dryness.

Unless a thorough assessment is carried out, bearing in mind the comments described above, the diagnosis of urethral pain syndrome does not seem credible. There are no data available to answer the inevitable question, 'how common is dysuria in the presence of negative rigorous investigation of the bladder and urethra?'

2.9.1 Treatment

There is no consensus on treatment. Management may require a multidisciplinary approach. Various modalities including antibiotics, alpha-blockers, acupuncture and laser therapy have been proved successful. Psychological support is important (5). An algorithm for diagnosing and managing urethral pain syndrome is given in Figure 5.

2.9.2 References

1. Huffman JW. The detailed anatomy of the paraurethral ducts in the adult human female. *Am J Obstet Gynec* 1948;55:86-101.
2. Parsons CL, Zupkas P, Parsons JK. Intravesical potassium sensitivity in patients with interstitial cystitis and urethral syndrome. *Urology* 2001;57(3):428-432.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11248610&dopt=Abstract
3. Gray RP, Malone-Lee J. Review: urinary tract infection in elderly people - time to review management? *Age & Ageing* 1995;24(4):341-345.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7484494&dopt=Abstract
4. Pappas PG. Laboratory in the diagnosis and management of urinary tract infections. *Med Clin North Am* 1991;75(2):313-325.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1996036&dopt=Abstract
5. Kaur H, Arunkalaivanan AS. Urethral pain syndrome and its management. *Obstet Gynecol Surv* 2007;62(5):348-351.
http://www.ncbi.nlm.nih.gov/pubmed/17425813?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

3. PELVIC PAIN IN GYNAECOLOGICAL PRACTICE

3.1 Introduction

The approach to pelvic pain presenting to the gynaecologist relies upon the same principles, namely to discover remediable causes and treat them using the most effective available therapies. However, the greatest therapeutic challenge will be provided by the 30% of patients in whom no cause can be found (1).

3.2 Clinical history

It is essential to start by taking a detailed medical history. The nature, frequency and site of the pain, and its relationship to precipitating factors and the menstrual cycle, may provide vital clues to the aetiology. A detailed

menstrual and sexual history, including any history of sexually transmitted diseases and vaginal discharge is mandatory. Discrete inquiry about previous sexual trauma may be appropriate.

3.3 Clinical examination

Abdominal and pelvic examination will exclude any gross pelvic pathology (tumours, scarring and reduced uterine mobility), as well as demonstrating the site of tenderness if present. Abnormalities in muscle function should also be sought.

3.3.1 Investigations

Vaginal and endocervical swabs to exclude infection are mandatory and cervical cytology screening is advisable. Pelvic ultrasound scanning provides further information about pelvic anatomy and pathology. Laparoscopy is the most useful invasive investigation to exclude gynaecological pathology (2) and to assist in the differential diagnosis (3).

3.4 Dysmenorrhoea

Pain in association with menstruation may be primary or secondary.

Primary dysmenorrhoea classically begins at the onset of ovulatory menstrual cycles and tends to decrease following childbirth (4). Explanation and reassurance may be helpful, together with the use of simple analgesics progressing to the use of non-steroidal anti-inflammatory drugs (NSAIDs), which are particularly helpful if they are started before the onset of menstruation. NSAIDs are effective in dysmenorrhoea probably because of their effects on prostaglandin synthetase. Suppression of ovulation using the oral contraceptive pill reduces dysmenorrhoea dramatically in most cases and may be used as a therapeutic test. Because of the chronic nature of the condition, potentially addictive analgesics should be avoided.

Secondary dysmenorrhoea suggests the development of a pathological process and it is essential to exclude endometriosis (5) and pelvic infection.

3.5 Infection

A history of possible exposure to infection should be sought and it is mandatory in all cases to obtain swabs to exclude chlamydia and gonorrhoea, as well as vaginal and genital tract pathogens (6). Patient's sexual contacts need to be traced in all cases with a positive culture. If there is any doubt about the diagnosis, laparoscopy may be very helpful.

Primary herpes simplex infection may present with severe pain (7), associated with an ulcerating lesion and inflammation, which may lead to urinary retention (8). Hospitalization and opiates may be needed to achieve adequate analgesia.

3.5.1 Treatment

Treatment of infection depends on the causative organisms. Subclinical chlamydial infection may lead to tubal pathology. Screening for this organism in sexually active young women may reduce the incidence of subsequent subfertility.

Chronic pelvic inflammatory disease is no longer common in developed countries, but still poses a significant problem with chronic pain in the Third World.

3.6 Endometriosis

The incidence of endometriosis is rising in the developed world. The precise aetiology is still a source of debate, but an association with nulliparity is well accepted.

The condition may be suspected from a history of secondary dysmenorrhoea and often dyspareunia, as well as the finding of scarring in the vaginal fornices on vaginal examination, with reduced uterine mobility and adnexal masses. Laparoscopy is the most useful diagnostic tool (9, 10).

Endometriotic lesions affecting the urinary bladder or causing ureteric obstructions can occur, as well as lesions affecting the bowel, which may lead to rectal bleeding in association with menstruation.

3.6.1 Treatment

As in primary dysmenorrhoea, analgesics and NSAIDs are helpful in easing pain at the time of menstruation. Hormone treatment with progestogens or the oral contraceptive pill may halt progress of endometriosis, but is not curative. A temporary respite may be obtained by using luteinizing hormone releasing hormone (LHRH) analogues to create an artificial menopause, though the resulting oestrogen deficiency may have marked long-term side effects, such as reduced bone density and osteoporosis in those taking more than six months worth of treatment. These drugs are used prior to surgery to improve surgical outcome and reduce surgical complications.

Surgery for endometriosis is challenging and the extensive removal of all endometriotic lesions is

essential. The best results are achieved laparoscopically, by highly trained and skilled laparoscopic surgeons, in specialist centres (11). A multidisciplinary team will be required for the treatment of extensive disease, including a pain management team.

The pain associated with endometriosis is often not proportionate to the extent of the condition and, even after extensive removal of the lesions and suppression of the condition, the pain may continue.

3.7 Gynaecological malignancy

The spread of gynaecological malignancy of the cervix, uterine body or ovary will cause pelvic pain depending on the site of spread. Treatment is of the primary condition, but all physicians dealing with pelvic pain must be fully aware of the possibility of gynaecological malignancy.

3.8 Injuries related to childbirth

Tissue trauma and soft tissue injuries occurring at the time of childbirth may lead to CPP related to the site of injury. Dyspareunia is a common problem leading to long-term difficulties with intercourse and female sexual dysfunction (12). Denervation of the pelvic floor with re-innervation may also lead to dysfunction and pain. Hypo-oestrogenism, as a result of breast feeding, may also contribute to pelvic floor pain and dysfunction.

Vulval pain and psychosexual problems are discussed extensively in other sections of this text.

Post-menopausal oestrogen deficiency may lead to pain associated with intercourse, which will respond to hormone replacement therapy.

3.9 Conclusion

Once all the above conditions have been excluded, the gynaecologist may well be left with patients with unexplained pelvic pain. It is imperative to consider pain associated with the urinary and gastrointestinal tract at the same time. For example, patients with bladder pain quite often present with dyspareunia due to bladder base tenderness.

Previously, pelvic congestion has been cited as a course of pelvic pain of unknown aetiology, but this diagnosis is not universally recognised (13, 14).

As previously stated in dealing with pelvic pain, the best results will be obtained from a multidisciplinary approach that considers all possible causes.

3.10 REFERENCES

1. Newham AP, van der Spuy ZM, Nugent F. Laparoscopic findings in women with pelvic pain. *S Afr Med J* 1996;86 (9 Suppl):1200-1203.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9180785&dopt=Abstract
2. Howard FM. The role of laparoscopy as a diagnostic tool in chronic pelvic pain. *Ballieres Best Pract Res Clin Obstet Gynaecol* 2000;14(3):467-494.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10962637&dopt=Abstract&itool=iconabstr
3. Porpora MG, Gomel V. The role of laparoscopy in the management of pelvic pain in women of reproductive age. *Fertil Steril* 1997;68(5):765-779.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9389799&dopt=Abstract
4. Visner SL, Blake RL Jr. Physician's knowledge and treatment of primary dysmenorrhoea. *J Fam Pract* 1985;21(6):462-466.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3934322&dopt=Abstract
5. Porpora MG, Konincks PR, Piazze J, Natili M, Colagrande S, Cosmi EV. Correlation between endometriosis and pelvic pain. *J AM Assoc Gynecol Laparosc* 1999;6(4):429-434.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10548700&dopt=Abstract
6. Ness RB, Soper DE, Holley RL, Peipert J, Randall H, Sweet LL, Sondheimer SJ, Hendrix SL, Amortegui A, Trucco G, Souger T, LA JR, Hillier SL, Bass DC, Kelsey K. Effectiveness of inpatient and outpatient strategies for women with pelvic inflammatory disease. *AM J Obstet Gynecol* 2002;186(5):929-937.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=12015517&dopt=Abstract

7. Corey L, Adams HC, Brown ZA, Holmes KK. Genital herpes simplex infections: clinical manifestations cause and complications. *Ann Intern Med* 1983;98(6):958-972.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6344712&dopt=Abstract
8. Robertson DH, McMillan A, Young H. In: *Clinical practice in sexually transmissible disease*. Edinburgh: Churchill Livingstone, 1989: 333.
9. Fauconnier A, Chapron C, Dubuisson JB, Viera M, Doussett B, Breart G. Relation between pain symptoms and the anatomic location of deep infiltrating endometriosis. *Fertil Steril* 2002;78(4):719-726.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=12372446&dopt=Abstract
10. Goldstein DP, De Cholonky C, Emans SJ. Adolescent endometriosis. *J Adolesc Health Care* 1980;1(1):37-41.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6458589&dopt=Abstract
11. Redwine DB, Wright JT. Laparoscopic treatment of complete obliteration of the cul-de-sac associated with endometriosis: long-term follow-up of en bloc resection. *Fertil Steril* 2001;76(2):358-365.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11476786&dopt=Abstract
12. Osborne JL. Presentation to the European Society of Female Urology. Verona, Italy, Oct 2001.
13. Beard RW, Kennedy RG, Gangar KF, Stones RW, Rogers V, Reginald PW, Anderson M. Bilateral oophorectomy and hysterectomy in the treatment of intractable pelvic pain associated with pelvic congestion. *Br J Obstet Gynaecol* 1991;98(10):988-992.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1751445&dopt=Abstract
14. Foong LC, Gamble J, Sutherland IA, Beard RW. Altered peripheral vascular response of women with and without pelvic pain due to congestion. *Br J Obstet Gynaecol* 2000;107(2):157-164.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10688497&dopt=Abstract

4. NEUROLOGICAL ASPECTS

4.1 Physiology of the urogenital system

The fused somatic and autonomic innervation of the pelvic organs reflects the human need to integrate urogenital functions into our social existence. Proper bladder control is essential for everyday life, while sexual behaviour is an activity upon which depend intimate relationships and the continuation of the species. Physiological functioning of the bladder and sexual organs therefore requires intact innervation, which extends from the frontal lobes of the cortex to the distant pelvic plexi.

The integration of the sympathetic, parasympathetic and somatic innervation in both female and male urogenital tracts is complex. Sympathetic innervation arises from the thoracolumbar outflow, while parasympathetic outflow and somatic innervation originate from the sacral segments of the spinal cord. Afferent nerves travel retrogradely with all three innervating systems. Integration of inputs from the different levels of central and peripheral innervation occurs in plexi, from which nerves arise to innervate pelvic organs. The sacral (predominantly somatic) and the pelvic (predominantly parasympathetic) plexus are intimately linked, with sympathetic connections from the superior and inferior hypogastric plexus.

The sacral plexus innervates the perineum, uterus in the female and the penis in the male through the pudendal nerve. The parasympathetic fibres arise from S2 to S4 to synapse with the ganglia in the pelvic plexus, which are located in the adventitia around the bladder base and in the bladder wall, and from which comes visceral innervation to the bladder and the internal genital organs. The superior hypogastric plexus (sympathetic), situated at the sacral promontory, is the origin of the left and right hypogastric nerves.

4.2 Physiology of the bladder

The urinary bladder is a reservoir whose functions are simply storage and emptying. However, these functions are only accomplished by an extensive and complex innervation that co-ordinates the activity of the striated muscles of the pelvic floor and the various effector organs making up the lower urinary tract (LUT) (1).

4.2.1 *Bladder filling*

Most of the time, the bladder is a low-pressure storage system, which accommodates urine entering from the ureters. The walls of the ureters contain smooth muscle arranged in spiral, longitudinal and circular bundles. They pass obliquely through the bladder wall thereby preventing reflux of urine into the ureters during a bladder contraction.

Under normal circumstances, urine entering the bladder does not cause an increase in intravesical pressure. The smooth muscle of the bladder wall (the detrusor muscle) exhibits plasticity when stretched. The relationship between detrusor pressure and bladder filling can be studied by performing subtracted cystometry, where the pressure difference between two fluid-filled catheters inserted into the bladder and the rectum is determined. A plot of these values against the volume of fluid infused is seen in Figure 6. In health, the detrusor pressure remains almost flat, as fluid fills the bladder that can normally accommodate around 500mL.

Detrusor muscle fibres condense in the region of the bladder neck, forming a well-defined circular collar in the male, and an obliquely/ longitudinally orientated muscle coalescing into the urethral wall in the female. The bladder neck therefore forms a proximal sphincter, which is more evident in the male than in the female, and which is thought to be important in preventing retrograde ejaculation.

Striated muscle comprises the external urethral sphincter, which forms a U-shape around the urethra with some fibres completely encircling it anteriorly, so that as the muscle contracts, the urethra becomes occluded. During the storage phase, continence is maintained by the high resistance offered by the bladder neck and urethra, together with the integrity of the external urethral sphincter (Figure 7). Tonic firing of pudendal motor units of the external urethral sphincter and pelvic floor ensure that a higher pressure is maintained within the urethra than within the bladder. Furthermore, dampening of the parasympathetic innervation of the detrusor by the activation of sympathetic efferents via a sacral to thoracolumbar intersegmental reflex pathway (2) prevents the bladder from contracting spontaneously or involuntarily, the so-called 'storage reflex'. This reflex not only inhibits the detrusor, but also causes contraction of the bladder neck and the proximal urethra. The situation is reversed when micturition is initiated (see below).

4.2.2 *Afferent innervation of the bladder*

Sensations of bladder fullness are conveyed to the spinal cord in the pelvic and hypogastric nerves (3). The afferent components of these nerves contain myelinated (A δ) and unmyelinated (C) axons. The A δ fibres respond to passive distension and active contraction (3) and thus convey information about bladder filling. The C-fibres, insensitive to bladder filling under physiological conditions (therefore termed 'silent' C-fibres) respond primarily to noxious stimuli such as chemical irritation of the urothelium (4) or cooling (5). The cell bodies of both these classes of axons are located in the dorsal root ganglia (DRG) at the level of S2 – S3 and T11 – L2 spinal segments. Bladder afferent activity enters the spinal cord through the dorsal horn and ascends rostrally to higher brain centres involved in bladder control (see below).

In the urinary bladder, sensory nerves have been identified in the suburothelial layer (predominantly in the bladder neck) as well as in the detrusor muscle (6-9). They form a plexus in the suburothelium with some terminal fibres possibly projecting into the urothelium (10-12). Afferent fibres also originate from the trigone and urethra and run in the hypogastric and pudendal nerves respectively.

The response of the bladder to stretch has been extensively investigated and recently a population of cells located in the suburothelial layer of the bladder, called myofibroblasts (13), have been identified. They may act as stretch sensory receptors.

4.2.3 *Efferent innervation of the bladder*

The LUT receives innervation from both the parasympathetic and the sympathetic branches of the autonomic nervous system. The pelvic nerves (arising from the parasympathetic pelvic plexus) cause contraction of the detrusor which effects bladder emptying, whereas parasympathetic innervation of the outflow tract exerts an inhibitory effect resulting in relaxation of the bladder neck and urethra (14). The sympathetic fibres are derived from the T11-T12 and L1-L2 in the spinal cord and run through either the inferior mesenteric ganglia or the hypogastric nerve, or pass through the paravertebral chain to enter the pelvic nerves at the base of the bladder and the urethra. The predominant effect of the sympathetic innervation is inhibition of the parasympathetic pathways at local or spinal level and mediation of contraction of the outflow tract.

The somatic nerve supply to the pelvic floor musculature and part of the urethra originates at S2 – S4 and is conveyed peripherally by the pudendal nerves. A distinct, medially placed motor nucleus at the same spinal level (Onuf's nucleus) supplies axons that innervate the external urethral sphincter.

4.2.4 *Central control of micturition*

Efficient storage and emptying of the bladder requires co-ordinated action of the detrusor and outflow tract. Storage control is achieved in infancy, but voiding is determined by the perceived state of bladder fullness and the social environment (15). The spinal reflexes involved in storage and micturition are relatively simple and are

controlled by higher brain centres. Functional brain imaging has shown that a wide complex of brain networks control the processes of bladder storage (16,17) and voiding (18,19), which ultimately results in the activation or inhibition of the pontine micturition centre.

It is from here that direct pathways descend to the sacral spinal cord and modulate the parasympathetic outflow to the detrusor and co-ordinate the somatic innervation to the external urethral sphincter (20). A desire to void is generated when the bladder volume reaches capacity (approximately 500 mL in humans) (21), but a micturition reflex is only triggered if higher cortical function assesses the situation as appropriate for voiding. Complete emptying is ensured by the resultant detrusor contraction, which is maintained throughout voiding (achieved by the detrusor's unique ability to sustain near-maximal force generation in the face of significant length change [22]), and concomitant relaxation of the outflow tract.

4.2.5 *Physiology of the genital organs*

The female reproductive organs are comprised of the vagina and vulva, important for sexual function, and the ovaries and uterus, necessary for ovulation and reproduction. The uterus is made up of a fibromuscular lower body or cervix and a muscular upper body, which is lined by a hormonally sensitive endometrial layer. The latter responds to the complex monthly hormonal cycle mediated by the hypothalamic-pituitary-ovarian axis acting in tandem with neurological control.

Innervation. Female reproductive organs are innervated in a topographic fashion by afferents which pass retrogradely to the pelvic or hypogastric plexus (23). The afferent nerves contribute to uterine and vaginal perceptions (nociception) that are modified by the reproductive status (24). These plexi communicate with the higher brain centres (the hypothalamus (25), the hippocampus and the limbic system) via the spinal cord, dorsal column nuclei and the solitary nucleus.

The vagina, a highly expandable fibromuscular tube, receives sensory fibres from the pudendal nerve (the perineal and posterior labial branches) and the ilioinguinal nerve. The blood vessels of the smooth muscle of the vaginal walls are supplied by autonomic fibres from the inferior hypogastric plexuses. The clitoris, which is considered homologous to the penis, is also composed of erectile tissue with two miniature corpora cavernosa. Covered with a prepuce, the free end of the clitoris, the glans, is highly sensitive to sexual stimulation. Sexual excitement induces vascular smooth muscle relaxation, (26) mediated by substances such as vasoactive intestinal peptide (VIP) (27) and nitric oxide, and resulting in increased pelvic blood flow, clitoral and labial engorgement and transudative vaginal lubrication. This sexual response is due, as in the male, to parasympathetic activity, and at orgasm, there is repeated contraction of the perineal skeletal muscle, supplied by the perineal branch of the pudendal nerve.

Women with complete spinal cord injury at the mid-thoracic level show perceptual responses to vaginal and/or cervical self-stimulation (e.g. pain suppression and sexual response, including orgasm), with increased activity in the solitary tract nucleus (28).

Menstruation. Animal studies have shown that the rat uterus is directly innervated by both autonomic and sensory nerves, including adrenergic (29) and cholinergic (30), as well as by different peptidergic fibres containing VIP, substance P, calcitonin-gene-related peptide and galanin (31,32). Uterine innervation undergoes profound remodelling during puberty, pregnancy and after delivery. However, the extent to which uterine innervation may change during the menstrual cycle is uncertain (33,34).

The human uterus is under direct control of the hormonal cycle, which influences the innervation of the uterine arteries (cholinergic, adrenergic and peptidergic), and regulates the spontaneous contractile activity of the smooth muscle of vessel walls, as well as the motor responses of these tissues to different vasoactive substances (35). A hormonal disturbance may cause dysfunctional bleeding by changing vessel growth as well as vascular and myometrial smooth muscle activity (36). An example of the latter is primary dysmenorrhoea, when there is an increased secretion of vasopressin (35), which acts on type V1 vasopressin receptors of the uterus, causing myometrial hyperactivity and vasoconstriction, with resultant uterine ischemia and pain.

Pregnancy and parturition. In the pregnant uterus, the motor and sensory innervation undergoes a profound denervation process, although the changes do not affect all types of nerves. Immunocytochemical studies have indicated that myometrial and perivascular VIP-containing fibres disappear at the end of pregnancy (37). In contrast, SP-containing primary afferent neurons do not degenerate during pregnancy (38). At the end of pregnancy, the numbers of both myometrial and perivascular adrenergic nerves are decreased in the rat (39,40) in the guinea pig (41,42), and in humans (43). The whole autonomic uterine innervation therefore undergoes substantial remodelling during pregnancy.

Urogenital pain. The innervation of the pelvis shows great convergence, demonstrating the existence of extensive cross-system, viscero-visceral interactions within the CNS, which, while organized for coherent

bodily functioning, serve as a substrate by which pathophysiology in one organ can influence physiology and responses to pathophysiology in other organs (24). Some cross-system effects reported in the literature include bladder inflammation, reducing the rate of uterine contractions and the effects of drugs on the uterus (44) and colon inflammation, producing signs of inflammation in an otherwise healthy bladder and uterus. How the pathophysiology of one pelvic organ influences the physiology of another is poorly understood, but improved knowledge and understanding of the convergence of peripheral and central innervation of the pelvis may have considerable clinical relevance.

Conclusion. The rich and complex innervation of the bladder and genital organs, which enables their physiological function, appears also to be a potential substrate for neural pain. Perhaps pain results when autonomic innervation becomes 'confused'.

Figure 6: Effector organs making up the lower urinary tract (LUT). This is a representation of the female LUT. In the male, the prostate gland is situated around the external urethral sphincter. (With permission of Mr. Vinal Kalsi).





Figure 7: Subtraction cystometry showing typical bladder function.



Pabd = intra-abdominal pressure measured with a rectal catheter; *Pves* = measured intravesical pressure; *Pdet* = detrusor pressure; *Pdet* = *Pves*-*Pabd*; *Vinfus* = infusion volume, 50 mL/min.

The detrusor pressure remains less than 10 cmH₂O until the first desire to void (FD) is reached, which is accompanied by a small rise in the pressure tracing. There is a concomitant rise in vesical pressure; however, there is no rise in intra-abdominal pressure. The pressure rise is due to contraction of the detrusor; however, it is small due to the compliant properties of the bladder. The normal desire to void (ND) is soon accompanied by cystometric capacity (about 500mL), at which time a void will be initiated. The undulations seen on the traces are a result of interference due to respiration.

4.3 Sexual dysfunction in men and women

Various classifications of sexual dysfunction have been proposed, the most recent being one which included hypoactive sexual desire, or disorders of sexual desire, sexual arousal, orgasms or sexual pain (45). Many of these disorders are common amongst the general population: the Male Massachusetts study showing an increasing prevalence of ED with age so that of a group of men aged 60-70 almost 60% had ED to a greater or lesser extent (46). The prevalence of FSD has been estimated to be between 25%-63%, the figure depending on the definition used and population studies. Amongst groups of patients with neurological disease the prevalence of all types of disorder is even higher, although precise figures are not known.

Neurological causes of sexual dysfunction include cortical disease, spinal cord trauma, stroke, epilepsy, multiple sclerosis, radical pelvic surgery and many more conditions. As the aetiology is diverse, so is the pathophysiology with damage to the thoraco-lumbar outflow in men during extensive surgery affecting ejaculation to damage of the hypothalamus and pituitary, following head injury, resulting in hypopituitarism and a concomitant de-sensitisation of the genital region.

The treatment in men and women with neurological or non-neurological disease includes pelvic floor exercises and electrical stimulation feedback with cognitive therapy. Male sexual dysfunction is discussed in detail within these guidelines.

Female sexual dysfunction is less easy to treat, but is affected by problems in the male, and it is now recommended that evaluation of the female should be addressed within the context of the couple in a sexual medicine clinic. Identifying and treating general medical conditions is vital to the effective management of both men and women. In addition to neurological related dysfunction in young women, pelvic floor dysfunction, as a

consequence of childbirth, must be taken into account as the importance of the menopause in the older woman. Despite the fact that hormone replacement therapy has been used extensively and effectively, there is still a small sub-set of women in whom this is not enough. In this group the libido is affected greatly and the use of testosterone products have been found to be important. Pharmacotherapy, like the PDE-5 inhibitors in men, has not been found to be useful in women.

4.4 References

1. de Groat WC, Booth AM. Physiology of the urinary bladder and urethra. *Ann Intern Med* 1980;92(2 Pt 2):312-315.
http://www.ncbi.nlm.nih.gov/pubmed/6243894?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
2. de Groat WC, Lalley PM. Reflex firing in the lumbar sympathetic outflow to activation of vesical afferent fibres. *J Physiol* 1972;226(2):289-309.
http://www.ncbi.nlm.nih.gov/pubmed/4508051?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
3. Janig W and Morrison, JF, Functional properties of spinal visceral afferents supplying abdominal and pelvic organs, with special emphasis on visceral nociception. *Prog Brain Res* 1986;67:87-114.
http://www.ncbi.nlm.nih.gov/pubmed/3823484?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
4. Habler HJ, Janig W, Koltzenburg M. Activation of unmyelinated afferent fibres by mechanical stimuli and inflammation of the urinary bladder in the cat. *J Physiol* 1990;425:545-562.
http://www.ncbi.nlm.nih.gov/pubmed/2213588?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
5. Fall M, Lindstrom S, Mazieres L. A bladder-to-bladder cooling reflex in the cat. *J Physiol* 1990;427:281-300.
http://www.ncbi.nlm.nih.gov/pubmed/2213600?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
6. Gosling JA, Dixon JS. Sensory nerves in the mammalian urinary tract. An evaluation using light and electron microscopy. *J Anat* 1974;117(Pt 1):133-144.
http://www.ncbi.nlm.nih.gov/pubmed/4844655?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
7. Smet PJ, Moore KH, Jonavicius J. Distribution and colocalization of calcitonin gene-related peptide, tachykinins, and vasoactive intestinal peptide in normal and idiopathic unstable human urinary bladder. *Lab Invest* 1997;77(1):37-49.
http://www.ncbi.nlm.nih.gov/pubmed/9251677?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
8. Avelino A, Cruz C, Nagy I, Cruz F. Vanilloid receptor 1 expression in the rat urinary tract. *Neuroscience* 2002;109(4):787-798.
http://www.ncbi.nlm.nih.gov/pubmed/11927161?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
9. Yiangou Y, Facer P, Ford A, Brady C, Wiseman O, Fowler CJ, Anand P. Capsaicin receptor VR1 and ATP-gated ion channel P2X3 in human urinary bladder. *BJU Int* 2001;87(9):774-779.
http://www.ncbi.nlm.nih.gov/pubmed/11412212?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
10. Birder LA, Kanai AJ, de Groat WC, Kiss S, Nealen ML, Burke NE, Dineley KE, Watkins S, Reynolds IJ, Caterina MJ, Vanilloid receptor expression suggests a sensory role for urinary bladder epithelial cells. *Proc Natl Acad Sci USA* 2001;98(23):13396-13401.
http://www.ncbi.nlm.nih.gov/pubmed/11606761?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
11. Lazzeri M, Vannucchi G, Zardo C, Spinelli M, Beneforti P, Turini D, Faussone-Pellegrini MS. Immunohistochemical evidence of vanilloid receptor 1 in normal human urinary bladder urothelium. *Eur Urol* 2004;46(6):792-799.
http://www.ncbi.nlm.nih.gov/pubmed/15548449?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
12. Gabella G, Davis C. Distribution of afferent axons in the bladders of rats. *J Neurocytol* 1998;27(3):141-155.
http://www.ncbi.nlm.nih.gov/pubmed/10640174?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

13. Wiseman OJ, Fowler CJ, Landon DN. The role of the human bladder lamina propria myofibroblast. *BJU Int* 2003;91(1):89-93.
http://www.ncbi.nlm.nih.gov/pubmed/12614258?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
14. Burnstock G. Innervation of bladder and bowel. *Ciba Found Symp* 1990;151:2-18; discussion 18-26.
http://www.ncbi.nlm.nih.gov/pubmed/1977565?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
15. Kavia RB, Dasgupta R, Fowler CJ. Functional imaging and the central control of the bladder. *J Comp Neurol* 2005;493(1):27-32.
http://www.ncbi.nlm.nih.gov/pubmed/16255006?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
16. Athwal BS, Berkley KJ, Hussain I, Brennan, A, Craggs M, Sakakibara R, Frackowiak RS, Fowler CJ. Brain responses to changes in bladder volume and urge to void in healthy men. *Brain* 2001;124(Pt 2): 369-377.
http://www.ncbi.nlm.nih.gov/pubmed/11157564?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
17. Matsuura S, Kakizaki H, Mitsui T, Shiga T, Tamaki N, Koyanagi T. Human brain region response to distention or cold stimulation of the bladder: a positron emission tomography study. *J Urol* 2002;168(5):2035-2039.
http://www.ncbi.nlm.nih.gov/pubmed/12394703?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
18. Blok BF, Willemssen AT, Holstege G. A PET study on brain control of micturition in humans. *Brain* 1997;120(Pt 1):111-121.
http://www.ncbi.nlm.nih.gov/pubmed/9055802?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
19. Blok BF, Sturms LM, Holstege G. Brain activation during micturition in women. *Brain* 1998;121(Pt 11):2033-2042.
http://www.ncbi.nlm.nih.gov/pubmed/9827764?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
20. Blok BF. Central pathways controlling micturition and urinary continence. *Urology* 2002;59(5 Suppl 1):13-7.
http://www.ncbi.nlm.nih.gov/pubmed/12007517?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
21. Wyndaele JJ. Normality in urodynamics studied in healthy adults. *J Urol* 1999;161(3):899-902.
http://www.ncbi.nlm.nih.gov/pubmed/10022710?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
22. Uvelius B, Gabella G. Relation between cell length and force production in urinary bladder smooth muscle. *Acta Physiol Scand* 1980;110(4):357-365.
http://www.ncbi.nlm.nih.gov/pubmed/7234441?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
23. Anaf V, Simon P, El Nakadi I, Fayt I, Buxant F, Simonart T, Peny MO, Noel JC. Relationship between endometriotic foci and nerves in rectovaginal endometriotic nodules. *Hum Reprod* 2000;15(8):1744-1750.
http://www.ncbi.nlm.nih.gov/pubmed/10920097?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
24. Berkley KJ. A life of pelvic pain. *Physiol Behav* 2005;86(3):272-280.
http://www.ncbi.nlm.nih.gov/pubmed/16139851?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
25. Akaishi T, Robbins A, Sakuma Y, Sato Y. Neural inputs from the uterus to the paraventricular magnocellular neurons in the rat. *Neurosci Lett* 1988;84(1):57-62.
http://www.ncbi.nlm.nih.gov/pubmed/3347371?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
26. Berman JR, Adhikari SP, Goldstein I. Anatomy and physiology of female sexual function and dysfunction: classification, evaluation and treatment options. *Eur Urol* 2000;38(1):20-29.
http://www.ncbi.nlm.nih.gov/pubmed/10859437?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
27. Levin RJ. VIP, vagina, clitoral and periurethral glans—an update on human female genital arousal. *Exp Clin Endocrinol* 1991;98(2):61-69.
http://www.ncbi.nlm.nih.gov/pubmed/1778234?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

28. Whipple B, Komisaruk BR. Brain (PET) responses to vaginal-cervical self-stimulation in women with complete spinal cord injury: preliminary findings. *J Sex Marital Ther* 2002;28(1):79-86.
http://www.ncbi.nlm.nih.gov/pubmed/11928182?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
29. Sjoberg NO. Dysmenorrhea and uterine neurotransmitters. *Acta Obstet Gynecol Scand* 1979;87:57-59.
http://www.ncbi.nlm.nih.gov/pubmed/37691?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
30. Stjernquist M, Owman CO. Cholinergic and adrenergic neural control of smooth muscle function in the non-pregnant rat uterine cervix. *Acta Physiol Scand* 1985;124(3):429-436.
http://www.ncbi.nlm.nih.gov/pubmed/4050475?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
31. Papka RE, Cotton JP, Traurig HH. Comparative distribution of neuropeptide tyrosine-, vasoactive intestinal polypeptide-, substance P-immunoreactive, acetylcholinesterase-positive and noradrenergic nerves in the reproductive tract of the female rat. *Cell Tissue Res* 1985 242(3):475-490.
http://www.ncbi.nlm.nih.gov/pubmed/2416449?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVAbstractPlusDrugs1
32. Shew RL, Papka RE, McNeill DL. Galanin and calcitonin gene-related peptide immunoreactivity in nerves of the rat uterus: localization, colocalization, and effects on uterine contractility. *Peptides* 1992;13(2):273-279.
http://www.ncbi.nlm.nih.gov/pubmed/1384006?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVAbstractPlusDrugs1
33. Sjoberg NO. New considerations on the adrenergic innervation of the cervix and uterus. *Acta Obstet Gynecol Scand* 1969;48 (Suppl 3):28+.
http://www.ncbi.nlm.nih.gov/pubmed/5380821?ordinalpos=22&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
34. Zoubina EV, Fan Q, Smith PG. Variations in uterine innervation during the estrous cycle in rat. *J Comp Neurol* 1998;397(4):561-571.
http://www.ncbi.nlm.nih.gov/pubmed/9699916?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVAbstractPlusDrugs1
35. Akerlund M. Vascularization of human endometrium. Uterine blood flow in healthy condition and in primary dysmenorrhoea. *Ann N Y Acad Sci* 1994;734:47-56.
http://www.ncbi.nlm.nih.gov/pubmed/7978951?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
36. Proctor ML, Latthe PM, Farquhar CM, Khan KS, Johnson NP. Surgical interruption of pelvic nerve pathways for primary and secondary dysmenorrhoea. *Cochrane Database Syst Rev* 2005(4):CD001896.
http://www.ncbi.nlm.nih.gov/pubmed/16235288?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVAbstractPlusDrugs1
37. Stjernquist M, Alm P, Ekman R, Owman C, Sjoberg NO, Sundler F. Levels of neural vasoactive intestinal polypeptide in rat uterus are markedly changed in association with pregnancy as shown by immunocytochemistry and radioimmunoassay. *Biol Reprod* 1985;33(1):157-163.
http://www.ncbi.nlm.nih.gov/pubmed/4063437?ordinalpos=8&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
38. Traurig H, Saria A, Lembeck F. Substance P in primary afferent neurons of the female rat reproductive system. *Naunyn Schmiedebergs Arch Pharmacol* 1984;326(4):343-346.
http://www.ncbi.nlm.nih.gov/pubmed/6207443?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
39. Moustafa, FA. Changes in cholinergic and noradrenergic nerves in the pregnant and postpartum uterus of the albino rat and guinea pig. *Acta Anat (Basel)* 1988;132(4):310-316.
http://www.ncbi.nlm.nih.gov/pubmed/3195314?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
40. Haase, EB, Buchman J, Tietz AE, Schramm LP. Pregnancy-induced uterine neuronal degeneration in the rat. *Cell Tissue Res* 1997;288(2):293-306.
http://www.ncbi.nlm.nih.gov/pubmed/9082965?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

41. Thorbert G, Alm P, Owman C, Sjöberg NO, Spörng B. Regional changes in structural and functional integrity of myometrial adrenergic nerves in pregnant guinea-pig, and their relationship to the localization of the conceptus. *Acta Physiol Scand* 1978;103(2):120-131.
http://www.ncbi.nlm.nih.gov/pubmed/676764?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
42. Alm P, Owman C, Sjöberg NO, Thorbert G. Uptake and metabolism of [3H]norepinephrine in uterine nerves of pregnant guinea pig. *Am J Physiol* 1979;236(5):C277-C285.
http://www.ncbi.nlm.nih.gov/pubmed/443367?ordinalpos=7&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
43. Wikland M, Lindblom B, Dahlström A, Haglid KG. Structural and functional evidence for the denervation of human myometrium during pregnancy. *Obstet Gynecol* 1984;64(4):503-509.
http://www.ncbi.nlm.nih.gov/pubmed/6384845?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
44. Dmitrieva N, Johnson OL, Berkley KJ. Bladder inflammation and hypogastric neurectomy influence uterine motility in the rat. *Neurosci Lett* 2001;313(1-2):49-52.
http://www.ncbi.nlm.nih.gov/pubmed/11684337?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
45. Levin RJ. The physiology of sexual arousal in the human female: a recreational and procreational synthesis. *Arch Sex Behav* 2002;31(5):405-411.
http://www.ncbi.nlm.nih.gov/pubmed/12238607?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
46. Feldman HA, Goldstein I, Hatzichristou DG, Krane RJ, McKinlay JB. Impotence and its medical and psychosocial correlates: results of the Massachusetts Male Ageing Study. *J Urol* 1994;151(1):54-61.

5 NEUROGENIC CONDITIONS

5.1 Introduction

It is clearly important for the patient to have been thoroughly examined by a urologist or gynaecologist and local pelvic pathology excluded. Once a structural cause has been eliminated, a neurological opinion is often sought, with the prime aim of the neurologist being to exclude any form of conus or sacral root pathology. MRI is the investigation of choice to show both neural tissue and surrounding structures.

If all examinations and investigations fail to reveal an abnormality, the diagnosis is likely to be one of the focal pain syndromes. These are persistent or recurrent or episodic pains referred to specific pelvic organs in the proven absence of infection, malignancy or other obvious pathology (see Table 1). Although these are well-recognized conditions, their pathophysiology is not understood. However, it seems likely that the problems relate in some way to the combined visceral, autonomic and somatic innervation of the pelvic organs.

5.2 Pudendal nerve entrapment

Chronic compression of the pudendal nerve in the ischiorectal fossa may result in a perineal pain located either anteriorly in the vagina or vulval region, or posteriorly in the anorectal region. The ICS has used the following definition, 'perineal pain is felt: in the female, between the posterior fourchette (posterior lip of the introitus) and the anus, and in the male, between the scrotum and the anus' (1).

The pain may include unpleasant sensations of numbness or a burning sensation, and may be exacerbated by sitting and relieved by standing. Neurological examination of the perineum is normal. If tested, the sacral reflexes are present and anal sphincter tone is normal. Neurophysiological examination is said to be helpful in some cases; sacral reflex latency (using electrical stimulation of the dorsal nerve of the clitoris and recording muscle activity in the perineum) and the pudendal nerve distal motor latency using the St Marks Stimulator has been recommended. These investigations require specialist neurophysiological expertise.

Pudendal nerve neuropathy is likely to be a probable diagnosis if the pain is unilateral, has a burning quality and is exacerbated by unilateral rectal palpation of the ischial spine, with delayed pudendal motor latency on that side only. However, such cases account for only a small proportion of all those presenting with perineal pain. Proof of diagnosis rests on pain relief following decompression of the nerve in Alcock's canal and is rarely achieved. The value of the clinical neurophysiological investigations is debatable; some centres in Europe claim that the investigations have great sensitivity (1,2), while other centres, which also have a specialized interest in pelvic floor neurophysiology, have not identified any cases. Further information may be gained by a diagnostic nerve block or MRI investigation.

5.3 Other neurogenic conditions

Other pelvic floor clinical neurophysiological investigations are more helpful in identifying changes of denervation and re-innervation. Lesions causing such disorders are usually associated with bladder and/or sexual dysfunction rather than isolated urogenital pain.

A major defect of the clinical neurophysiological investigations currently available is that they examine mostly large myelinated nerve fibre function rather than the unmyelinated and small myelinated fibres, which subserve autonomic innervation, pelvic organ sensation and pain (3).

5.4 References

1. Amarenco G, Kerdraon J. Pudendal nerve terminal sensitive latency: technique and normal values. *J Urol* 1999;161(1):103-106.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10037379&dopt=Abstract
2. Robert R, Prat-Pradal D, Labat JJ, Bensignor M, Raoul S, Rebai R, Leborgne J. Anatomic basis of chronic perineal pain: role of the pudendal nerve. *Surg Radiol Anat* 1998;20(2):93-98.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9658526&dopt=Abstract
3. Lee JC, Yang CC, Kromm BG, Berger RE. Neurophysiologic testing in chronic pelvic pain syndrome: a pilot study. *Urology* 2001;58(2):246-250.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11489711&dopt=Abstract

6. PELVIC FLOOR FUNCTION AND DYSFUNCTION

6.1 Introduction

The pelvic floor is made up of muscles and fascia. The muscles usually function as a composite, though the anterior and posterior components may act in isolation. The pelvic floor has three functions: support, contraction and relaxation.

6.2 Function

In its resting state, the pelvic floor supports the bladder and the urethra in the anterior compartment, the uterus and the vagina in the middle compartment, and the rectum and the anus in the posterior compartment. The integrity of the support function depends on the anatomical position of the muscles, on the resting 'tone' and on the integrity of the fascia (1). As with all skeletal muscles, tone is maintained by the efferent nerve fibres, and may vary with hormonal status (menstrual cycle, pregnancy, and menopause).

The support activated during a rise in intra-abdominal pressure is different from that at rest. When intra-abdominal pressure rises, the pelvic floor muscles must respond with a contraction occurring simultaneously or before the pressure rise. The latter is termed an anticipatory 'response' or feed-forward loop (2). Electromyography recordings show tonic motor unit activity at rest, with phasic recruitment of large motor units in response to coughing.

A contraction of the pelvic floor muscles results in an inward movement of the perineum and an upward movement of the pelvic organs. In many situations, other muscles such as the abdominal muscles, the adductor muscles and the gluteal muscles also contract. There are two types of contraction that can be distinguished: a voluntary contraction, resulting from impulses arising in the cerebral cortex, and a reflex contraction. These contractions not only maintain support of the pelvic organs, they close the urethra, anus and vagina, thus avoiding loss of urine or stool, and affording women a defensive mechanism. Additionally, detrusor inhibition occurs in parallel with pelvic floor muscle contraction.

A contraction of the pelvic floor muscles must have sufficient strength. Strength results from muscle capacity and neurogenic drive, reflected in the frequency of excitation and the number of activated motor units. An increase in muscle strength is achieved through the recruitment of more motor units. A contraction must be rapidly effective and remain so for a certain period (endurance).

Pelvic floor contractions play an important role in sexual function. During the arousal phase, pelvic floor muscle contractions are used to increase vasocongestion. During the last phase of the sexual response cycle, a series of involuntary contractions is associated with the physical sensations of orgasm (3).

Pelvic floor muscle relaxation results in a decrease or termination of the squeezing of the urethra, vagina and anus. The perineum and the pelvic organs return to their anatomical resting position. Pelvic floor muscle relaxation is the result of inhibition of tonically active motor units. Relaxation of the pelvic floor muscles

is needed for voiding, defaecation and for sexual intercourse.

6.3 Dysfunction

Pelvic floor dysfunction should be classified according to 'The standardisation of terminology of pelvic floor muscle function and dysfunction' (4). This is an international multidisciplinary report from the ICS. As in all ICS standardization documents, this is based on the triad of symptom, sign and condition. Symptoms are what the patient tells you; signs are found by physical examination. By palpation of the pelvic floor muscles, the contraction and relaxation are qualified. The voluntary contraction can be absent, weak, normal or strong. The voluntary relaxation can be absent, partly or completely. The involuntary contraction and relaxation is absent or present.

Based on these signs, pelvic floor muscles can be classified as follows:

- non-contracting pelvic floor
- non-relaxing pelvic floor
- non-contracting, non-relaxing pelvic floor.

Based on symptoms and signs, the following conditions are possible:

- normal pelvic floor muscles
- overactive pelvic floor muscles
- underactive pelvic floor muscles
- non-functioning pelvic floor muscles.

An example is as follows:

- *Symptom*: pain in the perineal region
- *Sign* non-relaxing pelvic floor (no relaxation, neither voluntary nor involuntary)
- *Condition*: overactive pelvic floor muscles.

An underactive pelvic floor means that the muscles do not contract when they need to. In practice, this leads to incontinence of urine or stool. An overactive pelvic floor means that the pelvic floor muscles do not relax when they should. This may result in complaints like low flow rates and constipation (5). Another symptom of overactivity is CPP and more specific dyspareunia.

Overactivity tends to develop over a protracted period, with many causes. In most cases, there is the problem of limited access to a toilet on demand, leading to postponement of voiding by contraction of the pelvic floor muscles. When they do eventually have the time to void, detrusor power is lacking. They start to use abdominal straining which results, through the guarding reflex, in contraction of the pelvic muscles (6).

Why an overactive pelvic floor causes pain has only partly been elucidated (7). A muscle that is continuously contracting will ache. Nerves that pass through the pelvic floor may be compressed, and vessels to the penis and scrotum may be obstructed. Both mechanisms will lead to pelvic pain. A contracting pelvic floor will increase afferent input to the sacral spinal cord, the pons and the cerebral cortex. In response, the CNS may modify efferent signals to the pelvis. This change in efferent activity may further aggravate the situation (8).

6.4 Myofascial trigger points

Repeated or chronic muscular overload can activate trigger points in the muscle. Trigger points are defined as hyperirritable spots associated with a hypersensitive palpable nodule in a taut band (9). Trigger points are painful on compression and give rise to characteristic referred pain and motor dysfunction. They prevent full lengthening of the muscles, they weaken the muscle and they lead to pain that is very recognisable for the patient.

Pain as a result of these trigger points is aggravated by specific movements and alleviated by certain positions. Patients know what activities and postures influence the pain. Trigger points can be located within the pelvic floor muscle (10). In a case of pelvic floor muscle trigger points, a patient will sit down cautiously, often on one buttock. Rising after a period of sitting will cause pain. Pain will be aggravated by pressure on the trigger point (e.g. pain related to sexual intercourse). Pain will also get worse after sustained or repeated contractions (e.g. pain related to voiding or defecation). On physical examination, trigger points can be palpated and compression will give local and referred pain. In patients with CPP, trigger points are often found in muscles related to the pelvis like abdominal, gluteal and piriformis muscle.

6.5 Therapy

Treating pelvic floor overactivity should be considered in the management of CPP (11). There are a number of methods, taught by specialized physiotherapists, which can be used to improve the function and co-ordination of the pelvic floor muscles. The use of biofeedback by means of pelvic floor muscle electromyography should be considered because it might help the patient to understand the dysfunction of the pelvic floor muscles. This

understanding will improve the result of the treatment.

Central trigger points are treated by stretching the muscle, which inactivates them. However, trigger points lying in the attachment of the muscle to the bone respond better to direct manual therapy. Muscle exercises are helpful, e.g. voluntary contractions followed by complete relaxation. Pressure on the trigger points and subsequent release is also effective (12,13). Stretching of the muscle will be more effective after pain relief by direct pressure on the trigger point. Injecting the trigger points with a local anaesthetic will show that the trigger points are really causing the pain; its will give an acute relief of pain and will unblock the muscle so that stretching becomes possible.

6.6 References

1. Olsen AL, Smith VJ, Bergstrom JO, Colling JC, Clark AL. Epidemiology of surgically managed pelvic organ prolapse and urinary incontinence. *Obstet Gynecol* 1997;89(4):501-506.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9083302&dopt=Abstract
2. Constantinou CE, Govan DE. Spatial distribution and timing of transmitted and reflexly generated urethral pressures in healthy women. *J Urol* 1982;127(5):964-969.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7201031&dopt=Abstract
3. Epstein M. Physiology of sexual function in women. In: Epstein M, ed. *Clinics in obstetrics and gynaecology*. London: WB Saunders, 1980: 7.
4. Messelink EJ, Benson T, Berghmans B et al. Standardisation of terminology of pelvic floor muscle function and dysfunction: report from the pelvic floor clinical assessment group of the International Continence Society. 2005;24:374-380.
5. Kaplan SA, Santarosa RP, D'Alisera PM, Fay BJ, Ikeguchi EF, Hendricks J, Klein L, Te AE. Pseudodyssynergia (contraction of the external sphincter during voiding) misdiagnosed as chronic nonbacterial prostatitis and the role of biofeedback as a therapeutic option. *J Urol* 1997;157(6):2234-2237.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9146624&dopt=Abstract
6. Messelink EJ. The overactive bladder and the role of the pelvic floor muscles. *BJU Int* 1999;83(Suppl 2):31-35.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10210602&dopt=Abstract
7. Howard FM. Pelvic floor pain syndrome. In: Howard FM, ed. *Pelvic Pain. Diagnosis and Management*. Philadelphia: Lippincott Williams & Wilkins, 2000: 429-432.
8. Zermann DH, Ishigooka M, Doggweiler R, Schmidt RA. Chronic prostatitis: a myofascial pain syndrome? *Infect Urol* 1999;12:84-86.
9. Carter J.E. Abdominal wall and pelvic myofascial trigger points. In: Howard FM, ed. *Pelvic Pain. Diagnosis and Management*. Lippincott Williams & Wilkins: Philadelphia, 2000: 314-358
10. Slocumb JC. Neurological factors in chronic pelvic pain: trigger points and the abdominal pelvic pain syndrome. *Am J Obstet Gynaecol* 1984;149(5):536-543.
http://www.ncbi.nlm.nih.gov/pubmed/6234807?ordinalpos=9&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
11. Glazer HI, Rodke G, Swencionis C, Hertz R, Young AW. Treatment of vulvar vestibulitis syndrome with electromyographic biofeedback of pelvic floor musculature. *J Reprod Med* 1995;40:283-290.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7623358&dopt=Abstract
12. Anderson RU, Wise D, Sawyer T, Chan C. Integration of myofascial trigger point release and paradoxical relaxation training treatment of chronic pelvic pain in men. *J Urol* 2005;174(1):155-160.
http://www.ncbi.nlm.nih.gov/pubmed/15947608?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
13. Srinivasan AK, Kaye JD, Moldwin R. Myofascial dysfunction associated with chronic pelvic floor pain: management strategies. *Curr Pain Headache Rep* 2007;11(5):359-364.
http://www.ncbi.nlm.nih.gov/pubmed/17894926?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

7. PSYCHOLOGICAL FACTORS IN PERSISTENT CHRONIC PELVIC PAIN

7.1 Introduction

This section describes the evidence for psychological factors, which play a part in the development or maintenance of persistent pelvic pain, in adjustment to pain, and in treatments including psychologically-based treatments.

The function of pain, particularly acute pain, is to demand action to stop or minimize damage, and to protect the injured part to promote healing. When pain persists without ongoing damage or disease, or after the damage has healed, it appears to have no such function. However, it can disrupt daily life and cause distress, particularly since the person with pain tends to use the acute pain model, and to fear damage, disease, and prolonged suffering. Cognitive and emotional processes are integral in pain processing (1,2). Psychological and social factors are involved as risk factors for the development of persistent pain from acute pain, in adaptation to pain, and in the decision to seek and to adhere (or not) to recommended treatment for the pain.

7.2 Psychological models of pain

A fully integrated model of pain, as described above, with changes in excitatory and inhibitory mechanisms at peripheral and central level, and changes in cortical and subcortical processing of pain, but not defined by tissue damage or disease severity, does not correspond well with the model of somatization, or somatoform pain.

This model assumes that pain can be partitioned into distinct biological and psychological processes which, added together, constitute the presenting complaint, and that absence of significant physical signs constitutes evidence for substantial psychological causation (3,4). This has been a particularly prominent model in pelvic pain, at least in women.

In fact, multiple models co-exist, often poorly specified. One model, unsubstantiated by psychiatric diagnostic systems, rules that pain is inauthentic if it is not supported by clinical signs and that the pain is therefore a covert expression of distress or psychological disorder. This model relies on a dualism, which has no place in scientific medicine and is unacceptable to most patients. Another model, commoner in psychodynamically informed systems, constructs pain (perhaps without physical signs, but not defined by their absence) as an expression of an unconscious conflict or source of distress.

Several models are compatible with our understanding of physiological processes as it has developed since Melzack and Wall's (1965) work (1). Psychophysiological reactivity, built on stress models, describes the activation of systems for defence against physical or psychological stress. Prolonged stress leads to repeated alerting and depletion of systems involved. This was demonstrated by Flor et al. (1992) (5), using electromyography in back muscles, in response to psychological/social stressors. This may be a mechanism for the production of pain in pelvic floor muscles, even if it is not the primary cause of pelvic pain. Pelvic floor overactivity is now the subject of investigation and treatment attempts (see Sections 2.5 and 6.5). In addition, Link et al. (2007) (6) have proposed an association between trauma (sexual and physical abuse in childhood) and bladder activity and urinary symptoms, presumably mediated by stress mechanisms.

The cognitive-behavioural model classically took pain as a starting point, and described modulation of the experience of pain, and reaction to it, in terms of cognition, emotion and behaviour. There are reliable associations between certain beliefs or types of belief, and associated behaviour and/or emotions, including extensive evidence for the importance of psychological factors in determining the level of pain experienced, levels of disability and restriction due to pain, levels of distress, and extent of healthcare use (2,7), demonstrated either in specific pain problems, such as back pain or fibromyalgia, or for mixed persistent pain populations, but with few studies in pelvic pain populations.

Psychological factors are better researched and described than social factors, and both suffer from overlapping constructs which neither fall into clear hierarchies of organization nor map well on to processes and pathways in the brain. Among the better established ones are attention (8) and fear and associated avoidance patterns (9).

The cognitive and behavioural model is compatible with the psychophysiological reactivity model. More recently, it has been extended by reference to neurotransmitters common to pain or to other symptoms, including urinary symptoms (10) and to mood pathways; and to dysregulation across multiple systems producing a range of 'medically unexplained symptoms', which concomitantly occur with mood disorders at a higher level than chance in individuals and in their first-degree relatives (11,12).

Psychological models, particularly cognitive and behavioural models, when integrated with a biomedical model, are described as biopsychosocial models.

7.3 Methodology

Both PubMed and PsychInfo were searched to access trials published in psychology journals not covered by medical databases, with a search covering the last 10 years, since psychological trials tend to take longer to run and to reach publication than medical and other clinical trials. The Cochrane database was also searched.

7.4 Psychological factors in assessment of pelvic pain

A careful review by Savidge and Slade (1997) (13) of CPP in women identifies as problematic the notion that women without physical findings to which pain can be causally attributed differ in psychological characteristics from women with physical findings. They critically examine the methodologies of studies purporting to show such differences, and the bias introduced by sampling and by unsuitable measures; they argue for better methodology in replication of these studies, particularly those sampling life events, and for greater use of idiographic methods. However, the report of anxiety, depression, and sexual problems is sufficiently common for these to be important in assessment and in planning treatment. Distress, described in the patient's terms or within a psychodiagnostic framework, is best understood in the context of pain and of the meaning of pain to the individual.

7.4.1 Psychological risk factors in the development of pelvic pain and adaptation to it

There is one systematic review (14) of risk factors for CPP in women: there appears to be no equivalent systematic review of pelvic pain in men. For non-cyclical pelvic pain in women, Latthe et al. drew on 40 studies ($n = 20,040$). Of the 48 risk factors included in the studies, pain was associated with (biomedical factors) pelvic pathology, miscarriage, and heavy menstrual flow; and (psychological factors):

- lifetime drug and alcohol abuse (OR 4.61, 95% CI 1.09-19.38)
- sexual or physical abuse (OR 1.51-3.49)
- psychological problems: anxiety (OR 2.28; 95% CI 1.41-3.70); depression (OR 2.69; 95% CI 1.86-3.88); multiple somatic problems (OR 4.83; 95% CI 2.50-9.33 and OR 8.01; 95% CI 5.16-12.44): the terms 'hysteria' and 'psychosomatic symptoms' are used but can best be understood as multiple somatic symptoms not associated with or indicative of any serious disease process.

Personality variables, length of education, and marital status were not reliably associated with pelvic pain in women. Interrelationships, such as between history of sexual abuse and depression, for instance, cannot be disentangled from the studies available.

A meta-analysis (15) confirms the reported association in retrospective studies between the report of childhood sexual abuse by adults and persistent pain; often this concerns childhood sexual abuse and pelvic pain (16). However, these studies are retrospective; interestingly, Latthe et al. (14) found poor quality papers were more likely to report this association than better quality ones. The only prospective investigation into the relationship between childhood sexual abuse, physical abuse, or neglect, and 'medically unexplained pain' including pelvic pain, used court records concerning sexual abuse before the age of 11 to establish a definite history, comparing those with such a history with demographically matched classmates (17). It concluded that physically and sexually abused individuals were not at risk for increased pain symptoms. Although those individuals with pain problems as adults were more likely to report earlier sexual or physical abuse or neglect, this did not correspond with the established early history of abuse. The correlation between childhood victimization and pain symptoms is less straightforward than previously thought, and may be more about retrospective explanatory frameworks used by women for pain which is 'medically unexplained' than about occurrence or extent of abuse.

In particular, findings of depression and/or post-traumatic stress disorder in adult women reporting childhood sexual abuse are common, with or without pain. Disentangling the influences and inferences requires prospective study or suitable comparison groups. See Savidge and Slade (1997) (13) for an excellent critique. Within pelvic pain populations, worthwhile studies can be done without comparison. For example, Poleshuck et al. (2005) (18) found that, in women with pelvic pain attending a clinic, the report of physical or sexual abuse in childhood was associated with greater psychological distress than in women without, but there were no differences in pain experience, or physical or social function. In summary, women with pelvic pain often have other 'medically unexplained' symptoms, and current or a lifetime anxiety and depression disorder; they may have a history of physical or sexual abuse in childhood but the significance of this for pelvic pain is unclear.

The report of pelvic pain appears to be more common after rape (19), and recent sexual assault should be considered, particularly if the woman expresses distress.

No studies were found of sexual or physical abuse in childhood and pelvic pain in men.

7.4.2 Anxiety

Findings of anxiety are common in samples of women with CPP (20), as are above-threshold scores on screening instruments for post-traumatic stress disorder (21).

In the latter study, higher post-traumatic stress disorder scores were associated with and contributed

up to 10% unique variance to poorer health status and more surgery visits. However, anxiety is likely to refer to pain being caused by fears of missed pathology, particularly cancer, and to uncertainties about treatment and the likely prognosis, treated or untreated.

Medical consultations and investigations are particular sources of anxiety, but also a focus of hopes of progress towards resolution of the problem. In a study of women with pelvic pain (22), laparoscopy was followed by a substantial reduction in pain, the size of which was predicted by beliefs about pain and about the seriousness of the woman's condition. Women with higher levels of pain at baseline and higher estimates of the seriousness of their condition reported greater pain reduction.

However, anxieties are not only about the possibility of serious pathology. Stones et al. (2000) reported that women with pelvic pain usually understand that nothing life-threatening has been or is likely to be found, but still want a diagnosis or explanation of the pain (23). Evidence from in-depth studies of women with pelvic pain underline the fact that they want to be assessed in terms of the effects of pain on their lives, to feel understood and have the pain legitimized. They hope for an explanation, and reassurance about the cause of pain and about the possibilities of treating it (24). When they feel disbelieved, and the emphasis is on the absence of physical findings rather than on their experience of pain, they are likely to feel invalidated and to drop out of treatment dissatisfied (25).

There appear to be no similar studies of men's concerns when consulting for persistent pelvic pain.

7.4.3 *Depression and catastrophizing*

Depression or depressed mood is also commonly found in both men and women with persistent pelvic pain (10), and has been reliably associated with urological symptoms (not necessarily painful) (10). However, this association may be no greater than in other persistent pain groups (26), when the comparison group is from a pain sample not from the healthy population.

In addition, there is an important methodological question to be asked of any study using diagnostic or standard assessment instruments, which is how symptoms have been attributed, many of which in depression are understood as neurovegetative signs. The core of clinical depression is negativity about the self. This describes relatively few people with persistent pain, who characteristically state that it is pain, not the self, at the base of their problem (27). Stones et al. (2000) suggested that 'psychological distress may be a consequence and not a cause of persistent pain: while identification of depression is important as part of treatment, caution is required before attribution of causality' (p. 416) (23).

Psychological factors contributing to adjustment to CPP/prostatitis in men was investigated in a large North American cohort study (28). Overall pain severity, sensory and affective aspects of pain was predicted by catastrophizing, particularly the helplessness dimension; pain distress was also predicted by depression. Impact of pain on activity was predicted by urinary symptoms, pain, and taking pain-contingent rest. An earlier study (28) has shown the importance of pain and depression for predicting lower quality of life in men with chronic prostatitis/pelvic pain.

7.4.4 *Impact of pain*

The impact of pain has been investigated almost entirely in terms of sexual problems. Women with pelvic pain reported these at a higher rate than did patients with other chronic pain disorders in a small UK study (29), with most reporting loss of interest in sex, dyspareunia, and pain following sexual intercourse. A much larger sample of women seeking healthcare, three-quarters for pelvic pain, yielded significantly lower quality of life scores than age-sex norms and widespread dissatisfaction with sexual activities, which was attributed to the pelvic problems (30). A small study of men with CPP (31) showed that pain severity predicted poorer sexual functioning, although this was less important than relationship variables.

There are short measures available to assess the impact of pain on the patient's life. The Brief Pain Inventory (www.mdanderson.org/pdf/bpif.pdf) consists of four 0 to 10 numerical scales for pain (current, average, worst, least) and seven scales for interference with specific aspects of life: general activity, mood, walking ability, normal work, relationships with other people, sleep, and enjoyment of life (32). A higher total score indicates greater pain and interference by pain with life. The EuroQoL is a quality-of-life scale (www.euroqol.org) available in various European languages and free for non-commercial use (33). It provides questions asking about mobility, self-care, pain, usual activities, and psychological status. There are also two questions on the impact of urinary symptoms and one on quality of life within the NIH Chronic Prostatitis Symptom Scale, CPSI (34).

7.5 **Summary: assessment recommendations** (Table 13)

A psychologist (or equivalent) is not required for this level of assessment, but access to regular discussion with a psychologist enables the clinician to interpret better the results of assessment.

7.5.1 Anxiety

It is important to obtain the patient's view of what is wrong or of what the patient is worried might be causing pain and other symptoms. Investment in establishing a trusting therapeutic relationship with the patient pays off when these questions are asked. Howard et al. (2003) suggest asking the patient, 'what do you believe or fear is the cause of your pain?' (35).

Investigations and results of examination should be explained clearly, in terms of what they can show, what they did or didn't show, and how this helps the investigations, attempts at diagnosis, or plans for treatment. This requires an adequate model of pain. Brief reassurance alone provides (at best) short-term relief of anxiety, after which the patient returns to seek help with the problem and the anxiety.

7.5.2 Depression

If the patient admits a depressed mood and attributes it to pain, it may be that the patient is interpreting information about and experience of pain and other symptoms in catastrophic ways. Good information can counteract this (as in anxiety). It may also be that the pain has had a serious impact on the patient's life; roles and satisfactions are lost because of pain, but can return with effective treatment. Encouragement to consider how to recover valued activities, with or without some pain relief, is helpful but the patient may require advice on how to do this from a pain management team.

7.5.3 Sexual and physical abuse in childhood

It is important to consider the possibility of physical and sexual abuse when taking the history, but disclosure can be difficult before a therapeutic relationship is established. It is not clear that pain, which the patient attributes to childhood sexual or physical abuse, should be managed any differently. Any disclosure of current physical or sexual abuse should be referred immediately to appropriate health, social or welfare services.

7.6 Psychological factors in treatment of pelvic pain (Table 14)

Untreated, there is a significant likelihood of symptom improvement. A follow-up study of women with pelvic pain referred to a clinic showed that 25% reported recovery (nearly half of them total recovery) over the 3 to 4 intervening years. However, neither pain nor distress at baseline, nor intervention received, was found to be associated with recovery (36).

Other sections cover the various physical (surgical, pharmacological, physiotherapeutic) interventions for male and female pelvic pain, and their outcomes. Psychological interventions may be directed:

1. At the pain itself, with the intended outcome of pain reduction and consequent reduction of impact of pain on life, or;
2. At adjustment to pain, with the intended outcome of improved mood and function and reduced healthcare use, with or without pain reduction.

The first category of interventions includes relaxation and biofeedback methods of controlling and decreasing pain by reducing muscle tension. Such methods are being applied to pelvic floor retraining, both in men (37,38) and women, sometimes alongside other physical therapies (see Section 6.5). The only RCT has been of a specific type of cognitively enhanced physical therapy applied to overall muscle tension, not to the pelvic floor. This Norwegian study (39) tested 'mensendieck somatocognitive therapy' combined with normal gynaecological treatment versus gynaecological treatment alone in women with CPP. Pain was reduced by 50% and motor function improved by 10 hours of physical therapy, with particular attention to tension and relaxation, and the thoughts and emotions that interfere with balanced posture and movement.

In the second category of interventions (see above), multicomponent pain management, involving education, physical retraining, behavioural change and increasing activity, relaxation, and cognitive therapy, is often applied to mixed groups of chronic pain patients, including those with pelvic pain, but there have been no RCTs of pelvic pain groups. There is a systematic review and meta-analysis (40), showing good outcome for mixed or back pain groups across pain experience, mood, coping, and activity. There is no reason to believe this does not apply to pelvic pain patients, but a trial of pelvic pain management alone is lacking.

The Cochrane database holds one systematic review and meta-analysis of treatments for pelvic pain, excluding pain due to endometriosis, irritable bowel syndrome, and chronic pelvic inflammatory disease (41). Treatment inclusion was broad: medical, surgical, physical, psychological, lifestyle, and complementary and alternative. Outcomes were mainly pain scores, quality of life and use of resources, including health care resources. This review, updated in 2005, includes 14 treatment trials covering pharmacotherapy; surgery; chiropractic; magnet therapy; counselling, psychoeducation, reassurance, and emotional disclosure; and a multicomponent pain management programme. Most trials were small, so the review covers 1,133 patients in all. Nine were of good methodological quality, but outcomes were mainly change in pain and other symptoms, with few including scales for mood (2), pain beliefs (1) or impact of pain (2).

From a psychological point of view, a minority of the trials in the Cochrane systematic review is of interest. The authors conclude in favour of educational counselling combined with ultrasound scan, which

improved pain and mood; and a multidisciplinary rehabilitative approach, including surgery, pharmacotherapy, physiotherapy, and psychosocial intervention, which improved function but not pain. An selective serotonin reuptake inhibitor antidepressant made no improvement in pain but improved function. Consultation using a photograph taken during laparoscopy had no effect; emotional disclosure (a stress reduction method) through writing brought about very small improvement in some pain scores.

It is regrettable that outcome measures correspond relatively poorly between studies with broadly similar aims, i.e. to restore the patient to a more normal lifestyle and better physical and psychological health. In the systematic review, the outcomes of pharmacotherapy trials and surgical and physical therapy trials are percentage pain reduction, self-rated pain score, physician-rated pain, and symptom scores, urinary measures such as peak flow rate (for persistent pelvic pain in men), pelvic tenderness (women); and uptake of further treatment following the trial treatment. A few studies included quality of life, but none measured mood change.

Mood change is a particular issue since, intentionally or not, any intervention, and even a good consultation can bring about cognitive, emotional, and/or behavioural change. Enabling the patient to understand what is causing the pain, and therefore the implications of the pain for everyday life and longer-term life goals, can be a major influence on the patient's successful management of pain. Furthermore, if all treatments sampled the same domains of pain in their evaluation, comparison across treatments, by medical personnel and by patients, would be more easily achieved (42).

Table 13: Psychological factors in the assessment of CPP

Assessment	Level of evidence	Grade of recommendation	Comment
<ul style="list-style-type: none"> Anxiety about cause of pain: 'Are you worried about what might be causing your pain? 	1a	C	Studies of women only: men's anxieties not studied
<ul style="list-style-type: none"> Depression attributed to pain: ask 'How has the pain affected your life?' 'How does the pain make you feel emotionally?' 	1a	C	Studies of women only: men's anxieties not studied
<ul style="list-style-type: none"> Multiple physical symptoms/ general health 	1a	C	
<ul style="list-style-type: none"> History of sexual or physical abuse 	1a	C	Current/recent abuse may be more important

Table 14: Treatment factors in the management of CPP

Treatment	Level of evidence	Grade of recommendation	Comment
<ul style="list-style-type: none"> Tension-reduction; relaxation, for pain reduction 	1b	A	Relaxation +/- biofeedback +/- physical therapy; mainly male pelvic pain
<ul style="list-style-type: none"> Multidisciplinary pain management for well-being 	(1a)	(A)	Pelvic pain patients treated within larger group: no specific pelvic pain trials

7.7 References

- Melzack R, Wall PD. Pain mechanisms: a new theory. *Science* 1965;150(699):971-979.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5320816&dopt=Abstract
- Chapman CR, Gavrin J. Suffering: the contributions of persistent pain. *Lancet* 1999;353(9171):2233-2237.
http://www.ncbi.nlm.nih.gov/pubmed/10393002?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- Grace VM. Pitfalls of the medical paradigm in chronic pelvic pain. *Baillieres Best Pract Res Clin Obstet Gynaecol* 2000;14(3):525-539.
http://www.ncbi.nlm.nih.gov/pubmed/10962640?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
- Sharpe M, Carson A. 'Unexplained' somatic symptoms, functional syndromes, and somatization: do we need a paradigm shift? *Ann Intern Med* 2001;134:926-930.

5. Flor H, Birbaumer N, Schugens MM, Lutzenberger W. Symptom-specific psychophysiological responses in chronic pain patients. *Psychophysiol* 1992;29:452-460.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1410176&dopt=Abstract
6. Link CL, Lutfey KE, Steers WD, McKinlay JD. Is abuse causally related to urologic symptoms? Results from the Boston Area Community Health (BACH) Survey. *Eur Urol* 2007;52:397-406.
<http://www.ncbi.nlm.nih.gov/pubmed/17383083>
7. Keefe FJ, Rumble ME, Scipio CD, Giordano LA, Perri LM. Psychological aspects of persistent pain: current state of the science. *J Pain* 2004;5(4):195-211.
http://www.ncbi.nlm.nih.gov/pubmed/15162342?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
8. Eccleston C, Crombez G. Pain demands attention: a cognitive-affective model of the interruptive function of pain. *Psychol Bull* 1999;125(3):356-366.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10349356&dopt=Abstract
9. Vlaeyen JW, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 2000;85(3):317-332.
http://www.ncbi.nlm.nih.gov/pubmed/10781906?ordinalpos=9&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
10. Fitzgerald MP, Link CL, Litman HJ, Trivison TG, McKinlay JB. Beyond the lower urinary tract: the association of urological and sexual symptoms with common illnesses. *Eur Urol* 2007;52(2):407-415.
http://www.ncbi.nlm.nih.gov/pubmed/17382458?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
11. Schur EA, Afari N, Furberg H, Olarte M, Goldberg J, Sullivan PF, Buchwald D. Feeling bad in more ways than one: comorbidity patterns of medically unexplained and psychiatric conditions. *J Gen Intern Med* 2007;22(6):818-821.
http://www.ncbi.nlm.nih.gov/pubmed/17503107?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
12. Warren JW, Jackson TL, Langenberg P, Meyers DJ, Xu J. Prevalence of interstitial cystitis in first-degree relatives of patients with interstitial cystitis. *Urology* 2004;63(1):17-21.
http://www.ncbi.nlm.nih.gov/pubmed/14751339?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
13. Savidge CJ, Slade P. Psychological aspects of chronic pelvic pain. *J Psychosom Res* 1997;42(5):433-444.
http://www.ncbi.nlm.nih.gov/pubmed/9194016?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
14. Latthe P, Mignini L, Gray R, Hills R, Khan K. Factors predisposing women to chronic pelvic pain: systematic review. *BMJ* 2006;332(7544):749-755.
http://www.ncbi.nlm.nih.gov/pubmed/16484239?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
15. Davis DA, Luecken LJ, Zautra AJ. Are reports of childhood abuse related to the experience of chronic pain in adulthood? A meta-analytic review of the literature. *Clin J Pain* 2005;21(5):398-405.
http://www.ncbi.nlm.nih.gov/pubmed/16093745?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
16. Hilden M, Schei B, Swahnberg K, Halmesmaki E, Langhoff-Roos J, Offerdal K, Pikarinen U, Sidenius K, Steingrimsdottir T, Stoum-Hinsverk H, Wigman B. A history of sexual abuse and health: a Nordic multicentre study. *BJOG* 2004;111(10):1121-1127.
http://www.ncbi.nlm.nih.gov/pubmed/15383115?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
17. Raphael KG, Widom CS, Lange G. Childhood victimization and pain in adulthood: a prospective investigation. *Pain* 2001;92(1-2):283-293.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11323150&dopt=Abstract
18. Poleshuck EL, Dworkin RH, Howard FM, Foster DC, Shield CG, Giles DE, Tu X. Contributions of physical and sexual abuse to women's experiences with chronic pelvic pain. *J Reprod Med* 2005;50(2):91-100.
http://www.ncbi.nlm.nih.gov/pubmed/15755045?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

19. Chandler HK, Ciccone DS, Raphael KG. Localization of pain and self-reported rape in a female community sample. *Pain Med* 2006;7(4):344-352.
http://www.ncbi.nlm.nih.gov/pubmed/16898946?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
20. Zondervan KT, Yudkin PL, Vessey MP, Jenkinson CP, Dawes MG, Barlow DH, Kennedy SH. The community prevalence of chronic pelvic pain in women and associated illness behaviour. *B J Gen Pract* 2001;51(468):541-547.
http://www.ncbi.nlm.nih.gov/pubmed/11462313?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
21. Meltzer-Brody S, Leserman J, Zolnoun D, Steege J, Green E, Teich A. Trauma and posttraumatic stress disorder in women with chronic pelvic pain. *Obstet Gynecol* 2007;109(4):902-908.
http://www.ncbi.nlm.nih.gov/pubmed/17400852?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
22. Elcombe S, Gath D, Day A. The psychological effects of laparoscopy on women with chronic pelvic pain. *Psychol Med* 1997;27(5):1041-1050.
http://www.ncbi.nlm.nih.gov/pubmed/9300510?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
23. Stones RW, Selfe SA, Fransman S, Horn SA. Psychosocial and economic impact of chronic pelvic pain. *Baillieres Best Pract Res Clin Obstet Gynaecol* 2000;14(3):415-431.
http://www.ncbi.nlm.nih.gov/pubmed/10962635?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
24. Price J, Farmer G, Harris J, Hope T, Kennedy S, Mayou R. Attitudes of women with chronic pelvic pain to the gynaecological consultation: a qualitative study. *BJOG* 2006;113(4):446-452.
http://www.ncbi.nlm.nih.gov/pubmed/16489938?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
25. McGowan L, Luker K, Creed F, Chew-Graham CA. 'How do you explain a pain that can't be seen?': the narratives of women with chronic pelvic pain and their disengagement with the diagnostic cycle. *B J Health Psychol* 2007;12(Pt 2):261-274.
http://www.ncbi.nlm.nih.gov/pubmed/17456285?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
26. Heinberg LJ, Fisher BJ, Wesselman U, Reed J, Haythornthwaite JA. Psychological factors in pelvic/urogenital pain: the influence of site of pain versus sex. *Pain* 2004;108(1-2):88-94.
http://www.ncbi.nlm.nih.gov/pubmed/15109511?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
27. Pincus T, Williams A. Models and measurements of depression in chronic pain. *J Psychosom Res* 1999;47(3):211-219.
http://www.ncbi.nlm.nih.gov/pubmed/10576470?ordinalpos=7&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
28. Tripp DA, Curtis Nickel J, Landis JR, Wang YL, Knauss JS, CPCR Study Group. Predictors of quality of life and pain in chronic prostatitis/chronic pelvic pain syndrome: findings from the National Institutes of Health Chronic Prostatitis Cohort Study. *BJU Int* 2004;94(9):1279-1282.
http://www.ncbi.nlm.nih.gov/pubmed/15610105?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
29. Collett BJ, Cordle J, Steward R, Jagger C. A comparative study of women with chronic pelvic pain, chronic nonpelvic pain and those with no history of pain attending general practitioners. *Br J Obstet Gynaecol* 1998;105(1):87-92.
http://www.ncbi.nlm.nih.gov/pubmed/9442168?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
30. Kuppermann M, Learman LA, Schembri M, Gregorich S, Jacoby A, Jackson RA, Gates E, Wassel-Fyr C, Lewis J, Washington AE. Effect of noncancerous pelvic problems on health-related quality of life and sexual functioning. *Obstet Gynecol* 2007;110(3):633-642.
http://www.ncbi.nlm.nih.gov/pubmed/17766611?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
31. Smith KB, Tripp D, Pukall C, Nickel JC. Predictors of sexual and relationship functioning in couples with Chronic Prostatitis/Chronic Pelvic Pain Syndrome. *J Sex Med* 2007;4(3):734-744.
http://www.ncbi.nlm.nih.gov/pubmed/17451490?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

32. Tan G, Jensen MP, Thornby JI, Shanti BF. Validation of the Brief Pain Inventory for chronic nonmalignant pain, *J Pain* 2004;5(2):133-137.
http://www.ncbi.nlm.nih.gov/pubmed/15042521?ordinalpos=18&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
33. Rabin R, de Charro F. EQ-5D: a measure of health status from the EuroQol group. *Ann Med* 2001;33(5):337-343.
http://www.ncbi.nlm.nih.gov/pubmed/11491192?ordinalpos=6&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
34. Turner JA, Ciol MA, Von Korff M, Berger R. Validity and responsiveness of the national institutes for health chronic prostatitis symptom index. *J Urol* 2003;169(2):580-583.
http://www.ncbi.nlm.nih.gov/pubmed/12544311?ordinalpos=19&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
35. Howard FM. Chronic pelvic pain. *Obstet Gynecol* 2003;101(3):594-611.
http://www.ncbi.nlm.nih.gov/pubmed/12636968?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
36. Weijenborg PTM, Greeven A, Dekker FW, Peters AW, ter Kuile MM. Clinical course of chronic pelvic pain in women. *Pain* 2007;132(Suppl 1):S117-123.
http://www.ncbi.nlm.nih.gov/pubmed/17689866?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
37. Anderson RU, Wise D, Sawyer T, Chan C. Integration of myofascial trigger point release and paradoxical relaxation training treatment of chronic pelvic pain in men. *J Urol* 2005;174(1):155-160.
http://www.ncbi.nlm.nih.gov/pubmed/15947608?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
38. Cornel EB, van Haast EP, Schaarsberg RW, Geels J. The effect of biofeedback physical therapy in men with chronic pelvic pain syndrome Type III. *Eur Urol* 2005;47(5):607-611.
http://www.ncbi.nlm.nih.gov/pubmed/15826751?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
39. Haugstad GK, Haugstad TS, Kirste UM, Leganger S, Klemmetsen I, Malt UF. Mensendieck somatocognitive therapy as treatment approach to chronic pelvic pain: Results of a randomized controlled intervention study. *Am J Obs Gynecol* 2006;194(5):1303-1310.
http://www.ncbi.nlm.nih.gov/pubmed/16647914?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
40. Morley SJ, Eccleston C, Williams A. Systematic review and meta-analysis of randomised controlled trials of cognitive behaviour therapy and behaviour therapy for chronic pain in adults, excluding headache. *Pain* 1999;80(1-2):1-13.
http://www.ncbi.nlm.nih.gov/pubmed/10204712?ordinalpos=5&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
41. Stones W, Cheong YC, Howard FM. Interventions for treatment chronic pelvic pain in women. *Cochrane Database Syst Rev* 2005;(2):CD000387.
<http://mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD000387/frame.html>
42. Dworkin RH, Turk DC, Farrar JT, Haythornthwaite JA, Jensen MP, Katz NP, Kerns RD, Stucki G, Allen RR, Bellamy N, Carr DB, Chandler J, Cowan P, Dionne R, Galer BS, Hertz S, Jadad AR, Kramer LD, Manning DC, Martin S, McCormick CG, McDermott MP, McGrath P, Quessy S, Rappaport BA, Robbins W, Robinson JP, Rothman M, Royal MA, Simon L, Stauffer JW, Stein W, Tollett J, Wernicke J, Witter J;IMMPACT. Topical review and recommendations: Core outcome measures for chronic pain clinical trials: IMMPACT recommendations. *Pain* 2005;113(1-2):9-19.
http://www.ncbi.nlm.nih.gov/pubmed/15621359?ordinalpos=10&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

8. GENERAL TREATMENT OF CHRONIC PELVIC PAIN

There is very little specific evidence for the role of analgesic and co-analgesic drugs in CPP, with one systematic review in women suggesting that further research is required (1). As CPP is thought to be modulated by similar mechanisms to those of somatic, visceral and neuropathic pain, the evidence below has been derived from the general chronic pain literature. This section has divided the drugs into their broad classes for

ease of reading and reference.

Many of the agents will be familiar to chronic pain clinicians and their involvement with difficult cases will help in the development of individualized pain management plans. Some drugs discussed below are best used by clinicians experienced in their use. The use of drugs for specific urogenital conditions is covered by the guidelines for that condition. This chapter looks at the general considerations for patients within the commonly used drug classes.

The evidence grading is the same as elsewhere in the EAU guidelines, looking at outcome of treatment. Recommendations are based on the likely benefit to a patient in terms of pain or improved quality of life rather than on the basis of disease modification. Table 15 provides an overview of available treatments together with evidence levels and graded recommendations.

8.1 Simple analgesics

8.1.1 Paracetamol

Paracetamol should be considered on its own. It has analgesic and antipyretic activity and is used in acute and chronic painful conditions (2). There is evidence for its use in somatic pain and arthritic pain in particular (3,4) but the benefits are limited. There is little evidence about its role in CPP.

8.1.2 Acidic antipyretic analgesics

The classical NSAIDs fall into this group and include salicylic acid. They are known to act on the cyclo-oxygenase (COX) enzyme. The early NSAIDs tended to have little selectivity for COX2 over COX1, and are therefore said to be associated with more side effects than the newer COX2 selective inhibitors. The COX1 enzyme is mainly involved in normal 'housekeeping' functions, such as mediating gastric mucosal integrity, and renal and platelet function. Blocking the COX1 enzyme is the cause of the platelet, gastric and renal complications that can occur with NSAIDs. It has been suggested that the COX2 enzyme is inducible as a result of tissue damage, and that it is the main enzyme involved in inflammation and peripheral sensitization of nociceptors. As a result, the analgesic efficacy of COX2 selective drugs should be as good as that of the non-selective drugs. This, however, has been disputed (5). More recent studies and reviews including those from the European and US drug advisory bodies highlight the cardiovascular risks associated with COX2 selective agents (6-8). The risk for cardiac risk is clear for COX2 agents and classical NSAIDs may also be incriminated (9). Thus, selective COX2 agents should not be prescribed in patients with increased risk of cardiovascular disease including congestive cardiac failure.

There is very little evidence for NSAIDs to be used in the management of CPP and even less evidence for COX2 selective drugs. Most analgesic studies have investigated dysmenorrhoea, in which NSAIDs were found to be superior to placebo and possibly paracetamol (10).

For practical purposes the NSAIDs may be divided into:

- non-selective, low potency (e.g. salicylic acid, ibuprofen, mefenamic acid)
- non-selective, high potency (e.g. ketoprofen, diclofenac, ketorolac)
- COX2 selective drugs (e.g. celecoxib, etoricoxib).

8.1.3 Guidelines for use of NSAIDs and COX2 selective agents

- Non-selective, low potency NSAIDs should be used in the first instance. They are most likely to be of help if there is an inflammatory component to the pain. More potent NSAIDs should be reserved for those conditions in which the low potency drugs have been tried and failed to produce significant benefit.
- COX2 selective drugs should be used with caution as an alternative to the non-selective drugs where there is an increased risk of gastric complications. They should be avoided in patients with known cardiovascular disease.
- NSAIDs should be taken with food and consideration must be given to the use of gastric protective agents.
- The benefits of the NSAIDs must be demonstrated to outweigh the risks.
- All NSAIDs are contraindicated in active gastrointestinal ulceration/bleeding and renal disease. They may exacerbate asthma and produce fluid retention.
- Even if stronger analgesics such as opioids are added, the NSAIDs can be continued as they are likely to have a synergistic action improving pain control above and beyond that obtained with opioids alone (11).
- Paracetamol should be considered as an alternative to, or given with, NSAIDs as it is well tolerated with few side effects

8.2 Neuropathic analgesics

8.2.1 Tricyclic antidepressants

There is very little evidence in CPP for tricyclic antidepressants in humans. What data is available includes these drugs as options in treatment (12-14). A study in cats does suggest that tricyclics may have a role in the management of cystitis (15). Most studies involve neuropathic pain. If there is a suggestion of nerve injury or central sensitization, the algorithm outlined in Figure 8 should be considered.

Saarto and Wiffen (16) reviewed antidepressants for neuropathic pain. They concluded that tricyclics are effective for neuropathic pain with limited evidence for the selective serotonin reuptake inhibitor antidepressant drugs and insufficient evidence for other antidepressants. The study suggests that HIV-related pain does not respond and further work is required with regard to more specific conditions.

Tricyclic antidepressants tend to be used in lower dosages than those required for treating depression. Amitriptyline is used up to a dose of 150 mg once daily.

8.2.2 Anticonvulsants

These drugs have been used in the management of pain for many years. Whereas there is little evidence to support the use of anticonvulsants in the management of genitourinary pain, they should be considered if there is a suggestion of neuropathic pain or central sensitization. There is no role for these drugs in acute pain (17).

Gabapentin has been introduced for pain management and has undergone a systematic review by Wiffen (18). It is said to have fewer serious side effects compared to the older anticonvulsants and in some countries is licensed for use in chronic neuropathic pain. There are claims that it produces a more natural sleep state at night than antidepressants.

The evidence (for neuropathic pain) does not demonstrate any superiority for gabapentin over carbamazepine (17). The numbers needed to harm (for minor harm) is 2.5 for gabapentin and 3.7 for carbamazepine. Many practitioners, however, would not use carbamazepine as a first-line anticonvulsant in pain management because of its potentially serious side effects (blood, hepatic or skin disorders).

Carbamazepine and other anticonvulsants (e.g. phenytoin or valproate) have been used for neuropathic pain but are best reserved for practitioners familiar with their use.

8.2.3 *N*-methyl-*D*-aspartate (NMDA) antagonists

The NMDA receptor channel complex is known to be an important channel for the development and maintenance of chronic pain. It is felt to be particularly important when there is evidence of central sensitization and opioid tolerance (19-23). These are the phenomena that alter signal transmission within the nervous system so that non-painful stimuli may become painful (allodynia) and pain from a painful stimulus is magnified (hyperalgesia). NMDA antagonists have been used in the management of neuropathic pain (24).

Ketamine has been useful in several chronic pain states including peripheral neuropathies with allodynia, stump and phantom pain, central pain, and cancer-related pain, with and without a neurological component, but its long-term role remains unclear (25).

Ketamine has been used as a general anaesthetic for over 30 years. It has also been used as an intravenous analgesic in burns units and accident and emergency units. Ketamine is thought to act primarily at the NMDA receptor, though it may also have action at sodium channels and opioid (κ and μ) receptors (23). Difficult urogenital pains may therefore be helped by ketamine if there is evidence of nerve injury or central sensitization.

Ketamine is not licensed for use in chronic pain and like the opioids has been used as a street drug of addiction. Ketamine should only be started by an experienced practitioner trained in its use. Similar care to that of opioids must be taken if a patient is to be managed at home.

8.2.4 Sodium channel blockade

In a significant number of patients with urogenital pain, nerve injury and neuropathic changes are thought to play a role. These may be associated with a reduction in some sodium channels and the development of novel sodium channels. As a result, injured afferents become prone to generating more prolonged and higher frequency discharges, with a reduced refractory period. These changes in the characteristics of sodium channels are thought to underlie the mechanisms of mechanosensitivity, thermosensitivity and chemosensitivity (26). They may be involved in some visceral hyperalgesia.

Human studies have demonstrated that intravenous lidocaine reduces neuropathic pain and sensory phenomena, such as allodynia (27). A positive lidocaine challenge may be followed by repeated infusions of lidocaine and the benefit from a single infusion may be prolonged.

A role for the oral analogue, mexiletine, has also to be defined (28); a positive response to intravenous lidocaine does not always indicate that mexiletine will work.

Challapalli et al. have reviewed the role of local anaesthetic agents in neuropathic pain. They are safe and better than placebo and as effective as other analgesics. Further research is needed into specific painful

conditions and outcome measures to assess whether pain relief is clinically significant (29). Again, these agents are being used outside their licence and are best instituted by clinicians familiar with them.

8.3 Opioids

There is now a general acceptance that opioids have a role in the management of chronic non-malignant pain (30). Studies have tended to be short term and a systematic review concluded that further research is required into the long-term use of opioids (31). The use of opioids in urogenital pain is poorly defined. Their use in neuropathic pain remains equivocal but a meta-analysis suggests clinically important benefits (32). The authors emphasize that more research is needed in long-term outcomes and side effects. There is also evidence suggesting that opioids may produce different responses with different types of pain (33).

Generally, slow-release preparations are preferred for chronic pain. Side effects are common but rarely serious. If, however, a particular agent causes side effects and clinical benefit, rotating to another opioid may be beneficial. Titrating the dose should be closely monitored to assess both benefit and side effects. Rotating from one opioid to another also requires close monitoring as there are no exact dose equivalents. General guidelines for the use of opioids in chronic pain have been published (34,35). The following guidelines are suggested, but the clinician involved should be familiar with the use of opioids in non-malignant pain.

8.3.1 Guidelines for the use of opioids in chronic/non-acute urogenital pain

1. All other reasonable treatments must have been tried and failed.
2. The decision to instigate long-term opioid therapy should be made by an appropriately trained specialist in consultation with another physician (preferably the patient's family doctor).
3. Where there is a history or suspicion of drug abuse, a psychiatrist or psychologist with an interest in pain management and drug addiction should be involved.
4. The patient should undergo a trial of opioids.
5. The dose required needs to be calculated by careful titration.
6. The patient should be made aware (and possibly give written consent):
 - I. That opioids are strong drugs and associated with addiction and dependency.
 - II. The opioids will normally only be prescribed from one source (preferably the family doctor).
 - III. The drugs will be prescribed for fixed periods of time and a new prescription will not be available until the end of that period.
 - IV. The patient will be subjected to spot urine and possibly blood checks to ensure that the drug is being taken as prescribed and that non-prescribed drugs are not being taken.
 - V. Inappropriate aggressive behaviour associated with demanding the drug will not be accepted.
 - VI. Hospital specialist review will normally occur at least once a year.
 - VII. The patient may be requested to attend a psychiatric/psychology review.
 - VIII. Failure to comply with the above may result in the patient being referred to a drug dependency agency and the use of therapeutic, analgesic opioids being stopped.
7. Morphine is the first-line drug, unless there are contraindications to morphine or special indications for another drug. The drug should be prescribed in a slow-release/modified release form. Short-acting preparations are undesirable and should be avoided where possible. Parenteral dosing is undesirable and should be avoided where possible.

8.3.2 Morphine

There is no compelling evidence that one opiate is better than another. Morphine is the traditional gold standard and the opioid many physicians are most familiar with. In an acute situation, the daily morphine requirement may be calculated by titration of rapid-release morphine. In chronic pain situations, starting with a low dose of slow-release morphine and titrating the dose every 3 days to 1 week is adequate.

8.3.3 Transdermal fentanyl

Transdermal fentanyl is used when oral absorption is restricted or when the patient suffers with intolerable side effects from other opioids. Patients with rapid bowel transit times (e.g. ileostomy) may find transdermal preparations beneficial. Patches are generally changed every 72 hours.

8.3.4 Methadone

Methadone is a strong analgesic with a long track record of use. It has opioid and NMDA-antagonistic activity (24). It is suggested that further work is needed to look at its role in neuropathic pain (33). Its use is supported as a fourth-line agent in treating neuropathic pain in a consensus document by the Canadian Pain Society (36). Rotating from other opioids to methadone is not an exact science because dosing ratios are not clearly

understood (37). Metabolite accumulation and cardiac side effects can be a problem. A practitioner familiar with its use as an analgesic should prescribe methadone.

8.3.5 *Oxycodone*

A slow-release preparation is available with evidence suggesting its benefit in neuropathic pain. Evidence suggests that oxycodone has benefits over morphine in some experimentally induced visceral pains (38). The pharmacology of oxycodone is different to morphine in experimental neuropathic states (39). An RCT has also demonstrated a role for oxycodone in neuropathic pain (diabetic neuropathy) (40).

8.3.6 *Other opioids and opioid-like agents*

Other opioids are available as slow- or modified-release preparations. They may be useful for opiate rotation if side effects or tolerance is a problem.

Buprenorphine and pentazocine both have agonist and antagonist properties and can induce withdrawal symptoms in patients used to opioids. Naloxone may only partly reverse respiratory depression. Buprenorphine topical patches are now available and may offer a similar advantage to topical fentanyl.

Codeine and dihydrocodeine are effective for the relief of mild-to-moderate pain. They are limited by side effects (notably constipation) and genetic variance of metabolism that affects analgesic efficacy. Approximately 10% of individuals will not effectively metabolize codeine resulting in inadequate analgesia (41).

Tramadol produces analgesia by two mechanisms, an opioid effect and an enhancement of serotonergic and adrenergic pathways (42,43). It has fewer typical opioid side effects (especially less respiratory depression, less constipation and less addiction potential) and is available in a slow-release preparation. A Cochrane review suggests that there is a role for tramadol in neuropathic pain management (44).

Figure 8: Guidelines for the use of neuropathic analgesics




Table 15: General treatment of CPP

Type of pain	Level of evidence	Grade of recommendation	Comment
Paracetamol for somatic pain	1b	A	Benefit is limited and based on arthritic pain
COX2 antagonists	1b	A	Avoid in patients with cardiovascular risk factors
NSAIDs for dysmenorrhoea	1a	B	Better than placebo but unable to distinguish between agents
Tricyclic antidepressants		1a	A Neuropathic pain
Tricyclic antidepressants		3	C Evidence suggests pelvic pain is similar to neuropathic pain
Anticonvulsants Gabapentin	1a	A	For neuropathic pain
Opioids for chronic non malignant pain	1a	A	Limited long-term data Should only be used by clinicians experienced in their use
Opioids for neuropathic pain	1a	A	Benefit is probably clinically significant Caution with use, as above

COX = cyclo-oxygenase; NSAID = non-steroidal anti-inflammatory drug.

8.4 References

1. Stones W, Cheong YC, Howard FM. Interventions for treating chronic pelvic pain in women. Cochrane Database System Rev 2005;(2):CD000387.
<http://mrw.interscience.wiley.com/cochrane/clsysrev/articles/CD000387/frame.html>
2. Bannwarth B, Pehourcq F. [Pharmacologic basis for using paracetamol: pharmacokinetic and pharmacodynamic issues.] *Drugs* 2003;63 Spec No 2:5-13. [French]
http://www.ncbi.nlm.nih.gov/pubmed/14758786?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
3. Bradley JD, Brandt KD, Katz BP, Kalasinski LA, Ryan SI. Treatment of knee osteoarthritis: relationship of clinical features of joint inflammation to the response to a nonsteroidal antiinflammatory drug or pure analgesic. *J Rheumatol* 1992;19(12):1950-1954.
http://www.ncbi.nlm.nih.gov/pubmed/1294745?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
4. Williams HJ, Ward JR, Egger MJ, Neuner R, Brooks RH, Clegg DO, Field EH, Skosey JL, Alarcon GS, Willkens RF, et al., Comparison of naproxen and acetaminophen in a two-year study of treatment of osteoarthritis of the knee. *Arthritis Rheum* 1993;36(9):1196-1206.
http://www.ncbi.nlm.nih.gov/pubmed/8216413?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
5. McCormack K, Twycross R. Cox-2-selective inhibitors and analgesia. *Pain Clinical Updates* 2002;10(1).
6. Jones SF, Power I. Postoperative NSAIDs and COX-2 inhibitors: cardiovascular risks and benefits. *Br J Anaesth* 2005;95(3):281-284.
http://www.ncbi.nlm.nih.gov/pubmed/16076922?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
7. Medicines and Healthcare products Regulatory Agency (MHRA) release. Cardiovascular Safety of Non-Steroidal Anti-inflammatory Drugs. Overview of key data. MHRA, 2005.
<http://www.mhra.gov.uk/Safetyinformation/Generalsafetyinformationandadvice/Product-specificinformationandadvice/CardiovascularsafetyofCOX-2inhibitorsandnon-selectiveNSAIDs/index.htm>
8. US Food and Drug Administration (FDA). Questions and Answers. FDA Regulatory Actions for the COX-2 Selective and Non-Selective Non-Steroidal Anti-inflammatory drugs (NSAIDs).
<http://www.fda.gov/cder/drug/infopage/COX2/COX2qa.htm>
9. Kearney PM, Baigent C, Godwin J, Halls H, Emberson JR, Patrono C. Do selective cyclo-oxygenase-2 inhibitors and traditional non-steroidal anti-inflammatory drugs increase the risk of atherothrombosis? Meta-analysis of randomised trials. *BMJ* 2006;332(7553):1302-1308.
http://www.ncbi.nlm.nih.gov/pubmed/16740558?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

10. Marjoribanks J, Proctor ML, Farquhar C. Nonsteroidal anti-inflammatory drugs for primary dysmenorrhoea. *Cochrane Database Syst Rev* 2003;(4):CD001751.
http://www.ncbi.nlm.nih.gov/pubmed/14583938?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
11. Christie MJ, Vaughan CW, Ingram SL. Opioids, NSAIDs and 5-lipoxygenase inhibitors act synergistically in brain via arachidonic acid metabolism. *Inflamm Res* 1999;48(1):1-4.
http://www.ncbi.nlm.nih.gov/pubmed/9987677?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
12. Greco CD. Management of adolescent chronic pelvic pain from endometriosis: a pain center perspective. *J Pediatr Adolesc Gynecol* 2003;16(3 Suppl):S17-S19.
http://www.ncbi.nlm.nih.gov/pubmed/12742182?ordinalpos=5&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
13. Phatak S, Foster HE Jr. The management of interstitial cystitis: an update. *Nat Clin Pract Urol* 2006;3(1):45-53.
http://www.ncbi.nlm.nih.gov/pubmed/16474494?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
14. Pontari MA. Chronic prostatitis/chronic pelvic pain syndrome in elderly men: toward better understanding and treatment. *Drugs Aging* 2003;20(15):1111-1125.
http://www.ncbi.nlm.nih.gov/pubmed/14651434?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
15. Chew DJ, Buffington CA, Kendall MS, DiBartola SP, Woodworth BE. Amitriptyline treatment for severe recurrent idiopathic cystitis in cats. *J Am Vet Med Assoc* 1998;213(9):1282-1286.
http://www.ncbi.nlm.nih.gov/pubmed/9810383?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
16. Saarto T, Wiffen PJ. Antidepressants for neuropathic pain. *Cochrane Database Syst Rev* 2007;17(4):CD005454.
http://www.ncbi.nlm.nih.gov/pubmed/17943857?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
17. Wiffen P, Collins S, McQuay H, Carroll D, Jadad A, Moore A. Anticonvulsant drugs for acute and chronic pain. *Cochrane Database Syst Rev* 2005;(3):CD001133.
http://www.ncbi.nlm.nih.gov/pubmed/16034857?ordinalpos=11&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
18. Wiffen PJ, McQuay HJ, Edwards JE, Moore RA. Gabapentin for acute and chronic pain. *Cochrane Database Syst Rev*. 2005;(3):CD005452.
http://www.ncbi.nlm.nih.gov/pubmed/16034978?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
19. Price DD, Mayer DJ, Mao J, Caruso FS. NMDA-receptor antagonists and opioid receptor interactions as related to analgesia and tolerance. *J Pain Symptom Manage* 2000;19(1 Suppl):S7-S11.
http://www.ncbi.nlm.nih.gov/pubmed/10687332?ordinalpos=9&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
20. Eide PK, Jørum E, Stubhaug A, Bremnes J, Breivik H. Relief of post-herpetic neuralgia with the N-methyl-D-aspartic acid receptor antagonist ketamine: a double-blind, cross-over comparison with morphine and placebo. *Pain* 1994;58(3):347-354.
http://www.ncbi.nlm.nih.gov/pubmed/7838584?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
21. Guirimand F, Dupont X, Brasseur L, Chauvin M, Bouhassira D. The effects of ketamine on the temporal summation (wind-up) of the R(III) nociceptive flexion reflex and pain in humans. *Anaesth Analg* 2000;90(2):408-414.
http://www.ncbi.nlm.nih.gov/pubmed/10648330?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
22. Laurido C, Pelissier T, Pérez H, Flores F, Hernández A. Effect of ketamine on spinal cord nociceptive transmission in normal and monoarthritic rats. *Neuroreport* 2001;12(8):1551-1554.
http://www.ncbi.nlm.nih.gov/pubmed/11409714?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
23. Mikkelsen S, Ilkjaer S, Brennum J, Borgbjerg FM, Dahl JB. The effect of naloxone on ketamine-induced effects on hyperalgesia and ketamine-induced side effects in humans. *Anaesthesiology* 1999;90(6):1539-1545.
http://www.ncbi.nlm.nih.gov/pubmed/10360849?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

24. Hewitt DJ. The use of NMDA-receptor antagonists in the treatment of chronic pain. *Clin J Pain* 2000;16(2 Suppl):S73-S79.
http://www.ncbi.nlm.nih.gov/pubmed/10870744?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
25. Visser E, Schug SA. The role of ketamine in pain management. *Biomed Pharmacother* 2006;60(7):341-348.
http://www.ncbi.nlm.nih.gov/pubmed/16854557?ordinalpos=8&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
26. Cummins T et al. Sodium channels as molecular targets in pain. In: Devor M, Rowbotham M, Wiesenfeld-Hallin Z, eds. *Proceedings of the 9th World Congress on Pain*. Seattle: IASP, 2000:77-91.
27. Baranowski AP, De Courcey J, Bonello E. A trial of intravenous lidocaine on the pain and allodynia of postherpetic neuralgia. *J Pain Symptom Manage* 1999;17(6):429-433.
http://www.ncbi.nlm.nih.gov/pubmed/10388248?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
28. Galer BS, Harle J, Rowbotham MC. Response to intravenous lidocaine infusion predicts subsequent response to oral mexiletine: a prospective study. *J Pain Symptom Manage* 1996;12(3):161-167.
http://www.ncbi.nlm.nih.gov/pubmed/8803379?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
29. Challapalli V, Tremont-Lukats IW, McNicol ED, Lau J, Carr DB. Systemic administration of local anesthetic agents to relieve neuropathic pain. *Cochrane Database Syst Rev* 2005;(4):CD003345.
http://www.ncbi.nlm.nih.gov/pubmed/16235318?ordinalpos=4&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
30. McQuay H. Opioids in pain management. *Lancet* 1999;353(9171):2229-2232.
http://www.ncbi.nlm.nih.gov/pubmed/10393001?ordinalpos=8&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
31. Kalso E, Edwards JE, Moore RA, McQuay HJ. Opioids in chronic non-cancer pain: systematic review of efficacy and safety. *Pain* 2004;s112(3):372-380.
http://www.ncbi.nlm.nih.gov/pubmed/15561393?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
32. Eisenberg E, McNicol E, Carr DB. Opioids for neuropathic pain. *Cochrane Database Syst Rev*. 2006;(3):CD006146.
http://www.ncbi.nlm.nih.gov/pubmed/16856116?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
33. Lemberg K, Kontinen VK, Viljakka K, Kylänlahti I, Yli-Kauhaluoma J, Kalso E. Morphine, oxycodone, methadone and its enantiomers in different models of nociception in the rat. *Anaesth Analg* 2006;102(6):1768-1774.
http://www.ncbi.nlm.nih.gov/pubmed/16717324?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
34. Kalso E, Allan L, DelleMijn PL, Faura CC, Ilias WK, Jensen TS, Perrot S, Plaghki LH, Zenz M. Recommendations for using opioids in chronic non-cancer pain. *Eur J Pain* 2003;7(5):381-386.
http://www.ncbi.nlm.nih.gov/pubmed/12935789?ordinalpos=5&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
35. The Pain Society, Recommendations for the appropriate use of opioids for persistent non-cancer pain. A consensus statement prepared on behalf of the Pain Society, the Royal College of Anaesthetists, the Royal College of General Practitioners and the Royal College of Psychiatrists. London: The Pain Society, 2004.
http://www.britishpainsociety.org/book_opioid_main.pdf
36. Moulin DE, Clark AJ, Gilon I, Ware MA, Watson CP, Sessle BJ, Coderre T, Morley-Forster PK, Stinson J, Boulanger A, Peng P, Finley GA, Taenzer P, Squire P, Dion D, Cholkani A, Gilani A, Gordon A, Henry J, Jovey R, Lynch M, Mailis-Gagnon A, Panju A, Rollman GB, Velly A; Canadian Pain Society. Pharmacological management of chronic neuropathic pain—consensus statement and guidelines from the Canadian Pain Society. *Pain Res Manag* 2007;12(1):13-21.
http://www.ncbi.nlm.nih.gov/pubmed/17372630?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
37. Fredheim OM, Borchgrevink PC, Klepstad P, Kaasa S, Dale O. Long term methadone for chronic pain: a pilot study of pharmacokinetic aspects. *Eur J Pain* 2007;11(6):599-604.
http://www.ncbi.nlm.nih.gov/pubmed/17113329?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

38. Staahl C, Dimcevski G, Andersen SD, Thorsgaard N, Christrup LL, Arendt-Nielsen L, Drewes AM. Differential effect of opioids in patients with chronic pancreatitis: an experimental pain study. *Scand J Gastroenterol* 2007;42(3):383-390.
http://www.ncbi.nlm.nih.gov/pubmed/17354119?ordinalpos=5&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
39. Nielsen CK, Ross FB, Lotfipour S, Saini KS, Edwards SR, Smith MT. Oxycodone and morphine have distinctly different pharmacological profiles: radioligand binding and behavioural studies in two rat models of neuropathic pain. *Pain* 2007;132(3):289-300.
http://www.ncbi.nlm.nih.gov/pubmed/17467904?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
40. Watson CP, Moulin D, Watt-Watson J, Gordon A, Eisenhoffer J. Controlled-release oxycodone relieves neuropathic pain: a randomized controlled trial in painful diabetic neuropathy. *Pain* 2003;105(1-2):71-78.
http://www.ncbi.nlm.nih.gov/pubmed/14499422?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
41. Mikus G, Weiss J. Influence of CYP2D6 genetics on opioid kinetics, metabolism and response. *Curr Pharmacogenomics* 2005;3(1):43-52.
42. Sagata K, Minami K, Yanagihara N, Shiraishi M, Toyohira Y, Ueno S, Shigematsu A. Tramadol inhibits norepinephrine transporter function at desipramine-binding sites in cultured bovine adrenal medullary cells. *Anaesth Analg* 2002;94(4):901-906.
http://www.ncbi.nlm.nih.gov/pubmed/11916794?ordinalpos=3&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
43. Desmeules JA, Pigué V, Collart L, Dayer P. Contribution of monoaminergic modulation to the analgesic effect of tramadol. *Br J Clin Pharmacol* 1996;41(1):7-12.
http://www.ncbi.nlm.nih.gov/pubmed/8824687?ordinalpos=2&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum
44. Hollingshead J, Dühmke RM, Cornblath DR. Tramadol for neuropathic pain. *Cochrane Database Syst Rev* 2006;(3):CD003726.
http://www.ncbi.nlm.nih.gov/pubmed/16856016?ordinalpos=1&itool=EntrezSystem2.PEntrez.Pubmed.Pubmed_ResultsPanel.Pubmed_RVDocSum

8.5 Nerve blocks

Neural blockade for pain management is usually carried out by a consultant in pain medicine with an anaesthetic background. Textbooks have been written on the techniques employed and individual specialists using neural blockade must be well versed in assessment of the patient, the indications for specific procedures, and the general and specific risks associated with the procedures, as well as possible advantages.

Procedures may be performed for diagnostic reasons, therapeutic benefit or possibly both. Diagnostic blocks can be difficult to interpret and a clear understanding of the many mechanisms by which a block may acting must be understood. Temporary but consistent responses to nerve blocks may lead a specialist to proceed with a neurolytic nerve block or to a pulsed radiofrequency neuromodulation procedure. Neurolytic nerve blocks are rarely indicated for a benign process, and to proceed with a neurolytic nerve block may result in disastrous results.

Published guidelines emphasize that all nerve blocks should be performed with appropriate attention to safety, including the presence of skilled support staff and appropriate monitoring and resuscitation equipment. The use of block needles, nerve location devices and imaging (i.e. X-ray image intensifier, ultrasound or computerized tomography) appropriate for the procedure is essential.

The evidence base for nerve blocks is not strong (1-5), but suggests that:

- Peripheral nerve blocks, such as ilioinguinal/iliohypogastric/genitofemoral, may be useful in the management of neuropathic pain associated with nerve damage, such as following hernia repairs.
- Blocks around the spermatic cord may be useful diagnostically prior to testicular denervation.
- Lumbar (L1) sympathetic blocks may be helpful in the management of testicular pain, renal pain and possibly a range of pelvic pain conditions with afferents that pass via the L1 level.
- Pudendal nerve blocks may be useful in the management of pudendal nerve injury related pain and possibly pelvic floor muscle spasm. Where pudendal neuralgia is suspected, pudendal nerve blocks may have a diagnostic role. Multiple other nerves close to the pudendal nerve may also be associated with neuropathic symptoms and differential nerve blocks using neurotracing may be of help in understanding the process'.
- Pre-sacral blocks and the ganglion impar block may have a role in the management of pelvic pathology, particularly cancer pain.
- Sacral root nerve blocks may be helpful in the diagnosis of those conditions that might respond to

sacral root stimulation.

The above list is not exhaustive and readers are advised to look at the major textbooks in this area (6).

8.5.1 References

1. Kennedy EM, Harms BA, Starling JR. Absence of maladaptive neuronal plasticity after genitofemoralilioinguinal neurectomy. *Surgery* 1994;116(4):665-670; discussion 670-671.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7940164&dopt=Abstract
2. Yamamoto M, Hibi H, Katsuno S, Miyake K. Management of chronic orchialgia of unknown etiology. *Int J Urol* 1995;2(1):47-49.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7542163&dopt=Abstract
3. Calvillo O, Skaribas IM, Rockett C. Computed tomography-guided pudendal nerve block. A new diagnostic approach to long-term anoperineal pain: a report of two cases. *Reg Anaesth Pain Med* 2000;25(4):420-423.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10925942&dopt=Abstract
4. Kovacs P, Gruber H, Piegger J, Bodner G. New, simple, ultrasound-guided infiltration of the pudendal nerve: ultrasonographic technique. *Dis Colon Rectum* 2001;44(9):1381-1385.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11584221&dopt=Abstract
5. McDonald JS, Spigos DG. Computed tomography-guided pudendal block for treatment of pelvic pain due to pudendal neuropathy. *Obstet Gynecol* 2000;95(2):306-309.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10674599&dopt=Abstract
6. Baranowski AP, Fall M, Abrams P, eds. *Urogenital Pain in Clinical Practice*. Taylor and Francis, 2007.

8.6 Transcutaneous electrical nerve stimulation (TENS)

The rationale behind using surface electrical nerve stimulation to relieve pain is the stimulation of myelinated afferents and thus activation of segmental inhibitory circuits. Urinary frequency may also be reduced. The favoured explanation of transcutaneous electrical nerve stimulation (TENS) draws on the gate-control theory of pain (1). Nevertheless, TENS may directly elicit reflex effects and influence autonomous functions. For example, relaxation of the bronchial muscles (2), the coronary arteries (3) and the urinary bladder have been observed in response to TENS (4).

TENS involves the use of a pulse generator with an amplifier and electrodes. The pulses may be delivered continuously or as trains of varying duration. Continuous stimulation seems to be preferable when treating pain.

The stimulation pulses used may have different properties. Square-wave pulses are notably effective in activating the nerve fibres and are most frequently used. Biphasic pulses are preferable as the zero net charge flow of this pulse helps to reduce electrochemical reactions at the electrode contact sites. Nevertheless, technical simplification has led to the use of unipolar rectangular pulses in many devices, apparently with few complications. The stimulus intensity required to activate a peripheral nerve varies with the pulse duration. In terms of charge transfer for a threshold effect, short pulses (0.1 ms) are most effective, but at the expense of higher pulse amplitudes (5). For most applications of nerve stimulation, the pulse frequency is a crucial variable. The frequencies used during TENS vary widely, from 1 Hz to 100 Hz. There are no systematic evaluation data to guide optimal electrical settings for TENS in urological practice.

Standard electrodes are made of carbon rubber. These are strong, flexible, durable and cheap, but must be attached by adhesive tape. Self-adhesive electrodes have been developed. These are especially advantageous for people with sensitive skin, but they are expensive. The size of the electrode has a bearing on the current density – a minimum of 4 cm² has been recommended for TENS (6). The electrode-skin impedance should be reduced by application of a generous layer of electrolyte gel to promote good contact under the entire electrode.

The stimulus intensity required to elicit sensory appreciation varies between individuals. The maximum tolerable intensity just below pain threshold should be used. While it is plausible that electrode positioning will affect the result of treatment, this property has not been evaluated. In BPS/IC, suprapubic (7,8), vaginal-anal (4,9) and tibial nerve sites (10,11) have been tested, all with some success.

Counselling of the patient before the start of the treatment is necessary. A specially trained nurse with the time necessary to communicate the technical instructions is a good option. The patient should be confident with the feeling of strong stimulation and view self-treatment without fear. The induction time for TENS to

produce analgesia varies widely. The effect is cumulative. Since onset and progression are usually rather slow in IC, the standard recommendation so far has been 0.5-2 hours of treatment twice daily. The duration of an individual treatment session depends on the severity of pain.

8.6.1 Results of suprapubic TENS in BPS/IC

Observations are scant. In the largest study published so far, 60 patients, 33 with classic IC and 27 with non-ulcer disease, were treated by suprapubic TENS (6). The electrodes were positioned about 10-15 cm apart, immediately above the pubic symphysis. They were attached by a long strip of adhesive tape going halfway around the body to enable the patient to be ambulant during stimulation. Follow-up ranged from 9 months to 17 years. Patients who responded reported more marked effects on bladder pain than on micturition frequency. Of the patients with classic IC, 54% were helped by the treatment. The outcome of TENS was less favourable in non-ulcer IC. Of 27 patients with non-ulcer IC, only 26% benefited from the treatment.

The present experience of electrical stimulation is based on open studies. There are difficulties in designing controlled studies of TENS, since the treatment is based on administration of stimulation of high intensity, at specific sites, over a very long period of time. Another problem is that it is not possible to measure pain precisely. Therefore, it is difficult to assess the efficacy of TENS in BPS/IC with accuracy.

Several controlled studies of post-operative pain have shown TENS to be superior to sham stimulation (12). TENS has also been shown to reduce the amount of halothane required to maintain adequate anaesthesia during hand surgery in unconscious patients, in whom psychological influences have been eliminated (13). The beneficial effect of TENS on classic IC clearly exceeds the level of the placebo effect observed in drug studies of IC (54% versus 13-20%) (14,15).

8.7 Sacral neuromodulation in pelvic pain syndromes

Sacral neuromodulation (SNS) has been shown to have benefits in patients with refractory motor urge incontinence (16,17), urinary retention, and CPP (18-20). Neuropathic pain and complex regional pain syndromes may also be treated successfully with neurostimulation applied to dorsal columns and peripheral nerves (21).

The mechanisms of action are the subject of many hypotheses, with very little evidence to support any particular one. The hypotheses include:

- blocking of pain transmission by direct effects in the spinothalamic tracts
- activation of descending inhibitory pathways
- effects on central sympathetic systems
- segmental inhibition through coarse fibre activation and brain stem loops
- inhibition by increasing gamma-aminobutyric acid levels in the dorsal horn
- thalamocortical mechanisms masking the nociceptive input (21,22).

Sacral root neuromodulation was introduced in the mid-1980s as a means of regaining bladder control in the face of disturbed function (23). Based on the neurophysiology of the bladder and urethra, it is a minimally invasive tool that bridges the gap between conservative options and invasive surgical procedures. The data on clinical applications are drawn exclusively from observational studies.

Sacral root neuromodulation draws on the observation that electrical stimulation of sacral nerves modulates neural reflexes of the pelvis (24). Acceptable application of the stimuli is the challenge. Neurostimulation of S3 or S4 sacral nerves using a transforaminal approach is a viable option for patients with refractory urinary voiding disorders.

Recently, SNS has also been investigated in IC. In an initial report on six patients (25), percutaneous neurostimulation significantly improved frequency, pain and urgency towards normal values, while urinary markers for IC were normalized. Maher et al. (26) reported a favourable response with significant improvement in pelvic pain, daytime frequency, nocturia, urgency and voided volume in 15 women with IC.

Because pelvic pain syndromes are viewed as a manifestation of disturbed neural function, patients with refractory pelvic floor dysfunction and pelvic pain have been treated with SNS and benefit has been reported (27). Sacral neuromodulation for CPP has been based on promising data from pilot studies and prospective, placebo-controlled studies are now justified.

8.7.1 References

1. Melzack R, Wall PD. Pain mechanisms: a new theory. *Science* 1965;150(699):971-979.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=5320816&dopt=Abstract
2. Sovijarvi AR, Poppius H. Acute bronchodilating effect of transcutaneous nerve stimulation in asthma. A peripheral reflex or psychogenic response. *Scand J Respir Dis* 1977;58(3):164-169.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=302028&dopt=Abstract

3. Mannheimer C, Carlsson CA, Vedin A, Wilhelmsson C. Transcutaneous electrical nerve stimulation (TENS) in angina pectoris. *Pain* 1986;26(3):291-300.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3534690&dopt=Abstract
4. Fall M, Carlsson CA, Erlandson BE. Electrical stimulation in interstitial cystitis. *J Urol* 1980;123(2):192-195.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6965508&dopt=Abstract
5. Fall M, Lindström S. Electrical stimulation. A physiologic approach to the treatment of urinary incontinence. *Urol Clin North Am* 1991;18(2):393-407.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2017820&dopt=Abstract
6. Fall M, Lindstrom S. Transcutaneous electrical nerve stimulation in classic and nonulcer interstitial cystitis. *Urol Clin North Am* 1994;21(1):131-139.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8284836&dopt=Abstract
7. Fall M. Conservative management of chronic interstitial cystitis: transcutaneous electrical nerve stimulation and transurethral resection. *J Urol* 1985;133(5):774-778.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3872946&dopt=Abstract
8. Fall M. Transcutaneous electrical nerve stimulation in interstitial cystitis. Update on clinical experience. *Urology* 1987;29(4 Suppl):40-42.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=3494331&dopt=Abstract
9. Eriksen BC. Painful bladder disease in women: effect of maximal electric pelvic floor stimulation. *Neurourol Urodynam* 1989;8:362-363.
<http://www3.interscience.wiley.com/cgi-bin/jhome/35693>
10. Geirsson G, Wang YH, Lindstrom S, Fall M. Traditional acupuncture and electrical stimulation of the posterior tibial nerve. A trial in chronic interstitial cystitis. *Scand J Urol Nephrol* 1993;27(1):67-70.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=8493470&dopt=Abstract
11. McGuire EJ, Zhang SC, Horwinski ER, Lytton B. Treatment of motor and sensory detrusor instability by electrical stimulation. *J Urol* 1983;129(1):78-79.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6600794&dopt=Abstract
12. Woolf CJ. Segmental afferent fibre-induced analgesia: transcutaneous electrical nerve stimulation (TENS) and vibration. In: Melzack R, Wall PD, eds. *Textbook of Pain*. 2nd ed. Edinburgh; Churchill-Livingstone: 884-896.
13. Bourke DL, Smith BA, Erickson J, Gwartz B, Lessard L. TENS reduces halothane requirements during hand surgery. *Anaesthesiology* 1984;61(6):769-772.
http://www.ncbi.nlm.nih.gov/80/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=6391280&dopt=Abstract
14. Mulholland SG, Hanno P, Parsons CL, Sant GR, Staskin DR. Pentosan polysulfate sodium for therapy of interstitial cystitis. A double-blind placebo-controlled clinical study. *Urology* 1990;35(6):552-558.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=1693797&dopt=Abstract
15. Holm-Bentzen M, Jacobsen F, Nerstrom B, Lose G, Kristensen JK, Pedersen RH, Krarup T, Feggetter J, Bates P, Barnard R et al. A prospective double-blind clinically controlled multicenter trial of sodium pentosanpolysulfate in the treatment of interstitial cystitis and related painful bladder disease. *J Urol* 1987;138(3):503-507.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2442415&dopt=Abstract
16. Kennedy EM, Harms BA, Starling JR. Absence of maladaptive neuronal plasticity after genitofemoralilioinguinal neurectomy. *Surgery* 1994;116(4):665-70; discussion 670-671.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=7940164&dopt=Abstract

17. Janknegt RA, Hassouna MM, Siegel SW, Schmidt RA, Gajewski JB, Rivas DA, Elhilali MM, Milam DC, van Kerrebroeck PE, Dijkema HE, Lycklama a Nyeholt AA, Fall M, Jonas U, Catanzaro F, Fowler CJ, Oleson KA. Long-term effectiveness of sacral nerve stimulation for refractory urge incontinence. *Eur Urol* 2001;39(1):101-106.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11173947&dopt=Abstract
18. Paszkiewicz EJ, Siegel SW, Kirkpatrick C, Hinkel B, Keeisha J, Kirkemo A. Sacral nerve stimulation in patients with chronic, intractable pelvic pain. *Urology* 2001;57(6 Suppl 1):124.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11378113&dopt=Abstract
19. Edlund C, Hellstrom M, Pecker R, Fall M. First Scandinavian experience of electrical sacral nerve stimulation in the treatment of the overactive bladder. *Scand J Urol Nephrol* 2000;34(6):366-376.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11195901&dopt=Abstract
20. Shaker HS, Hassouna M. Sacral root neuromodulation in idiopathic nonobstructive chronic urinary retention. *J Urol* 1998;159(5):1476-1478.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=9554336&dopt=Abstract
21. Kemler MA, Barendse GA, van Kleef M, de Vet HC, Rijk CP, Furnee CA, van den Wildenberg FA. Spinal cord stimulation in patients with chronic reflex sympathetic dystrophy. *N Engl J Med* 2000;343(9):618-624.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10965008&dopt=Abstract
22. Kemler MA, Barendse GA, van Kleef M, Egbrink MG. Pain relief in complex regional pain syndrome due to spinal cord stimulation does not depend on vasodilation. *Anaesthesiology* 2000;92(6):1653-1660.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10839916&dopt=Abstract&itool=iconabstr
23. Schmidt RA. Applications of neuromodulation. *Urol Neuro Urodyn* 1988;7:585.
24. Schmidt RA, Senn E, Tanagho EA. Functional evaluation of sacral nerve root integrity. Report of a technique. *Urology* 1990;35(5):388-392.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=2336766&dopt=Abstract
25. Chai TC, Zhang C, Warren JW, Keay S. Percutaneous sacral third nerve root neurostimulation improves symptoms and normalizes urinary HB-EGF levels and antiproliferative activity in patients with interstitial cystitis. *Urology* 2000;55(5):643-646.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=10792070&dopt=Abstract
26. Maher CF, Carey MP, Dwyer PL, Schluter PL. Percutaneous sacral nerve root neuromodulation for intractable interstitial cystitis. *J Urol* 2001;165(3):884-886.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=11176493&dopt=Abstract
27. Aboseif S, Tamaddon K, Chalfin S, Freedman S, Kaptein J. Sacral neuromodulation as an effective treatment for refractory pelvic floor dysfunction. *Urology* 2002;60(1):52-56.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&list_uids=12100921&dopt=Abstract

9. ABBREVIATIONS USED IN THE TEXT

This list is not comprehensive for the most common abbreviations

ABP	acute bacterial prostatitis
BCG	Bacillus Calmette-Guérin
BPS	bladder pain syndrome
BTX	Botulinum Toxin A
CBP	chronic bacterial prostatitis
CFU	colony-forming units
CNS	central nervous system
COX	cyclo-oxygenase
CPP(S)	chronic pelvic pain (syndrome)
CPSI	chronic Prostatitis Symptom Index
CyA	cyclosporin A
DMSO	dimethyl sulphoxide
DRE	digital rectal examination
ED	erectile dysfunction
EMDA	electromotive drug administration
EPS	expressed prostatic secretions
ESSIC	European Society for the Study of IC/PBS
FD	first desire to void
GAG	glycosaminoglycan
GI	gastrointestinal
HBO	hyperbaric oxygen
IASP	International Association for the Study of Pain
IC	interstitial cystitis
ICA	Interstitial Cystitis Association
ICDB	Interstitial Cystitis Data Base
ICS	International Continence Society
ICSI	Interstitial Cystitis Symptom Index (also known as the O'Leary Sant Symptom Index)
IL-6	interleukin-6
I-PSS	International Prostate Symptom Score
ISSVD	International Society for the Study of Vulvovaginal Disease
LHRH	luteinizing hormone releasing hormone
LUT	lower urinary tract
LUTS	lower urinary tract symptoms
ND	normal desire to void
Nd-YAG	neodymium-yttrium-aluminium-garnet
NIDDK	National Institute of Diabetes and Digestive and Kidney Diseases
NIH	National Institutes of Health
NMDA	N-methyl-D-aspartate
NSAID	non-steroidal anti-inflammatory drugs
PBS	painful bladder syndrome
PCA	prostate cancers
PFM	pelvic floor muscles
PPMT	pre- and post-massage test
PPS	pentosanpolysulphate sodium
PSA	prostate specific antigen
PUGO	IASP special interest group, Pain of Urogenital Origin
PVR	post-void residual urine
RCT	randomized controlled trial
RTX	resiniferatoxin
SNS	sacral neuromodulation
SPIN	Specialists in Pain International Network
TENS	transcutaneous electrical nerve stimulation
TUNA	transurethral needle ablation of the prostate
TUR	transurethral resection
VAS	visual analogue scale
VB2	pre-prostatic massage urine
VB3	post-prostatic massage urine

VIP	vasoactive intestinal peptide
WBC	white blood cells
WHO	World Health Organization

